



# wwPDB NMR Structure Validation Summary Report ⓘ

Apr 26, 2016 – 02:29 PM BST

PDB ID : 1EIH  
Title : NMR STRUCTURE OF HOLO CELLULAR RETINOL-BINDING PROTEIN II  
Authors : Lu, J.; Lin, C.L.; Tang, C.; Ponder, J.W.; Kao, J.L.; Cistola, D.P.; Li, E.  
Deposited on : 2000-02-25

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*


The overall completeness of chemical shifts assignment is 88%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	134	

## 2 Ensemble composition and analysis ⓘ

This entry contains 25 models. Model 2 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:134 (132)	0.49	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 3, 5, 6, 7, 8, 9, 10, 14, 16, 18, 21, 23
2	1, 4, 11, 12, 13, 15, 17, 19, 20, 22, 24
Single-model clusters	25


### 3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2219 atoms, of which 1103 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called CELLULAR RETINOL-BINDING PROTEIN II.

Mol	Chain	Residues	Atoms						Trace
1	A	134	Total	C	H	N	O	S	0
			2168	686	1073	189	214	6	

- Molecule 2 is RETINOL (three-letter code: RTL) (formula:  $C_{20}H_{30}O$ ).




Mol	Chain	Residues	Atoms			
2	A	1	Total	C	H	O
			51	20	30	1

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: CELLULAR RETINOL-BINDING PROTEIN II



### 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 2. Colouring as in section 4.1 above.

- Molecule 1: CELLULAR RETINOL-BINDING PROTEIN II



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry & simulated annealing*.

Of the 30 calculated structures, 25 were deposited, based on the following criterion: *FINAL PENALTY FUNCTION VALUES WITHIN 2 STANDARD DEVIATIONS FROM THE MEAN*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
TINKER	structure solution	3.3
TINKER	refinement	3.3

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 4682
Number of chemical shift lists	3
Total number of shifts	2344
Number of shifts mapped to atoms	2319
Number of unparsed shifts	0
Number of shifts with mapping errors	25
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	88%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RTL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.06±0.02	6±1/1098 (0.6±0.1%)	2.16±0.05	40±7/1474 (2.7±0.5%)
All	All	1.06	160/27450 (0.6%)	2.16	1004/36850 (2.7%)

5 of 8 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	110	TRP	CD2-CE2	-6.90	1.33	1.41	22	23
1	A	9	TRP	CD2-CE2	-6.77	1.33	1.41	15	23
1	A	107	TRP	CD2-CE2	-6.69	1.33	1.41	24	17
1	A	89	TRP	CG-CD2	-6.58	1.32	1.43	8	25
1	A	110	TRP	CG-CD2	-6.58	1.32	1.43	7	25

5 of 176 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	36	ARG	NE-CZ-NH1	16.60	128.60	120.30	16	11
1	A	89	TRP	CD1-CG-CD2	15.84	118.97	106.30	11	23
1	A	105	ARG	NE-CZ-NH1	15.77	128.19	120.30	19	13
1	A	89	TRP	CG-CD1-NE1	-13.30	96.80	110.10	12	25
1	A	110	TRP	CD1-CG-CD2	13.03	116.72	106.30	2	22

There are no chirality outliers.

There are no planarity outliers.

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1080	1055	1054	46±8
2	A	21	30	30	5±3
All	All	27525	27125	27100	1199

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

5 of 590 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:17:PHE:CD2	1:A:35:VAL:HG22	0.93	1.99	25	5
1:A:26:ILE:HG21	1:A:30:THR:HG21	0.90	1.40	9	2
1:A:109:GLN:CG	1:A:118:LEU:HD13	0.89	1.96	20	7
1:A:109:GLN:HG2	1:A:118:LEU:HD13	0.89	1.44	20	7
1:A:37:LEU:HD22	1:A:38:THR:N	0.83	1.88	2	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/134 (98%)	86±4 (66±3%)	31±4 (24±3%)	14±3 (11±2%)	1	9
All	All	3275/3350 (98%)	2157 (66%)	771 (24%)	347 (11%)	1	9

5 of 64 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	6	ASN	22
1	A	44	VAL	20

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	94	LEU	16
1	A	115	LYS	15
1	A	104	ASN	14

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	118/120 (98%)	70±4 (60±3%)	48±4 (40±3%)	<div>05</div>
All	All	2950/3000 (98%)	1760 (60%)	1190 (40%)	<div>05</div>

5 of 109 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	31	ARG	25
1	A	9	TRP	25
1	A	53	LYS	25
1	A	107	TRP	25
1	A	63	LEU	25

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	RTL	A	135	-	21,21,21	1.76±0.15	1±0 (4±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	RTL	A	135	-	26,28,28	2.39±0.14	2±1 (6±3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RTL	A	135	-	-	0±0,14,31,31	0±0,1,1,1

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	135	RTL	C1-C6	8.45	1.42	1.53	10	25

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )	Models	
								Worst	Total
2	A	135	RTL	C4-C5-C6	7.03	115.00	122.73	14	10
2	A	135	RTL	C11-C10-C9	6.36	117.98	127.22	1	12
2	A	135	RTL	C17-C1-C16	5.88	89.42	108.36	16	21
2	A	135	RTL	C2-C1-C6	5.29	118.35	110.48	19	2

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation ⓘ

The completeness of assignment taking into account all chemical shift lists is 88% for the well-defined parts and 87% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 4682

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping ⓘ

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	25
Number of shifts mapped to atoms	0
Number of unparsed shifts	0
Number of shifts with mapping errors	25
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Chain not found in structure. First 5 (of 25) occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
UNMAPPED	1	RTL	H151	3.56	0.025	1
UNMAPPED	1	RTL	H20	0.4	0.025	1
UNMAPPED	1	RTL	H31	1.85	0.025	1
UNMAPPED	1	RTL	H8	6.32	0.025	1
UNMAPPED	1	RTL	C16	29.1	0.15	1

#### 7.1.2 Chemical shift referencing ⓘ

No chemical shift referencing corrections were calculated (not enough data).

### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 1677. 0 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	0/660 (0%)	0/264 (0%)	0/264 (0%)	0/132 (0%)
Sidechain	0/874 (0%)	0/507 (0%)	0/318 (0%)	0/49 (0%)
Aromatic	0/143 (0%)	0/75 (0%)	0/62 (0%)	0/6 (0%)
Overall	0/1677 (0%)	0/846 (0%)	0/644 (0%)	0/187 (0%)

### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

### 7.1.5 Random Coil Index (RCI) plots [i](#)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned\_chem\_shift\_list\_1). RCI is only applicable to proteins.

## 7.2 Chemical shift list 2

File name: BMRB entry 4682

Chemical shift list name: *assigned\_chem\_shift\_list\_2*

### 7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1634
Number of shifts mapped to atoms	1634
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	11

### 7.2.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, <i>ppm</i>	Suggested action
$^{13}\text{C}_\alpha$	133	$-0.34 \pm 0.09$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	122	$-0.22 \pm 0.14$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	133	$1.20 \pm 0.16$	Should be applied
$^{15}\text{N}$	132	$-4.71 \pm 0.23$	Should be applied

### 7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 87%, i.e. 1460 atoms were assigned a chemical shift out of a possible 1677. 20 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	660/660 (100%)	264/264 (100%)	264/264 (100%)	132/132 (100%)
Sidechain	729/874 (83%)	457/507 (90%)	257/318 (81%)	15/49 (31%)
Aromatic	71/143 (50%)	67/75 (89%)	0/62 (0%)	4/6 (67%)
Overall	1460/1677 (87%)	788/846 (93%)	521/644 (81%)	151/187 (81%)

### 7.2.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.


Mol	Chain	Res	Type	Atom	Shift, <i>ppm</i>	Expected range, <i>ppm</i>	Z-score
1	A	117	TYR	HB3	0.45	4.75 – 0.95	-6.3
1	A	102	LYS	HG2	3.01	2.67 – 0.07	6.3
1	A	116	LEU	HB3	-0.70	3.34 – -0.26	-6.2
1	A	96	CYS	HB3	0.06	5.25 – 0.55	-6.0
1	A	96	CYS	HB2	0.32	5.20 – 0.70	-5.8
1	A	3	LYS	HE3	1.81	3.86 – 1.96	-5.8
1	A	39	GLN	HE22	4.43	9.27 – 4.77	-5.8
1	A	85	THR	HB	2.32	5.82 – 2.52	-5.6
1	A	102	LYS	HG3	2.92	2.76 – -0.04	5.6
1	A	41	LYS	HE3	1.88	3.86 – 1.96	-5.4
1	A	125	GLN	NE2	121.00	120.91 – 102.81	5.0

### 7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication

of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



## 7.3 Chemical shift list 3

File name: BMRB entry 4682

Chemical shift list name: *assigned\_chem\_shift\_list\_3*

### 7.3.1 Bookkeeping ⓘ

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	685
Number of shifts mapped to atoms	685
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	10

### 7.3.2 Chemical shift referencing ⓘ

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	0	—	—
$^{13}\text{C}_\beta$	0	—	—
$^{13}\text{C}'$	0	—	—
$^{15}\text{N}$	132	$-4.79 \pm 0.46$	Should be applied

### 7.3.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 37%, i.e. 626 atoms were assigned a chemical shift out of a possible 1677. 0 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	389/660 (59%)	257/264 (97%)	0/264 (0%)	132/132 (100%)
Sidechain	229/874 (26%)	212/507 (42%)	0/318 (0%)	17/49 (35%)
Aromatic	8/143 (6%)	4/75 (5%)	0/62 (0%)	4/6 (67%)
Overall	626/1677 (37%)	473/846 (56%)	0/644 (0%)	153/187 (82%)

### 7.3.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	59	ARG	NE	125.80	92.63 – 76.73	25.9
1	A	128	ARG	NE	125.50	92.63 – 76.73	25.7
1	A	117	TYR	HB3	0.38	4.75 – 0.95	-6.5
1	A	92	ASN	HD22	4.07	9.59 – 4.69	-6.3
1	A	96	CYS	HB3	0.01	5.25 – 0.55	-6.1
1	A	109	GLN	HE22	4.30	9.27 – 4.77	-6.0
1	A	96	CYS	HB2	0.29	5.20 – 0.70	-5.9
1	A	85	THR	HB	2.26	5.82 – 2.52	-5.8
1	A	39	GLN	HE22	4.46	9.27 – 4.77	-5.7
1	A	45	GLN	HB2	0.74	3.30 – 0.80	-5.2

### 7.3.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



