



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:27 AM GMT

PDB ID : 3EOA
Title : Crystal structure the Fab fragment of Efalizumab in complex with LFA-1 I domain, Form I
Authors : Li, S.; Ding, J.
Deposited on : 2008-09-26
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	<div> <div>%</div> <div> <div></div> <div>72%</div> <div>25%</div> <div>.</div> </div> </div>
1	L	214	<div> <div>3%</div> <div> <div></div> <div>71%</div> <div>23%</div> <div>6%</div> </div> </div>
2	B	220	<div> <div>%</div> <div> <div></div> <div>81%</div> <div>15%</div> <div>..</div> </div> </div>
2	H	220	<div> <div>%</div> <div> <div></div> <div>78%</div> <div>18%</div> <div>..</div> </div> </div>
3	I	181	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>20%</div> <div>..</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	J	181	<div><div></div><div>3%</div><div>78%</div><div>17%</div><div><div></div><div></div><div></div><div></div><div></div></div></div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9804 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Efalizumab Fab fragment, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	214	Total	C	N	O	S	0	0	0
			1646	1030	276	334	6			
1	A	214	Total	C	N	O	S	0	0	0
			1646	1030	276	334	6			

- Molecule 2 is a protein called Efalizumab Fab fragment, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	215	Total	C	N	O	S	0	0	0
			1640	1043	273	317	7			
2	B	215	Total	C	N	O	S	0	0	0
			1640	1043	273	317	7			

- Molecule 3 is a protein called Integrin alpha-L.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	I	179	Total	C	N	O	S	0	0	0
			1438	929	231	274	4			
3	J	179	Total	C	N	O	S	0	0	0
			1438	929	231	274	4			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	80	Total	O	0	0
			80	80		
4	B	64	Total	O	0	0
			64	64		
4	H	66	Total	O	0	0
			66	66		
4	I	60	Total	O	0	0
			60	60		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	J	50	Total	O	0	0
			50	50		
4	L	36	Total	O	0	0
			36	36		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Efalizumab Fab fragment, light chain




- Molecule 1: Efalizumab Fab fragment, light chain




- Molecule 2: Efalizumab Fab fragment, heavy chain




- Molecule 2: Efalizumab Fab fragment, heavy chain







• Molecule 3: Integrin alpha-L



• Molecule 3: Integrin alpha-L



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.72Å 81.70Å 281.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 49.93 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.0 (50.00-2.80) 98.1 (49.93-2.80)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.09 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.225 , 0.264 0.224 , 0.233	Depositor DCC
R_{free} test set	1856 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	36.6	Xtriage
Anisotropy	0.487	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	1 of 37157 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	9804	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.04 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.1410e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	1/1681 (0.1%)	0.52	0/2280
1	L	0.54	3/1681 (0.2%)	0.56	1/2280 (0.0%)
2	B	0.37	0/1683	0.54	0/2291
2	H	0.37	0/1683	0.53	0/2291
3	I	0.46	0/1465	0.57	1/1970 (0.1%)
3	J	0.52	1/1465 (0.1%)	0.58	1/1970 (0.1%)
All	All	0.45	5/9658 (0.1%)	0.55	3/13082 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	J	302	LEU	N-CA	9.21	1.64	1.46
1	L	121	SER	CB-OG	7.37	1.51	1.42
1	L	50	SER	CB-OG	5.34	1.49	1.42
1	L	169	LYS	CE-NZ	5.15	1.61	1.49
1	A	143	GLU	CG-CD	5.04	1.59	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	262	GLY	N-CA-C	6.38	129.04	113.10
3	J	302	LEU	N-CA-C	5.34	125.41	111.00
1	L	171	SER	N-CA-C	5.26	125.19	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1646	0	1608	32	0
1	L	1646	0	1608	32	0
2	B	1640	0	1591	20	0
2	H	1640	0	1591	29	0
3	I	1438	0	1449	19	0
3	J	1438	0	1449	21	0
4	A	80	0	0	0	0
4	B	64	0	0	0	0
4	H	66	0	0	0	0
4	I	60	0	0	3	0
4	J	50	0	0	0	0
4	L	36	0	0	0	0
All	All	9804	0	9296	145	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 145 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:J:129:ASN:HB3	3:J:165:SER:HB2	1.53	0.89
2:B:164:SER:H	2:B:205:ASN:HD21	1.23	0.87
2:H:83:MET:HB3	2:H:86:LEU:HD21	1.57	0.85
1:A:2:ILE:HG22	1:A:4:MET:HE3	1.58	0.85
3:I:287:LYS:HE3	4:I:366:HOH:O	1.75	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/214 (99%)	201 (95%)	10 (5%)	1 (0%)	34	69
1	L	212/214 (99%)	195 (92%)	12 (6%)	5 (2%)	7	25
2	B	211/220 (96%)	201 (95%)	9 (4%)	1 (0%)	34	69
2	H	211/220 (96%)	204 (97%)	7 (3%)	0	100	100
3	I	177/181 (98%)	165 (93%)	8 (4%)	4 (2%)	8	26
3	J	177/181 (98%)	163 (92%)	12 (7%)	2 (1%)	17	50
All	All	1200/1230 (98%)	1129 (94%)	58 (5%)	13 (1%)	17	50

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	84	ALA
1	L	170	ASP
1	L	213	GLU
2	B	113	GLN
3	I	262	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/189 (100%)	171 (90%)	18 (10%)	11	30
1	L	189/189 (100%)	167 (88%)	22 (12%)	7	20
2	B	182/186 (98%)	174 (96%)	8 (4%)	35	69
2	H	182/186 (98%)	172 (94%)	10 (6%)	27	59
3	I	159/161 (99%)	141 (89%)	18 (11%)	7	22
3	J	159/161 (99%)	144 (91%)	15 (9%)	11	31
All	All	1060/1072 (99%)	969 (91%)	91 (9%)	13	36

5 of 91 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	25	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	1	GLU
3	J	231	THR
2	H	51	ILE
2	H	124	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	3	GLN
2	H	179	GLN
3	J	247	ASN
2	H	39	GLN
2	H	113	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/214 (100%)	-0.11	2 (0%) 85 79	14, 29, 43, 66	0
1	L	214/214 (100%)	0.41	6 (2%) 56 44	22, 38, 53, 69	0
2	B	215/220 (97%)	0.00	3 (1%) 78 69	13, 26, 52, 66	0
2	H	215/220 (97%)	0.28	2 (0%) 85 79	20, 31, 45, 61	0
3	I	179/181 (98%)	-0.02	3 (1%) 73 63	13, 27, 45, 53	0
3	J	179/181 (98%)	0.09	5 (2%) 56 44	11, 27, 51, 61	0
All	All	1216/1230 (98%)	0.11	21 (1%) 73 63	11, 31, 48, 69	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	212	GLY	3.2
2	B	136	SER	3.1
1	A	212	GLY	2.7
3	I	163	ASN	2.6
3	J	163	ASN	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.