



wwPDB EM Map/Model Validation Report ⓘ

Apr 10, 2016 – 01:39 PM BST

PDB ID : 3EQ3
EMDB ID: : EMD-1565
Title : Model of tRNA(Trp)-EF-Tu in the ribosomal pre-accommodated state revealed by cryo-EM
Authors : Frank, J.; Li, W.; Agirrezabala, X.
Deposited on : 2008-09-30
Resolution : 9.00 Å(reported)
Based on PDB ID : 2AVY, 2AW4, 1QZA, 1OB2

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.
For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/EMValidationReportHelp>

MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : trunk27241

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY










The reported resolution of this entry is 9.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	114402	924
RNA backbone	3027	244

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	X	393	 100%
2	L	123	 100%
3	I	141	 100%
4	Y	74	 100%
5	A	9	 100%
6	C	11	 100%
7	B	48	 100%
8	D	28	 100%
9	E	17	 100%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 844 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Elongation factor Tu.

Mol	Chain	Residues	Atoms		AltConf	Trace
1	X	393	Total	C	0	393
			393	393		

- Molecule 2 is a protein called 30S ribosomal protein S12.

Mol	Chain	Residues	Atoms		AltConf	Trace
2	L	123	Total	C	0	123
			123	123		

- Molecule 3 is a protein called 50S ribosomal protein L11.

Mol	Chain	Residues	Atoms		AltConf	Trace
3	I	141	Total	C	0	141
			141	141		

- Molecule 4 is a RNA chain called tRNA.

Mol	Chain	Residues	Atoms		AltConf	Trace
4	Y	74	Total	P	0	74
			74	74		

- Molecule 5 is a RNA chain called Fragment h18 of the 16S rRNA.

Mol	Chain	Residues	Atoms		AltConf	Trace
5	A	9	Total	P	0	9
			9	9		

- Molecule 6 is a RNA chain called Fragment h44 of the 16S rRNA.

Mol	Chain	Residues	Atoms		AltConf	Trace
6	C	11	Total	P	0	11
			11	11		

- Molecule 7 is a RNA chain called Fragment H43-44 of the 23S rRNA.

Mol	Chain	Residues	Atoms		AltConf	Trace
7	B	48	Total	P	0	48
			48	48		

- Molecule 8 is a RNA chain called Fragment H95 of the 23S rRNA.

Mol	Chain	Residues	Atoms		AltConf	Trace
8	D	28	Total	P	0	28
			28	28		

- Molecule 9 is a RNA chain called Fragment H69 of the 23S rRNA.

Mol	Chain	Residues	Atoms		AltConf	Trace
9	E	17	Total	P	0	17
			17	17		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Elongation factor Tu

Chain X:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: 30S ribosomal protein S12

Chain L:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: 50S ribosomal protein L11

Chain I:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: tRNA

Chain Y:  100%

There are no outlier residues recorded for this chain.

- Molecule 5: Fragment h18 of the 16S rRNA

Chain A:  100%

There are no outlier residues recorded for this chain.

- Molecule 6: Fragment h44 of the 16S rRNA

Chain C:  100%

There are no outlier residues recorded for this chain.

- Molecule 7: Fragment H43-44 of the 23S rRNA

Chain B:  100%

There are no outlier residues recorded for this chain.

- Molecule 8: Fragment H95 of the 23S rRNA

Chain D:  100%

There are no outlier residues recorded for this chain.

- Molecule 9: Fragment H69 of the 23S rRNA

Chain E:  100%

There are no outlier residues recorded for this chain.

4 Experimental information

Property	Value	Source
Reconstruction method	Not provided	Depositor
Imposed symmetry	POINT, Not provided	Depositor
Number of images	Not provided	Depositor
Resolution determination method	Not provided	Depositor
CTF correction method	Not provided	Depositor
Microscope	Not provided	Depositor
Voltage (kV)	Not provided	Depositor
Electron dose ($e^-/\text{\AA}^2$)	Not provided	Depositor
Minimum defocus (nm)	Not provided	Depositor
Maximum defocus (nm)	Not provided	Depositor
Magnification	Not provided	Depositor
Image detector	AMT	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	393	0	0	0	0
2	L	123	0	0	0	0
3	I	141	0	0	0	0
4	Y	74	0	0	0	0
5	A	9	0	0	0	0
6	C	11	0	0	0	0
7	B	48	0	0	0	0
8	D	28	0	0	0	0
9	E	17	0	0	0	0
All	All	844	0	0	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	Y	0/74	-	-
5	A	0/9	-	-
6	C	0/11	-	-
7	B	0/48	-	-
8	D	0/28	-	-
9	E	0/17	-	-
All	All	0/187	-	-

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.