



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:30 AM GMT

PDB ID : 3EVU  
Title : Crystal structure of Calcium bound dimeric GCAMP2, (#1)  
Authors : Wang,Q.; Shui,B.; Kotlikoff,M.I.; Sondermann,H.  
Deposited on : 2008-10-13  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1609 (1.76-1.76)
Clashscore	102246	1730 (1.76-1.76)
Ramachandran outliers	100387	1711 (1.76-1.76)
Sidechain outliers	100360	1711 (1.76-1.76)
RSRZ outliers	91569	1610 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	449	<div> <div>3%</div> <div>79%</div> <div>8%</div> <div>12%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3682 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin light chain kinase, Green fluorescent protein, Calmodulin chimera.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	394	Total	C	N	O	S	0	8	0
			3162	1990	528	628	16			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	45	ASN	GLN	CONFLICT	UNP P11799
A	60	LEU	-	LINKER	UNP P11799
A	61	GLU	-	LINKER	UNP P11799
A	76	ALA	VAL	CONFLICT	UNP P42212
A	88	GLY	SER	CONFLICT	UNP P42212
A	93	TYR	ASP	CONFLICT	UNP P42212
A	119	LYS	ALA	CONFLICT	UNP P42212
A	144	LEU	HIS	CONFLICT	UNP P42212
A	152	GLY	-	LINKER	UNP P42212
A	153	GLY	-	LINKER	UNP P42212
A	154	THR	-	LINKER	UNP P42212
A	155	GLY	-	LINKER	UNP P42212
A	156	GLY	-	LINKER	UNP P42212
A	157	SER	-	LINKER	UNP P42212
A	158	MET	-	LINKER	UNP P42212
A	159	VAL	-	LINKER	UNP P42212
A	222	LEU	PHE	CONFLICT	UNP P42212
A	224	CRO	SER	SER REMARK 999	UNP P42212
A	224	CRO	TYR	SEE REMARK 999	UNP P42212
A	224	CRO	TYR	SEE REMARK 999	UNP P42212
A	251	ILE	VAL	CONFLICT	UNP P42212
A	303	THR	-	LINKER	UNP P42212
A	304	ARG	-	LINKER	UNP P62161

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total 4	Ca 4	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	516	Total 516	O 516	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Myosin light chain kinase, Green fluorescent protein, Calmodulin chimera



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	127.94Å 47.31Å 69.64Å 90.00° 100.70° 90.00°	Depositor
Resolution (Å)	42.52 – 1.75 42.52 – 1.75	Depositor EDS
% Data completeness (in resolution range)	97.3 (42.52-1.75) 97.3 (42.52-1.75)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.15 (at 1.75Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.161 , 0.197 0.158 , 0.195	Depositor DCC
$R_{free}$ test set	4184 reflections (11.48%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.8	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 57.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 41444 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3682	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.40	0/3217	0.53	0/4329

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3162	0	3094	42	0
2	A	4	0	0	0	0
3	A	516	0	0	23	0
All	All	3682	0	3094	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:446:GLN:HG2	3:A:804:HOH:O	1.54	1.08

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60[A]:LEU:HD12	1:A:60[A]:LEU:H	1.37	0.88
1:A:66:MET:CE	3:A:689:HOH:O	2.21	0.87
1:A:66:MET:HE2	3:A:689:HOH:O	1.79	0.79
1:A:164:GLU:HG3	3:A:667:HOH:O	1.83	0.78
1:A:251:ILE:CD1	1:A:269:GLU:HG2	2.14	0.78
1:A:248:GLU:HG2	3:A:954:HOH:O	1.87	0.74
1:A:301:TYR:CD2	1:A:303:THR:HG23	2.24	0.72
1:A:49:HIS:HD2	1:A:52:ARG:HH21	1.38	0.71
1:A:142:ILE:O	1:A:235:HIS:HE1	1.73	0.70
1:A:301:TYR:HD2	1:A:303:THR:HG23	1.59	0.67
1:A:49:HIS:CD2	1:A:52:ARG:HH21	2.12	0.67
1:A:181:ASN:HD21	1:A:288:PHE:H	1.45	0.65
1:A:66:MET:HE3	3:A:689:HOH:O	1.88	0.65
1:A:161:LYS:HA	3:A:667:HOH:O	1.96	0.64
1:A:307:LEU:HD23	3:A:527:HOH:O	1.97	0.64
1:A:79:LYS:HG2	1:A:93:TYR:CD1	2.32	0.64
1:A:284:LYS:HE2	3:A:580:HOH:O	2.03	0.58
1:A:85:GLU:HG3	3:A:772:HOH:O	2.04	0.58
1:A:307:LEU:HD12	3:A:928:HOH:O	2.06	0.56
1:A:91:LEU:HD23	1:A:93:TYR:OH	2.07	0.53
1:A:235:HIS:HD2	3:A:618:HOH:O	1.92	0.53
1:A:387:GLU:HG3	3:A:774:HOH:O	2.08	0.53
1:A:307:LEU:HB2	3:A:928:HOH:O	2.09	0.52
1:A:183:HIS:HD2	3:A:656:HOH:O	1.93	0.51
1:A:85:GLU:O	1:A:86:ASP:HB2	2.11	0.49
1:A:68:ASP:OD1	1:A:75:LYS:HE3	2.13	0.48
1:A:60[A]:LEU:HD13	3:A:925:HOH:O	2.14	0.46
1:A:119:LYS:NZ	1:A:309:GLU:HG3	2.30	0.46
1:A:70:GLN:HG2	3:A:797:HOH:O	2.16	0.45
1:A:397:LYS:HG2	3:A:735:HOH:O	2.16	0.45
1:A:99:THR:HG23	3:A:864:HOH:O	2.17	0.45
1:A:450:ALA:HB1	3:A:473:HOH:O	2.17	0.44
1:A:184:LYS:HG3	3:A:660:HOH:O	2.18	0.44
1:A:308:THR:OG1	1:A:311:GLN:HG3	2.19	0.43
1:A:128:ARG:CZ	1:A:205:ILE:HD13	2.50	0.41
1:A:79:LYS:NZ	3:A:859:HOH:O	2.54	0.41
1:A:57:LEU:O	1:A:60[A]:LEU:HD11	2.20	0.41
1:A:119:LYS:HZ3	1:A:309:GLU:CD	2.24	0.41
1:A:389:ARG:CZ	3:A:679:HOH:O	2.69	0.41
1:A:79:LYS:HG2	1:A:93:TYR:HD1	1.85	0.40
1:A:60[A]:LEU:CD1	1:A:60[A]:LEU:H	2.15	0.40



There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	393/449 (88%)	387 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	346/382 (91%)	342 (99%)	4 (1%)	78	62

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60[A]	LEU
1	A	60[B]	LEU
1	A	181	ASN
1	A	184	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	181	ASN
1	A	183	HIS
1	A	235	HIS
1	A	311	GLN
1	A	345	ASN
1	A	414	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	CRO	A	224	1	23,23,24	4.19	5 (21%)	29,32,34	4.62	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	A	224	1	-	0/12/31/32	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	224	CRO	C2-N3	-3.42	1.32	1.39
1	A	224	CRO	CG2-CB2	2.35	1.51	1.46
1	A	224	CRO	O2-C2	2.83	1.29	1.23

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	224	CRO	C1-N2	4.26	1.38	1.32
1	A	224	CRO	CB2-CA2	18.76	1.51	1.35

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	224	CRO	O2-C2-CA2	-12.70	124.09	130.95
1	A	224	CRO	C2-CA2-N2	-7.86	102.64	108.91
1	A	224	CRO	CG2-CB2-CA2	-7.32	120.71	130.22
1	A	224	CRO	CB2-CA2-N2	3.42	134.78	128.67
1	A	224	CRO	CA2-N2-C1	3.59	108.96	105.71
1	A	224	CRO	CA2-C2-N3	17.22	112.03	103.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	393/449 (87%)	-0.08	14 (3%)	46 52	10, 20, 47, 101	1 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	303	THR	13.3
1	A	145	GLY	9.1
1	A	146	MET	8.3
1	A	307	LEU	4.1
1	A	143	THR	3.6
1	A	378	LYS	3.5
1	A	377	ARG	2.6
1	A	309	GLU	2.5
1	A	144	LEU	2.3
1	A	161	LYS	2.3
1	A	310	GLU	2.3
1	A	125	ASN	2.3
1	A	379	MET	2.3
1	A	450	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CRO	A	224	22/23	0.98	0.08	-	9,13,18,22	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	A	505	1/1	1.00	0.06	-0.46	14,14,14,14	0
2	CA	A	506	1/1	1.00	0.06	-1.19	15,15,15,15	0
2	CA	A	507	1/1	0.98	0.05	-2.17	21,21,21,21	0
2	CA	A	508	1/1	1.00	0.04	-3.39	14,14,14,14	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.