



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:33 PM GMT

PDB ID : 4F1E
Title : The Crystal Structure of a Human MitoNEET mutant with Asp 67 replaced by a Gly
Authors : Baxter, E.I.; Zuris, J.A.; Wang, C.; Axelrod, H.L.; Cohen, A.E.; Paddock, M.L.; Nechushtai, R.; Onuchic, J.N.; Jennings, P.A.
Deposited on : 2012-05-06
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	76	 79% 5% • 14%
1	B	76	 83% • 14%
1	C	76	 87% 13%
1	D	76	 86% 14%
1	E	76	 86% 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	76	
1	G	76	
1	H	76	
1	I	76	
1	J	76	
1	K	76	
1	L	76	
1	M	76	
1	N	76	
1	O	76	
1	P	76	
1	Q	76	
1	R	76	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FES	B	201	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9244 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CDGSH iron-sulfur domain-containing protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	65	Total	C	N	O	S	0	0	0
			506	320	91	90	5			
1	B	65	Total	C	N	O	S	0	0	0
			513	325	91	92	5			
1	C	66	Total	C	N	O	S	0	0	0
			506	320	90	91	5			
1	D	65	Total	C	N	O	S	0	0	0
			508	321	90	92	5			
1	E	65	Total	C	N	O	S	0	0	0
			508	321	92	90	5			
1	F	65	Total	C	N	O	S	0	0	0
			508	321	92	90	5			
1	G	66	Total	C	N	O	S	0	0	0
			507	321	92	89	5			
1	H	66	Total	C	N	O	S	0	0	0
			514	325	91	93	5			
1	I	65	Total	C	N	O	S	0	0	0
			508	322	91	90	5			
1	J	66	Total	C	N	O	S	0	1	0
			510	322	90	93	5			
1	K	66	Total	C	N	O	S	0	0	0
			509	323	92	89	5			
1	L	64	Total	C	N	O	S	0	0	0
			501	318	91	87	5			
1	M	65	Total	C	N	O	S	0	0	0
			501	318	92	86	5			
1	N	65	Total	C	N	O	S	0	0	0
			510	323	92	90	5			
1	O	62	Total	C	N	O	S	0	0	0
			486	308	87	87	4			
1	P	63	Total	C	N	O	S	0	0	0
			495	313	89	88	5			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Q	64	Total	C	N	O	S	0	0	0
			495	315	89	87	4			
1	R	64	Total	C	N	O	S	0	0	0
			498	314	89	91	4			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
B	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
C	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
D	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
E	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
F	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
G	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
H	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
I	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
J	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
K	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
L	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
M	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
N	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
O	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
P	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
Q	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45
R	67	GLY	ASP	ENGINEERED MUTATION	UNP Q9NZ45

- Molecule 2 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			4	2	2		
2	B	1	Total	Fe	S	0	0
			4	2	2		
2	C	1	Total	Fe	S	0	0
			4	2	2		
2	D	1	Total	Fe	S	0	0
			4	2	2		
2	E	1	Total	Fe	S	0	0
			4	2	2		
2	F	1	Total	Fe	S	0	0
			4	2	2		
2	G	1	Total	Fe	S	0	0
			4	2	2		
2	H	1	Total	Fe	S	0	0
			4	2	2		
2	I	1	Total	Fe	S	0	0
			4	2	2		
2	J	1	Total	Fe	S	0	0
			4	2	2		
2	K	1	Total	Fe	S	0	0
			4	2	2		
2	L	1	Total	Fe	S	0	0
			4	2	2		
2	M	1	Total	Fe	S	0	0
			4	2	2		
2	N	1	Total	Fe	S	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	O	1	Total 4	Fe 2	S 2	0	0
2	P	1	Total 4	Fe 2	S 2	0	0
2	Q	1	Total 4	Fe 2	S 2	0	0
2	R	1	Total 4	Fe 2	S 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	11	Total 11	O 11	0	0
3	B	6	Total 6	O 6	0	0
3	C	7	Total 7	O 7	0	0
3	D	7	Total 7	O 7	0	0
3	E	5	Total 5	O 5	0	0
3	F	3	Total 3	O 3	0	0
3	G	4	Total 4	O 4	0	0
3	H	4	Total 4	O 4	0	0
3	I	11	Total 11	O 11	0	0
3	J	5	Total 5	O 5	0	0
3	K	8	Total 8	O 8	0	0
3	L	5	Total 5	O 5	0	0
3	M	3	Total 3	O 3	0	0
3	N	8	Total 8	O 8	0	0
3	O	1	Total 1	O 1	0	0

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	R	1	Total	O	0	0
			1	1		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CDGSH iron-sulfur domain-containing protein 1

Chain A: 




- Molecule 1: CDGSH iron-sulfur domain-containing protein 1

Chain B: 





- Molecule 1: CDGSH iron-sulfur domain-containing protein 1

Chain C: 





- Molecule 1: CDGSH iron-sulfur domain-containing protein 1

Chain D: 




- Molecule 1: CDGSH iron-sulfur domain-containing protein 1

Chain E: 





- Molecule 1: CDGSH iron-sulfur domain-containing protein 1



Chain F: 




- Molecule 1: CDGSH iron-sulfur domain-containing protein 1





- Molecule 1: CDGSH iron-sulfur domain-containing protein 1





- Molecule 1: CDGSH iron-sulfur domain-containing protein 1





- Molecule 1: CDGSH iron-sulfur domain-containing protein 1




- Molecule 1: CDGSH iron-sulfur domain-containing protein 1




- Molecule 1: CDGSH iron-sulfur domain-containing protein 1




- Molecule 1: CDGSH iron-sulfur domain-containing protein 1








- Molecule 1: CDGSH iron-sulfur domain-containing protein 1





- Molecule 1: CDGSH iron-sulfur domain-containing protein 1





- Molecule 1: CDGSH iron-sulfur domain-containing protein 1



- Molecule 1: CDGSH iron-sulfur domain-containing protein 1



- Molecule 1: CDGSH iron-sulfur domain-containing protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.40Å 110.86Å 207.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.54 – 2.40 49.54 – 2.40	Depositor EDS
% Data completeness (in resolution range)	(Not available) (49.54-2.40) 99.4 (49.54-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 2.39Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.193 , 0.227 0.207 , 0.238	Depositor DCC
R_{free} test set	2624 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	54.2	Xtriage
Anisotropy	0.511	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 49.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 51537 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9244	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.51	0/518	0.67	0/696
1	B	0.53	0/525	0.66	0/705
1	C	0.49	0/517	0.63	0/695
1	D	0.50	0/519	0.64	0/697
1	E	0.48	0/520	0.67	0/698
1	F	0.48	0/520	0.65	0/699
1	G	0.47	0/519	0.65	0/698
1	H	0.50	0/525	0.71	0/704
1	I	0.49	0/520	0.68	0/698
1	J	0.49	0/525	0.64	0/709
1	K	0.52	0/521	0.69	0/700
1	L	0.50	0/513	0.62	0/689
1	M	0.52	0/513	0.65	0/689
1	N	0.49	0/522	0.65	0/701
1	O	0.48	0/498	0.66	0/670
1	P	0.49	0/507	0.67	0/682
1	Q	0.46	0/507	0.65	0/683
1	R	0.45	0/510	0.61	0/688
All	All	0.49	0/9299	0.66	0/12501

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	506	0	482	5	0
1	B	513	0	491	2	0
1	C	506	0	479	0	0
1	D	508	0	489	0	0
1	E	508	0	489	0	0
1	F	508	0	488	1	0
1	G	507	0	480	2	0
1	H	514	0	496	2	0
1	I	508	0	486	0	0
1	J	510	0	469	0	0
1	K	509	0	484	2	0
1	L	501	0	486	6	0
1	M	501	0	483	2	0
1	N	510	0	492	2	0
1	O	486	0	455	1	0
1	P	495	0	472	1	0
1	Q	495	0	464	1	0
1	R	498	0	462	0	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
2	D	4	0	0	0	0
2	E	4	0	0	0	0
2	F	4	0	0	0	0
2	G	4	0	0	0	0
2	H	4	0	0	0	0
2	I	4	0	0	0	0
2	J	4	0	0	0	0
2	K	4	0	0	0	0
2	L	4	0	0	0	0
2	M	4	0	0	0	0
2	N	4	0	0	0	0
2	O	4	0	0	0	0
2	P	4	0	0	0	0
2	Q	4	0	0	0	0
2	R	4	0	0	0	0
3	A	11	0	0	1	0
3	B	6	0	0	0	0
3	C	7	0	0	0	0
3	D	7	0	0	0	0
3	E	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	3	0	0	0	0
3	G	4	0	0	0	0
3	H	4	0	0	0	0
3	I	11	0	0	0	0
3	J	5	0	0	0	0
3	K	8	0	0	1	0
3	L	5	0	0	0	0
3	M	3	0	0	0	0
3	N	8	0	0	0	0
3	O	1	0	0	0	0
3	R	1	0	0	0	0
All	All	9244	0	8647	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:GLN:NE2	3:A:308:HOH:O	2.23	0.72
1:L:94:THR:CG2	1:L:96:ASP:HB2	2.24	0.68
1:L:94:THR:HG23	1:L:96:ASP:HB2	1.76	0.68
1:N:79:LYS:HG2	1:N:84:ASP:HB3	1.75	0.67
1:A:106:LYS:HE3	1:B:61:ASP:OD2	1.95	0.67
1:L:90:HIS:O	1:L:94:THR:HG22	2.04	0.57
1:H:107:GLU:O	1:H:108:THR:HG22	2.11	0.51
1:G:106:LYS:HE3	1:H:61:ASP:OD2	2.11	0.50
1:Q:79:LYS:HG2	1:Q:84:ASP:HB3	1.93	0.50
1:N:79:LYS:CG	1:N:84:ASP:HB3	2.41	0.49
1:L:90:HIS:CE1	1:L:94:THR:HG21	2.49	0.48
1:K:79:LYS:NZ	3:K:308:HOH:O	2.48	0.47
1:A:57:VAL:HG22	1:B:100:PRO:HD2	1.99	0.45
1:O:100:PRO:HD2	1:P:57:VAL:HG22	2.01	0.42
1:A:64:ASP:CG	1:M:58:HIS:HE2	2.22	0.42
1:L:70:VAL:HG12	1:L:83:CYS:HB3	2.02	0.42
1:F:51:LYS:HA	1:F:51:LYS:HD3	1.88	0.41
1:L:90:HIS:NE2	1:L:94:THR:HG21	2.35	0.41
1:G:51:LYS:HD3	1:G:51:LYS:HA	1.96	0.41
1:A:81:PRO:HD2	1:M:79:LYS:HD2	2.01	0.41
1:K:51:LYS:HA	1:K:51:LYS:HD3	1.96	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	63/76 (83%)	62 (98%)	1 (2%)	0	100	100
1	B	63/76 (83%)	62 (98%)	1 (2%)	0	100	100
1	C	64/76 (84%)	63 (98%)	1 (2%)	0	100	100
1	D	63/76 (83%)	62 (98%)	1 (2%)	0	100	100
1	E	63/76 (83%)	62 (98%)	1 (2%)	0	100	100
1	F	63/76 (83%)	60 (95%)	1 (2%)	2 (3%)	5	4
1	G	64/76 (84%)	62 (97%)	2 (3%)	0	100	100
1	H	64/76 (84%)	62 (97%)	2 (3%)	0	100	100
1	I	63/76 (83%)	62 (98%)	1 (2%)	0	100	100
1	J	65/76 (86%)	64 (98%)	1 (2%)	0	100	100
1	K	64/76 (84%)	62 (97%)	2 (3%)	0	100	100
1	L	62/76 (82%)	60 (97%)	2 (3%)	0	100	100
1	M	63/76 (83%)	62 (98%)	1 (2%)	0	100	100
1	N	63/76 (83%)	62 (98%)	1 (2%)	0	100	100
1	O	60/76 (79%)	57 (95%)	3 (5%)	0	100	100
1	P	61/76 (80%)	57 (93%)	3 (5%)	1 (2%)	12	16
1	Q	62/76 (82%)	56 (90%)	5 (8%)	1 (2%)	12	16
1	R	62/76 (82%)	61 (98%)	1 (2%)	0	100	100
All	All	1132/1368 (83%)	1098 (97%)	30 (3%)	4 (0%)	39	56

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Q	95	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	P	50	GLN
1	F	66	GLY
1	F	67	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	52/67 (78%)	51 (98%)	1 (2%)	65	83
1	B	53/67 (79%)	53 (100%)	0	100	100
1	C	51/67 (76%)	51 (100%)	0	100	100
1	D	53/67 (79%)	53 (100%)	0	100	100
1	E	53/67 (79%)	53 (100%)	0	100	100
1	F	53/67 (79%)	53 (100%)	0	100	100
1	G	51/67 (76%)	51 (100%)	0	100	100
1	H	53/67 (79%)	53 (100%)	0	100	100
1	I	52/67 (78%)	52 (100%)	0	100	100
1	J	51/67 (76%)	51 (100%)	0	100	100
1	K	51/67 (76%)	51 (100%)	0	100	100
1	L	52/67 (78%)	52 (100%)	0	100	100
1	M	51/67 (76%)	51 (100%)	0	100	100
1	N	53/67 (79%)	53 (100%)	0	100	100
1	O	49/67 (73%)	49 (100%)	0	100	100
1	P	52/67 (78%)	51 (98%)	1 (2%)	65	83
1	Q	49/67 (73%)	49 (100%)	0	100	100
1	R	51/67 (76%)	51 (100%)	0	100	100
All	All	930/1206 (77%)	928 (100%)	2 (0%)	95	99

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	LYS
1	P	63	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FES	A	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	B	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	C	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	D	500	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	E	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	F	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	G	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	H	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	I	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	J	201	1	0,4,4	0.00	-	0,4,4	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FES	K	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	L	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	M	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	N	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	O	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	P	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	Q	201	1	0,4,4	0.00	-	0,4,4	0.00	-
2	FES	R	201	1	0,4,4	0.00	-	0,4,4	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FES	A	201	1	-	0/0/4/4	0/1/1/1
2	FES	B	201	1	-	0/0/4/4	0/1/1/1
2	FES	C	201	1	-	0/0/4/4	0/1/1/1
2	FES	D	500	1	-	0/0/4/4	0/1/1/1
2	FES	E	201	1	-	0/0/4/4	0/1/1/1
2	FES	F	201	1	-	0/0/4/4	0/1/1/1
2	FES	G	201	1	-	0/0/4/4	0/1/1/1
2	FES	H	201	1	-	0/0/4/4	0/1/1/1
2	FES	I	201	1	-	0/0/4/4	0/1/1/1
2	FES	J	201	1	-	0/0/4/4	0/1/1/1
2	FES	K	201	1	-	0/0/4/4	0/1/1/1
2	FES	L	201	1	-	0/0/4/4	0/1/1/1
2	FES	M	201	1	-	0/0/4/4	0/1/1/1
2	FES	N	201	1	-	0/0/4/4	0/1/1/1
2	FES	O	201	1	-	0/0/4/4	0/1/1/1
2	FES	P	201	1	-	0/0/4/4	0/1/1/1
2	FES	Q	201	1	-	0/0/4/4	0/1/1/1
2	FES	R	201	1	-	0/0/4/4	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	65/76 (85%)	-0.17	0 100 100	39, 54, 85, 102	0
1	B	65/76 (85%)	-0.20	0 100 100	40, 57, 84, 109	0
1	C	66/76 (86%)	0.02	0 100 100	42, 71, 106, 120	0
1	D	65/76 (85%)	-0.03	1 (1%) 76 75	44, 69, 107, 137	0
1	E	65/76 (85%)	-0.15	0 100 100	47, 60, 93, 105	0
1	F	65/76 (85%)	-0.11	1 (1%) 76 75	49, 64, 96, 142	0
1	G	66/76 (86%)	0.11	3 (4%) 37 38	53, 76, 95, 122	0
1	H	66/76 (86%)	0.10	1 (1%) 76 75	51, 73, 101, 117	0
1	I	65/76 (85%)	-0.00	1 (1%) 76 75	44, 62, 89, 124	0
1	J	66/76 (86%)	-0.16	0 100 100	48, 66, 93, 108	0
1	K	66/76 (86%)	0.15	1 (1%) 76 75	45, 71, 106, 122	0
1	L	64/76 (84%)	0.28	4 (6%) 23 24	44, 69, 102, 124	0
1	M	65/76 (85%)	0.03	0 100 100	47, 70, 96, 120	0
1	N	65/76 (85%)	-0.02	0 100 100	46, 71, 98, 110	0
1	O	62/76 (81%)	0.24	0 100 100	54, 83, 117, 120	0
1	P	63/76 (82%)	0.83	10 (15%) 3 2	68, 96, 127, 149	0
1	Q	64/76 (84%)	0.43	3 (4%) 35 36	60, 89, 121, 135	0
1	R	64/76 (84%)	1.16	13 (20%) 1 1	61, 93, 134, 144	0
All	All	1167/1368 (85%)	0.14	38 (3%) 50 50	39, 71, 114, 149	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	P	91	ASN	6.5
1	P	93	GLU	5.8
1	R	64	ASP	5.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	Q	45	ILE	5.0
1	R	47	LEU	4.4
1	L	45	ILE	4.3
1	P	98	VAL	3.9
1	R	94	THR	3.9
1	P	92	GLU	3.7
1	R	65	LEU	3.7
1	R	98	VAL	3.6
1	R	95	GLY	3.4
1	K	45	ILE	3.2
1	P	95	GLY	3.2
1	R	48	HIS	3.0
1	D	107	GLU	3.0
1	P	45	ILE	3.0
1	R	91	ASN	3.0
1	H	47	LEU	2.8
1	R	78	LYS	2.7
1	P	75	TRP	2.4
1	Q	55	LYS	2.4
1	P	96	ASP	2.4
1	P	94	THR	2.4
1	L	91	ASN	2.3
1	R	49	ILE	2.2
1	R	45	ILE	2.2
1	G	44	MET	2.2
1	I	47	LEU	2.2
1	L	43	ALA	2.2
1	F	106	LYS	2.2
1	G	43	ALA	2.1
1	L	89	LYS	2.1
1	P	97	ASN	2.1
1	R	107	GLU	2.1
1	G	75	TRP	2.1
1	R	97	ASN	2.0
1	Q	93	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	FES	B	201	4/4	1.00	0.17	2.25	51,51,53,53	0
2	FES	A	201	4/4	0.98	0.16	1.95	41,43,43,44	0
2	FES	J	201	4/4	1.00	0.17	1.50	56,57,58,58	0
2	FES	I	201	4/4	1.00	0.16	1.14	52,52,53,53	0
2	FES	D	500	4/4	0.99	0.16	0.78	60,61,61,62	0
2	FES	O	201	4/4	0.99	0.16	0.33	67,67,68,68	0
2	FES	E	201	4/4	0.99	0.13	0.24	50,50,51,51	0
2	FES	N	201	4/4	0.99	0.14	0.18	59,60,60,61	0
2	FES	F	201	4/4	0.99	0.14	0.14	57,57,58,60	0
2	FES	L	201	4/4	0.99	0.13	-0.10	56,56,57,57	0
2	FES	M	201	4/4	1.00	0.13	-0.24	57,57,58,59	0
2	FES	C	201	4/4	0.99	0.14	-0.34	63,64,64,66	0
2	FES	H	201	4/4	0.98	0.13	-0.45	66,67,67,68	0
2	FES	P	201	4/4	0.98	0.15	-0.48	93,94,94,95	0
2	FES	Q	201	4/4	0.99	0.13	-0.51	77,77,78,78	0
2	FES	K	201	4/4	1.00	0.13	-0.75	58,58,59,59	0
2	FES	G	201	4/4	0.97	0.10	-1.90	69,70,70,72	0
2	FES	R	201	4/4	0.98	0.10	-2.16	89,90,90,90	0

6.5 Other polymers [i](#)

There are no such residues in this entry.