



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:27 PM GMT

PDB ID : 4F23  
Title : Influenza A virus hemagglutinin H16 HA0 structure with an alpha-helix con-  
formation in the cleavage site: a potential drug target  
Authors : Lu, X.; Shi, Y.; Gao, F.; Xiao, H.; Qi, J.; Gao, G.F.  
Deposited on : 2012-05-07  
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	515	<div> <div></div> <div>80%17% ..</div> </div>
1	B	515	<div> <div>%</div> <div>74%22% ...</div> </div>
1	C	515	<div> <div>%</div> <div>82%15% ..</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	509	Total	C	N	O	S	0	3	0
			4040	2522	719	781	18			
1	B	509	Total	C	N	O	S	0	1	0
			4025	2513	716	778	18			
1	C	509	Total	C	N	O	S	0	0	0
			4019	2510	715	776	18			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	EXPRESSION TAG	UNP Q5DL24
A	-4	HIS	-	EXPRESSION TAG	UNP Q5DL24
A	-3	HIS	-	EXPRESSION TAG	UNP Q5DL24
A	-2	HIS	-	EXPRESSION TAG	UNP Q5DL24
A	-1	HIS	-	EXPRESSION TAG	UNP Q5DL24
A	0	HIS	-	EXPRESSION TAG	UNP Q5DL24
A	1	ALA	-	SEE REMARK 999	UNP Q5DL24
A	2	ASP	-	SEE REMARK 999	UNP Q5DL24
A	3	GLY	-	SEE REMARK 999	UNP Q5DL24
A	4	ILE	-	SEE REMARK 999	UNP Q5DL24
A	5	GLN	-	SEE REMARK 999	UNP Q5DL24
A	505	ARG	-	SEE REMARK 999	UNP Q5DL24
A	506	LEU	-	SEE REMARK 999	UNP Q5DL24
A	507	VAL	-	SEE REMARK 999	UNP Q5DL24
A	508	PRO	-	SEE REMARK 999	UNP Q5DL24
A	509	ARG	-	SEE REMARK 999	UNP Q5DL24
B	-5	HIS	-	EXPRESSION TAG	UNP Q5DL24
B	-4	HIS	-	EXPRESSION TAG	UNP Q5DL24
B	-3	HIS	-	EXPRESSION TAG	UNP Q5DL24
B	-2	HIS	-	EXPRESSION TAG	UNP Q5DL24
B	-1	HIS	-	EXPRESSION TAG	UNP Q5DL24
B	0	HIS	-	EXPRESSION TAG	UNP Q5DL24
B	1	ALA	-	SEE REMARK 999	UNP Q5DL24

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	2	ASP	-	SEE REMARK 999	UNP Q5DL24
B	3	GLY	-	SEE REMARK 999	UNP Q5DL24
B	4	ILE	-	SEE REMARK 999	UNP Q5DL24
B	5	GLN	-	SEE REMARK 999	UNP Q5DL24
B	505	ARG	-	SEE REMARK 999	UNP Q5DL24
B	506	LEU	-	SEE REMARK 999	UNP Q5DL24
B	507	VAL	-	SEE REMARK 999	UNP Q5DL24
B	508	PRO	-	SEE REMARK 999	UNP Q5DL24
B	509	ARG	-	SEE REMARK 999	UNP Q5DL24
C	-5	HIS	-	EXPRESSION TAG	UNP Q5DL24
C	-4	HIS	-	EXPRESSION TAG	UNP Q5DL24
C	-3	HIS	-	EXPRESSION TAG	UNP Q5DL24
C	-2	HIS	-	EXPRESSION TAG	UNP Q5DL24
C	-1	HIS	-	EXPRESSION TAG	UNP Q5DL24
C	0	HIS	-	EXPRESSION TAG	UNP Q5DL24
C	1	ALA	-	SEE REMARK 999	UNP Q5DL24
C	2	ASP	-	SEE REMARK 999	UNP Q5DL24
C	3	GLY	-	SEE REMARK 999	UNP Q5DL24
C	4	ILE	-	SEE REMARK 999	UNP Q5DL24
C	5	GLN	-	SEE REMARK 999	UNP Q5DL24
C	505	ARG	-	SEE REMARK 999	UNP Q5DL24
C	506	LEU	-	SEE REMARK 999	UNP Q5DL24
C	507	VAL	-	SEE REMARK 999	UNP Q5DL24
C	508	PRO	-	SEE REMARK 999	UNP Q5DL24
C	509	ARG	-	SEE REMARK 999	UNP Q5DL24

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	B	2	Total	C	N	O	0	0
			28	16	2	10		
2	C	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	676	Total	O	0	0
			676	676		
4	B	650	Total	O	0	0
			650	650		
4	C	654	Total	O	0	0
			654	654		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Hemagglutinin




#### • Molecule 1: Hemagglutinin



#### • Molecule 1: Hemagglutinin





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.33Å 240.94Å 69.67Å 90.00° 119.79° 90.00°	Depositor
Resolution (Å)	25.00 – 1.70 24.15 – 1.71	Depositor EDS
% Data completeness (in resolution range)	97.5 (25.00-1.70) 97.6 (24.15-1.71)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.09 (at 1.71Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.160 , 0.192 (Not available) , (Not available)	Depositor DCC
$R_{free}$ test set	NotAvailable	DCC
Wilson B-factor (Å <sup>2</sup> )	16.2	Xtriage
Anisotropy	0.759	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 31.3	EDS
Estimated twinning fraction	0.659 for H, K, L 0.209 for -H-L, K, H 0.132 for L, K, -H-L 0.217 for -h-l,k,h 0.217 for l,k,-h-l 0.097 for h,-k,-h-l 0.095 for -h-l,-k,l 0.090 for l,-k,h	Xtriage
Reported twinning fraction	0.659 for H, K, L 0.209 for -H-L, K, H 0.132 for L, K, -H-L	Depositor
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.23$	Xtriage
Outliers	0 of 210913 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	14190	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.27	11/4123 (0.3%)	1.03	12/5583 (0.2%)
1	B	1.28	11/4108 (0.3%)	1.04	13/5565 (0.2%)
1	C	1.30	15/4102 (0.4%)	1.00	6/5556 (0.1%)
All	All	1.29	37/12333 (0.3%)	1.02	31/16704 (0.2%)

The worst 5 of 37 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	431	GLU	CB-CG	8.63	1.68	1.52
1	C	348	GLY	N-CA	-7.75	1.34	1.46
1	C	256	TYR	CE1-CZ	-7.22	1.29	1.38
1	A	208	TRP	CE3-CZ3	7.21	1.50	1.38
1	B	493	GLU	CB-CG	6.67	1.64	1.52

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	509	ARG	NE-CZ-NH1	10.96	125.78	120.30
1	A	265	ARG	NE-CZ-NH2	-9.77	115.41	120.30
1	A	265	ARG	NE-CZ-NH1	9.27	124.94	120.30
1	C	265	ARG	NE-CZ-NH2	-9.08	115.76	120.30
1	B	509	ARG	NE-CZ-NH2	-8.48	116.06	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4040	0	3923	70	0
1	B	4025	0	3908	107	0
1	C	4019	0	3903	82	0
2	A	28	0	25	0	0
2	B	28	0	25	0	0
2	C	28	0	25	0	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0
3	C	14	0	13	0	0
4	A	676	0	0	30	0
4	B	650	0	0	48	0
4	C	654	0	0	40	0
All	All	14190	0	11848	243	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 243 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:480:ILE:HD12	4:B:1261:HOH:O	1.25	1.27
1:A:90:ALA:HB3	4:A:1086:HOH:O	1.42	1.20
1:B:467:SER:HB2	4:B:872:HOH:O	1.58	1.04
1:B:459:ILE:HB	4:B:872:HOH:O	1.58	1.03
1:A:202:LEU:HB3	4:A:1211:HOH:O	1.60	1.00

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	510/515 (99%)	495 (97%)	14 (3%)	1 (0%)	52	32
1	B	508/515 (99%)	488 (96%)	17 (3%)	3 (1%)	30	12
1	C	507/515 (98%)	488 (96%)	18 (4%)	1 (0%)	52	32
All	All	1525/1545 (99%)	1471 (96%)	49 (3%)	5 (0%)	46	26

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	19	ASP
1	B	339	GLU
1	C	348	GLY
1	B	348	GLY
1	B	338	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	442/445 (99%)	428 (97%)	14 (3%)	46	24
1	B	440/445 (99%)	423 (96%)	17 (4%)	39	16
1	C	439/445 (99%)	424 (97%)	15 (3%)	44	21
All	All	1321/1335 (99%)	1275 (96%)	46 (4%)	43	20

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	173	ARG
1	B	288	ASN
1	C	388	ASN
1	B	234	LEU
1	B	272	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	41	ASN
1	B	388	ASN
1	C	445	ASN
1	B	92	HIS
1	B	277	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

6 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	601	1,2	14,14,15	0.75	0	15,19,21	0.71	1 (6%)
2	NAG	A	602	2	14,14,15	1.26	2 (14%)	15,19,21	1.63	3 (20%)
2	NAG	B	601	1,2	14,14,15	0.53	0	15,19,21	1.45	4 (26%)
2	NAG	B	602	2	14,14,15	0.60	0	15,19,21	1.46	1 (6%)
2	NAG	C	601	1,2	14,14,15	0.86	0	15,19,21	0.94	0
2	NAG	C	602	2	14,14,15	0.61	0	15,19,21	1.46	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	601	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	602	2	-	0/6/23/26	0/1/1/1
2	NAG	B	601	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	602	2	-	0/6/23/26	0/1/1/1
2	NAG	C	601	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	602	2	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	602	NAG	O4-C4	2.11	1.48	1.43
2	A	602	NAG	C1-C2	2.59	1.56	1.52

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	602	NAG	C2-N2-C7	-2.92	119.28	123.04
2	A	602	NAG	C3-C4-C5	-2.86	105.20	110.20
2	B	601	NAG	O6-C6-C5	-2.49	103.11	111.33
2	B	601	NAG	O7-C7-C8	-2.33	117.79	122.06
2	C	602	NAG	C1-O5-C5	-2.18	109.48	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	603	1	14,14,15	1.83	1 (7%)	15,19,21	1.30	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	B	603	1	14,14,15	1.03	1 (7%)	15,19,21	1.23	2 (13%)
3	NAG	C	603	1	14,14,15	1.08	1 (7%)	15,19,21	1.59	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	603	1	-	0/6/23/26	0/1/1/1
3	NAG	B	603	1	-	0/6/23/26	0/1/1/1
3	NAG	C	603	1	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	603	NAG	C1-C2	2.71	1.56	1.52
3	B	603	NAG	C1-C2	3.06	1.56	1.52
3	A	603	NAG	C1-C2	6.02	1.60	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	603	NAG	C4-C3-C2	-2.87	106.76	111.23
3	B	603	NAG	C2-N2-C7	-2.75	119.51	123.04
3	A	603	NAG	O3-C3-C4	2.44	115.83	110.34
3	B	603	NAG	C1-O5-C5	2.54	115.47	112.25
3	C	603	NAG	C1-O5-C5	5.44	119.15	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	509/515 (98%)	-0.14	2 (0%) 93 94	9, 18, 28, 44	0
1	B	509/515 (98%)	-0.16	3 (0%) 90 92	9, 19, 30, 45	0
1	C	509/515 (98%)	-0.17	4 (0%) 87 90	9, 19, 29, 42	0
All	All	1527/1545 (98%)	-0.16	9 (0%) 90 92	9, 19, 29, 45	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	ALA	5.7
1	A	18	THR	5.6
1	C	1	ALA	4.2
1	B	17	SER	3.1
1	C	17	SER	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	C	601	14/15	0.84	0.12	0.67	22,32,35,35	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	B	601	14/15	0.92	0.09	-1.30	26,29,33,36	0
2	NAG	A	602	14/15	0.76	0.19	-	27,31,37,39	0
2	NAG	B	602	14/15	0.88	0.14	-	34,37,40,40	0
2	NAG	A	601	14/15	0.95	0.08	-	19,22,27,31	0
2	NAG	C	602	14/15	0.88	0.14	-	31,35,41,44	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	A	603	14/15	0.89	0.13	-	26,29,33,37	0
3	NAG	C	603	14/15	0.88	0.18	-	32,36,39,41	0
3	NAG	B	603	14/15	0.82	0.15	-	30,34,36,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.