



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:32 AM GMT

PDB ID : 3F3T  
Title : Kinase domain of cSrc in complex with inhibitor RL38 (Type III)  
Authors : Gruetter, C.; Klueter, S.; Getlik, M.; Rauh, D.  
Deposited on : 2008-10-31  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	<div> <div>2%</div> <div>66%</div> <div>22%</div> <div>•</div> <div>9%</div> </div>
1	B	286	<div> <div>11%</div> <div>64%</div> <div>21%</div> <div>5%</div> <div>9%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4347 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	261	Total	C	N	O	S	0	0	0
			2097	1345	350	384	18			
1	B	259	Total	C	N	O	S	0	0	0
			2092	1344	351	381	16			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	EXPRESSION TAG	UNP P00523
A	249	HIS	-	EXPRESSION TAG	UNP P00523
A	250	MET	-	EXPRESSION TAG	UNP P00523
A	345	CYS	SER	ENGINEERED	UNP P00523
B	248	GLY	-	EXPRESSION TAG	UNP P00523
B	249	HIS	-	EXPRESSION TAG	UNP P00523
B	250	MET	-	EXPRESSION TAG	UNP P00523
B	345	CYS	SER	ENGINEERED	UNP P00523

- Molecule 2 is 1-[1-(3-AMINOPHENYL)-3-TERT-BUTYL-1H-PYRAZOL-5-YL]-3-NAPHTHALEN-1-YLUREA (three-letter code: 1AU) (formula: C<sub>24</sub>H<sub>25</sub>N<sub>5</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			30	24	5	1		
2	A	1	Total	C	N	O	0	0
			30	24	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	55	Total	O	0	0
			55	55		
3	B	43	Total	O	0	0
			43	43		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	42.01Å 63.49Å 74.99Å 78.71° 89.71° 89.91°	Depositor
Resolution (Å)	36.71 – 2.50 36.77 – 2.49	Depositor EDS
% Data completeness (in resolution range)	100.0 (36.71-2.50) 88.8 (36.77-2.49)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.33 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.244 , 0.295 0.277 , 0.315	Depositor DCC
$R_{free}$ test set	872 reflections (3.64%)	DCC
Wilson B-factor (Å <sup>2</sup> )	36.4	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , -4.6	EDS
Estimated twinning fraction	0.030 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 24908 reflections	Xtriage
$F_o, F_c$ correlation	0.84	EDS
Total number of atoms	4347	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.69% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 1AU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.81	1/2148 (0.0%)	0.82	2/2908 (0.1%)
1	B	0.80	0/2141	0.83	2/2896 (0.1%)
All	All	0.80	1/4289 (0.0%)	0.82	4/5804 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	487	CYS	CB-SG	-7.03	1.70	1.82

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	350	LEU	CA-CB-CG	6.59	130.45	115.30
1	B	350	LEU	CA-CB-CG	5.84	128.74	115.30
1	B	360	LEU	CB-CG-CD2	5.13	119.73	111.00
1	A	322	LEU	CA-CB-CG	5.08	126.99	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2097	0	2082	49	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2092	0	2085	67	0
2	A	60	0	50	5	0
3	A	55	0	0	4	0
3	B	43	0	0	1	0
All	All	4347	0	4217	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (119) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:296:THR:O	1:B:307:PHE:CZ	1.72	1.42
1:B:257:LYS:C	1:B:257:LYS:HD2	1.64	1.16
1:B:280:GLU:HB3	1:B:296:THR:OG1	1.50	1.10
1:B:257:LYS:O	1:B:257:LYS:HD2	1.53	1.09
1:B:296:THR:O	1:B:307:PHE:HZ	1.43	1.02
1:A:508:THR:HG22	1:A:511:TYR:H	1.22	1.01
1:B:266:SER:HB3	1:B:287:ASN:OD1	1.59	0.99
1:B:297:LEU:HD12	1:B:297:LEU:H	1.28	0.96
1:B:280:GLU:HB3	1:B:296:THR:HG1	1.23	0.95
1:B:257:LYS:CD	1:B:257:LYS:C	2.42	0.85
1:B:287:ASN:ND2	1:B:287:ASN:O	2.09	0.85
1:B:257:LYS:HD3	1:B:258:ASP:O	1.81	0.80
1:B:457:THR:HG23	1:B:460:ARG:H	1.46	0.79
1:A:526:GLN:OE1	3:A:128:HOH:O	2.01	0.78
1:B:457:THR:CG2	1:B:460:ARG:H	1.99	0.75
1:A:508:THR:CG2	1:A:511:TYR:H	1.99	0.72
1:B:296:THR:O	1:B:307:PHE:CE2	2.43	0.71
1:B:286:TRP:HD1	1:B:290:THR:HB	1.56	0.70
1:A:526:GLN:HE21	1:A:526:GLN:H	1.38	0.69
1:A:384:HIS:O	1:A:385:ARG:HB2	1.91	0.69
2:A:1:1AU:NAD	3:A:132:HOH:O	2.26	0.69
1:B:262:ILE:HD13	1:B:286:TRP:HZ3	1.57	0.69
1:B:471:VAL:O	1:B:475:VAL:HG13	1.93	0.68
1:B:297:LEU:HD23	1:B:334:ILE:HG23	1.75	0.68
1:B:297:LEU:HD12	1:B:297:LEU:N	2.02	0.67
1:A:262:ILE:HG12	1:A:327:ALA:HB1	1.74	0.67
1:B:310:GLU:HB2	1:B:406:GLY:HA2	1.77	0.67
1:B:262:ILE:HD13	1:B:286:TRP:CZ3	2.30	0.66
1:A:319:HIS:HD2	1:A:321:LYS:H	1.43	0.66

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:298:LYS:HB2	1:B:299:PRO:HD3	1.77	0.66
1:A:526:GLN:NE2	1:A:526:GLN:H	1.93	0.66
1:B:264:ARG:NH2	1:B:333:PRO:O	2.31	0.63
1:B:298:LYS:HB2	1:B:299:PRO:CD	2.28	0.63
1:B:453:THR:O	1:B:457:THR:HB	1.98	0.63
1:A:275:GLN:HG3	1:A:280:GLU:HG2	1.81	0.62
1:A:476:GLU:OE2	1:A:501:LYS:NZ	2.31	0.62
1:B:384:HIS:O	1:B:385:ARG:HB2	2.00	0.61
1:B:286:TRP:CD1	1:B:286:TRP:C	2.74	0.60
1:A:272:LYS:HE3	1:A:280:GLU:OE2	2.01	0.60
1:A:493:ASP:O	1:A:497:GLN:HG3	2.02	0.58
1:B:295:LYS:HB3	1:B:336:ILE:HG23	1.86	0.58
1:A:345:CYS:SG	2:A:534:1AU:NAD	2.77	0.57
1:B:303:SER:HB2	1:B:304:PRO:HA	1.87	0.57
1:A:355:GLY:O	1:A:458:LYS:HD2	2.05	0.57
1:A:319:HIS:CD2	1:A:321:LYS:H	2.21	0.56
1:B:473:ASP:HB3	1:B:477:ARG:NH2	2.21	0.56
1:B:257:LYS:HB2	1:B:261:GLU:HG3	1.88	0.56
1:A:323:VAL:HG21	1:A:393:LEU:HD12	1.87	0.55
1:A:386:ASP:OD1	1:A:388:ARG:NE	2.35	0.55
1:B:262:ILE:CD1	1:B:286:TRP:HZ3	2.20	0.55
1:A:350:LEU:O	1:A:355:GLY:HA3	2.07	0.55
1:B:464:PRO:HB3	3:B:125:HOH:O	2.07	0.54
1:B:286:TRP:CD1	1:B:290:THR:HB	2.40	0.53
1:A:275:GLN:HG3	1:A:280:GLU:CG	2.39	0.53
1:A:388:ARG:HB3	1:A:428:TRP:CD1	2.43	0.52
1:B:362:GLN:O	1:B:366:MET:HG3	2.09	0.52
1:B:303:SER:HA	1:B:304:PRO:C	2.28	0.52
1:A:264:ARG:NH2	1:A:331:GLU:O	2.42	0.52
1:B:308:LEU:HG	1:B:312:GLN:HE21	1.73	0.52
2:A:534:1AU:OAE	2:A:534:1AU:HAL	2.10	0.51
1:A:427:LYS:NZ	3:A:114:HOH:O	2.43	0.51
1:B:457:THR:CG2	1:B:460:ARG:N	2.73	0.51
1:B:473:ASP:HB3	1:B:477:ARG:HH21	1.76	0.51
1:B:388:ARG:HB3	1:B:428:TRP:CD1	2.46	0.50
1:B:426:ILE:C	1:B:426:ILE:HD12	2.32	0.49
1:A:441:ILE:O	1:A:445:VAL:HG23	2.12	0.49
1:A:323:VAL:HG23	1:A:402:VAL:O	2.13	0.49
1:B:457:THR:HG23	1:B:457:THR:O	2.12	0.49
1:B:385:ARG:HD2	1:B:407:LEU:O	2.13	0.49
1:A:508:THR:HG23	1:A:510:GLU:H	1.78	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:322:LEU:HD13	2:A:1:1AU:HAB	1.95	0.48
1:A:261:GLU:HA	1:A:328:VAL:O	2.13	0.48
1:B:314:MET:SD	1:B:336:ILE:HD11	2.53	0.48
1:B:289:THR:HG23	1:B:289:THR:O	2.14	0.48
1:B:305:GLU:HB2	1:B:308:LEU:HD22	1.96	0.47
1:A:428:TRP:HE1	1:A:454:GLU:CD	2.19	0.47
1:B:283:MET:HG3	1:B:340:TYR:CE1	2.50	0.47
1:A:502:ASP:HB3	1:A:505:GLU:HG3	1.96	0.46
1:B:297:LEU:HD22	1:B:299:PRO:HD2	1.98	0.46
1:A:257:LYS:HB3	1:A:261:GLU:HG3	1.96	0.46
1:B:287:ASN:C	1:B:287:ASN:HD22	2.17	0.46
1:A:407:LEU:H	1:A:407:LEU:HD12	1.80	0.46
1:A:508:THR:HG22	1:A:511:TYR:N	2.07	0.46
1:A:466:MET:HA	1:A:470:GLU:OE1	2.16	0.45
1:A:476:GLU:HA	1:A:476:GLU:OE2	2.16	0.45
1:B:297:LEU:C	1:B:297:LEU:CD1	2.85	0.45
1:A:309:GLN:HB2	1:A:309:GLN:HE21	1.61	0.45
1:B:297:LEU:HD21	1:B:333:PRO:HA	1.97	0.45
1:B:500:ARG:HD3	1:B:505:GLU:HB3	1.98	0.45
1:B:285:THR:HG22	1:B:289:THR:HA	1.99	0.45
1:B:495:MET:O	1:B:498:CYS:N	2.49	0.45
1:B:286:TRP:O	1:B:286:TRP:CD1	2.70	0.44
1:B:310:GLU:HG3	1:B:405:PHE:C	2.38	0.44
1:A:310:GLU:O	1:A:314:MET:HG3	2.18	0.43
1:B:332:GLU:H	1:B:332:GLU:HG3	1.68	0.43
1:B:311:ALA:HB2	1:B:336:ILE:HD12	1.99	0.43
1:B:480:ARG:NH2	1:B:499:TRP:O	2.44	0.43
1:B:385:ARG:HH11	1:B:424:PHE:HZ	1.64	0.43
1:A:508:THR:CG2	1:A:510:GLU:HB3	2.49	0.43
1:B:297:LEU:C	1:B:297:LEU:HD13	2.39	0.43
1:B:287:ASN:HA	1:B:288:GLY:HA2	1.57	0.43
1:A:323:VAL:HG21	1:A:403:ALA:HB2	2.00	0.42
1:A:428:TRP:O	1:A:447:SER:OG	2.28	0.42
1:A:332:GLU:HA	1:A:333:PRO:C	2.40	0.42
1:B:297:LEU:HG	1:B:334:ILE:H	1.85	0.42
1:B:298:LYS:H	1:B:299:PRO:HD2	1.85	0.42
1:A:384:HIS:O	1:A:385:ARG:CB	2.62	0.42
1:B:354:MET:HE2	1:B:354:MET:HB3	1.81	0.42
1:B:311:ALA:HA	1:B:314:MET:HB2	2.02	0.41
1:A:466:MET:HE2	1:A:471:VAL:HA	2.01	0.41
1:A:524:GLU:HA	1:A:526:GLN:NE2	2.36	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:371:ALA:O	1:A:375:ALA:N	2.53	0.41
1:A:508:THR:HG23	1:A:510:GLU:N	2.36	0.41
1:A:330:SER:HA	1:A:334:ILE:HG22	2.03	0.41
1:A:486:GLU:HB2	1:A:533:LEU:HD12	2.03	0.41
1:B:261:GLU:OE1	1:B:330:SER:HB3	2.21	0.40
1:A:324:GLN:NE2	3:A:3:HOH:O	2.55	0.40
2:A:1:1AU:CAL	2:A:1:1AU:OAE	2.67	0.40
1:A:501:LYS:HA	1:A:501:LYS:HD2	1.76	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/286 (90%)	246 (96%)	10 (4%)	1 (0%)	39	61
1	B	251/286 (88%)	237 (94%)	13 (5%)	1 (0%)	39	61
All	All	508/572 (89%)	483 (95%)	23 (4%)	2 (0%)	39	61

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	298	LYS
1	A	425	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/245 (92%)	209 (92%)	17 (8%)	17	31
1	B	225/245 (92%)	198 (88%)	27 (12%)	6	12
All	All	451/490 (92%)	407 (90%)	44 (10%)	10	19

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	258	ASP
1	A	277	CYS
1	A	303	SER
1	A	305	GLU
1	A	309	GLN
1	A	312	GLN
1	A	320	GLU
1	A	332	GLU
1	A	350	LEU
1	A	356	LYS
1	A	360	LEU
1	A	438	ARG
1	A	473	ASP
1	A	508	THR
1	A	524	GLU
1	A	526	GLN
1	A	528	GLN
1	B	257	LYS
1	B	261	GLU
1	B	268	ARG
1	B	269	LEU
1	B	271	VAL
1	B	286	TRP
1	B	287	ASN
1	B	291	ARG
1	B	297	LEU
1	B	308	LEU
1	B	309	GLN
1	B	316	LYS
1	B	330	SER
1	B	334	ILE
1	B	336	ILE
1	B	350	LEU
1	B	351	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	356	LYS
1	B	360	LEU
1	B	396	GLU
1	B	455	LEU
1	B	457	THR
1	B	469	ARG
1	B	472	LEU
1	B	475	VAL
1	B	477	ARG
1	B	501	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	287	ASN
1	A	309	GLN
1	A	312	GLN
1	A	319	HIS
1	A	324	GLN
1	A	397	ASN
1	A	526	GLN
1	A	528	GLN
1	B	312	GLN
1	B	381	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	1AU	A	1	-	32,33,33	2.12	4 (12%)	40,48,48	1.55	5 (12%)
2	1AU	A	534	-	32,33,33	2.20	6 (18%)	40,48,48	1.28	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1AU	A	1	-	-	0/16/18/18	0/4/4/4
2	1AU	A	534	-	-	0/16/18/18	0/4/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	1AU	NAR-NBC	-8.65	1.23	1.39
2	A	534	1AU	NAR-NBC	-8.59	1.24	1.39
2	A	534	1AU	CAW-NBC	-4.60	1.34	1.44
2	A	1	1AU	CAW-NBC	-4.20	1.35	1.44
2	A	1	1AU	CAQ-CAY	-3.64	1.34	1.39
2	A	534	1AU	CAQ-CAY	-3.55	1.34	1.39
2	A	534	1AU	CAX-NAS	-2.10	1.35	1.41
2	A	534	1AU	CAP-CAW	2.15	1.41	1.38
2	A	1	1AU	CBD-CAY	3.06	1.57	1.52
2	A	534	1AU	CBD-CAY	3.30	1.57	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	1AU	CAK-CAW-CAP	-4.68	117.26	121.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	534	1AU	CAW-NBC-CAZ	-3.67	124.80	129.02
2	A	534	1AU	CBD-CAY-NAR	2.00	122.70	120.58
2	A	1	1AU	CAJ-CAV-CAP	2.13	121.44	118.66
2	A	1	1AU	NAT-CAU-NAS	2.71	116.79	112.53
2	A	534	1AU	CAP-CAW-NBC	3.11	122.92	119.13
2	A	1	1AU	CAP-CAW-NBC	3.12	122.94	119.13
2	A	1	1AU	CAH-CAK-CAW	3.14	122.83	118.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	1AU	3	0
2	A	534	1AU	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	261/286 (91%)	0.48	7 (2%) 58 62	2, 8, 26, 35	0
1	B	259/286 (90%)	0.73	32 (12%) 5 5	2, 8, 34, 44	0
All	All	520/572 (90%)	0.60	39 (7%) 17 19	2, 8, 32, 44	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	289	THR	5.9
1	B	307	PHE	5.3
1	B	279	GLY	4.7
1	B	334	ILE	4.4
1	B	408	ALA	3.8
1	B	297	LEU	3.6
1	B	306	ALA	3.4
1	B	335	TYR	3.3
1	B	328	VAL	3.2
1	B	304	PRO	3.1
1	B	424	PHE	3.1
1	B	299	PRO	3.1
1	B	268	ARG	3.1
1	B	298	LYS	3.0
1	B	332	GLU	2.9
1	B	272	LYS	2.9
1	A	313	VAL	2.9
1	A	328	VAL	2.9
1	B	258	ASP	2.8
1	B	331	GLU	2.8
1	A	424	PHE	2.8
1	B	271	VAL	2.6
1	B	283	MET	2.6
1	A	257	LYS	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	309	GLN	2.5
1	B	409	ARG	2.5
1	A	268	ARG	2.5
1	B	407	LEU	2.4
1	B	308	LEU	2.4
1	B	275	GLN	2.4
1	B	296	THR	2.3
1	B	313	VAL	2.3
1	B	333	PRO	2.3
1	B	270	GLU	2.2
1	B	303	SER	2.2
1	B	330	SER	2.1
1	A	299	PRO	2.1
1	B	284	GLY	2.1
1	B	329	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	1AU	A	534	30/30	0.88	0.24	1.78	25,27,28,29	0
2	1AU	A	1	30/30	0.90	0.20	0.06	15,19,24,24	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.