



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:59 AM GMT

PDB ID : 2F48
Title : Crystal Structure of A Novel Fructose 1,6-Bisphosphate and AlF₃ containing Pyrophosphate-dependent Phosphofructo-1-kinase Complex from *Borrelia burgdorferi*
Authors : Wang, M.
Deposited on : 2005-11-22
Resolution : 2.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4587 (2.14-2.10)
Clashscore	102246	5132 (2.14-2.10)
Ramachandran outliers	100387	5080 (2.14-2.10)
Sidechain outliers	100360	5081 (2.14-2.10)
RSRZ outliers	91569	4597 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	 2% 88% 10% ..
1	B	555	 6% 86% 13% ..

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 9434 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called diphosphate--fructose-6-phosphate 1-phosphotransferase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	551	Total	C	N	O	S	0	0	0
			4365	2814	714	819	18			
1	B	552	Total	C	N	O	S	0	0	0
			4360	2810	713	819	18			

- Molecule 2 is SUGAR (BETA FRUCTOSE 1,6-DIPHOSPHATE) (three-letter code: FBP) (formula: $C_6H_{14}O_{12}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			20	6	12	2		

- Molecule 3 is ALUMINUM FLUORIDE (three-letter code: AF3) (formula: AlF_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	Al	F	0	0
			4	1	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	412	Total	O	0	0
			412	412		
4	B	273	Total	O	0	0
			273	273		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: diphosphate-fructose-6-phosphate 1-phosphotransferase



- Molecule 1: diphosphate-fructose-6-phosphate 1-phosphotransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.41Å 100.98Å 150.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.72 – 2.11 55.78 – 2.11	Depositor EDS
% Data completeness (in resolution range)	94.3 (58.72-2.11) 94.3 (55.78-2.11)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.200 , 0.258 0.200 , 0.255	Depositor DCC
R_{free} test set	3533 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	32.5	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 69268 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9434	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FBP, AF3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/4454	0.62	0/6004
1	B	0.54	0/4449	0.61	0/6001
All	All	0.58	0/8903	0.62	0/12005

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4365	0	4402	37	0
1	B	4360	0	4383	46	0
2	A	20	0	10	0	0
3	B	4	0	0	0	0
4	A	412	0	0	11	0
4	B	273	0	0	6	0
All	All	9434	0	8795	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:176:ASP:HB3	4:A:1150:HOH:O	1.33	1.21
1:B:340:LYS:CB	1:B:341:GLY:HA3	1.69	1.20
1:B:340:LYS:HB3	1:B:341:GLY:CA	1.81	1.08
1:A:210:ARG:HD3	4:A:1089:HOH:O	1.57	1.01
1:A:486:MET:SD	4:A:1188:HOH:O	2.18	1.00
1:B:340:LYS:HB3	1:B:341:GLY:HA3	0.94	0.91
1:B:211:ASN:HD22	1:B:213:HIS:H	1.23	0.86
1:A:501:LYS:HE3	4:A:1231:HOH:O	1.82	0.79
1:B:3:THR:HA	4:B:931:HOH:O	1.88	0.72
1:B:344:ILE:HA	1:B:347:MET:HG2	1.74	0.67
1:B:211:ASN:ND2	1:B:213:HIS:H	1.96	0.64
1:A:59:SER:OG	4:A:1081:HOH:O	2.16	0.63
1:A:9:GLU:HG3	1:B:541:ILE:HD11	1.79	0.63
1:B:159:LEU:HD11	1:B:194:GLU:HG2	1.84	0.59
1:A:36:THR:HG23	4:A:1009:HOH:O	2.02	0.59
1:B:370:ILE:HD12	1:B:392:VAL:HG23	1.84	0.59
1:B:340:LYS:CB	1:B:341:GLY:CA	2.55	0.58
1:B:412:ARG:NE	1:B:414:GLU:OE2	2.37	0.58
1:A:509:ARG:NH1	1:A:551:LEU:O	2.37	0.57
1:B:481:PRO:HB2	4:B:1081:HOH:O	2.04	0.57
1:A:226:ILE:HD11	1:A:433:ALA:CB	2.35	0.56
1:B:495:LYS:NZ	4:B:1083:HOH:O	2.36	0.56
1:A:45:LEU:HD11	1:A:487:ASN:HA	1.88	0.55
1:A:323:SER:HB3	1:A:359:MET:HE1	1.89	0.55
1:B:381:ARG:HD2	1:B:385:GLY:O	2.07	0.54
1:B:120:ASN:ND2	1:B:165:ASN:HD21	2.06	0.54
1:B:267:HIS:HE1	4:B:1007:HOH:O	1.91	0.53
1:A:18:PRO:HD2	1:A:21:LEU:HD12	1.91	0.53
1:B:487:ASN:HD21	1:B:500:LYS:HD3	1.74	0.53
1:A:29:SER:HA	1:A:65:SER:HB2	1.92	0.52
1:B:36:THR:O	1:B:54:GLY:HA2	2.10	0.52
1:A:528:LEU:HD22	1:B:3:THR:HB	1.92	0.52
1:A:149:ILE:HG23	1:A:154:HIS:HB3	1.90	0.52
1:B:30:LEU:HD22	1:B:60:PHE:HE2	1.76	0.51
1:A:203:LYS:O	1:A:204:THR:HB	2.11	0.51
1:B:49:PHE:HB3	1:B:52:THR:O	2.11	0.51
1:A:37:GLU:CD	1:A:42:ARG:HH11	2.13	0.51
1:B:315:ILE:HG13	1:B:394:THR:OG1	2.11	0.51
1:B:289:ASP:OD1	1:B:405:ARG:NH1	2.42	0.50
1:B:211:ASN:ND2	1:B:214:ILE:H	2.11	0.49
1:A:528:LEU:CD2	1:B:3:THR:HB	2.45	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:ASP:O	1:A:180:THR:HB	2.14	0.47
1:A:46:LYS:HE2	1:A:46:LYS:O	2.14	0.47
1:B:226:ILE:HD11	1:B:433:ALA:CB	2.44	0.47
1:B:449:ASN:OD1	1:B:477:ALA:HB1	2.16	0.46
1:B:369:PHE:O	1:B:373:GLU:HG2	2.15	0.46
1:B:205:ILE:HD13	1:B:431:ARG:HB3	1.98	0.46
1:B:213:HIS:HA	1:B:466:LYS:NZ	2.31	0.45
1:A:534:GLN:O	1:A:542:VAL:HG23	2.16	0.45
1:A:480:VAL:HG13	1:A:484:MET:SD	2.56	0.45
1:B:226:ILE:CD1	1:B:433:ALA:CB	2.94	0.45
1:A:192:ASN:HB2	4:A:1223:HOH:O	2.16	0.45
1:A:27:ASN:HA	1:A:63:GLY:O	2.17	0.45
1:A:9:GLU:HG2	4:A:1206:HOH:O	2.16	0.44
1:A:77:ILE:HD11	1:A:173:ILE:HD11	2.00	0.43
1:B:382:ASP:HB2	1:B:383:PRO:HD2	2.01	0.43
1:B:30:LEU:HD22	1:B:60:PHE:CE2	2.53	0.43
1:B:467:ASN:HB3	1:B:470:LEU:HD23	2.00	0.43
1:A:43:GLN:CD	1:A:43:GLN:H	2.22	0.43
1:A:541:ILE:HB	4:A:1077:HOH:O	2.19	0.43
1:B:274:SER:HB3	1:B:317:PHE:CG	2.54	0.43
1:B:120:ASN:HD21	1:B:165:ASN:HD21	1.67	0.42
1:A:337:GLY:HA2	1:A:340:LYS:HG3	2.00	0.42
1:A:207:ALA:HA	1:A:217:SER:OG	2.19	0.42
1:A:483:THR:HA	4:A:1188:HOH:O	2.19	0.42
1:B:320:GLU:OE1	1:B:358:TYR:OH	2.29	0.42
1:A:57:ILE:HD13	1:A:481:PRO:HD2	2.02	0.42
1:B:320:GLU:HB2	4:B:891:HOH:O	2.20	0.42
1:B:373:GLU:OE2	1:B:391:ARG:HD3	2.20	0.42
1:B:315:ILE:HD13	1:B:392:VAL:HG22	2.02	0.42
1:B:500:LYS:HE2	4:B:1091:HOH:O	2.20	0.42
1:A:203:LYS:O	1:A:204:THR:CB	2.68	0.42
1:B:212:ASP:OD2	1:B:212:ASP:N	2.53	0.42
1:A:226:ILE:CD1	1:A:433:ALA:CB	2.98	0.41
1:B:75:ILE:HD12	1:B:98:ILE:HG21	2.02	0.41
1:B:392:VAL:HA	1:B:393:PRO:HD3	1.93	0.41
1:A:203:LYS:HB2	1:A:218:PHE:CE1	2.56	0.41
1:A:339:PHE:HD2	1:A:347:MET:HG2	1.86	0.41
1:A:57:ILE:N	4:A:1009:HOH:O	2.53	0.41
1:B:491:ARG:NH2	1:B:496:LYS:HD2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	549/555 (99%)	529 (96%)	18 (3%)	2 (0%)	39	36
1	B	550/555 (99%)	522 (95%)	24 (4%)	4 (1%)	26	20
All	All	1099/1110 (99%)	1051 (96%)	42 (4%)	6 (0%)	34	29

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	344	ILE
1	A	204	THR
1	B	34	GLU
1	B	340	LYS
1	A	427	GLY
1	B	427	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	483/488 (99%)	465 (96%)	18 (4%)	41	41
1	B	481/488 (99%)	460 (96%)	21 (4%)	35	32
All	All	964/976 (99%)	925 (96%)	39 (4%)	38	37

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LYS
1	A	23	LYS
1	A	34	GLU
1	A	46	LYS
1	A	66	SER
1	A	67	LEU
1	A	120	ASN
1	A	122	LYS
1	A	157	LYS
1	A	195	ASN
1	A	210	ARG
1	A	218	PHE
1	A	310	VAL
1	A	340	LYS
1	A	348	LYS
1	A	487	ASN
1	A	520	LYS
1	A	541	ILE
1	B	10	ARG
1	B	11	GLN
1	B	20	ILE
1	B	30	LEU
1	B	35	ASN
1	B	42	ARG
1	B	46	LYS
1	B	122	LYS
1	B	143	SER
1	B	157	LYS
1	B	194	GLU
1	B	204	THR
1	B	212	ASP
1	B	218	PHE
1	B	338	GLU
1	B	344	ILE
1	B	368	LEU
1	B	386	ASN
1	B	414	GLU
1	B	429	GLU
1	B	476	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	120	ASN
1	B	120	ASN
1	B	156	ASN
1	B	211	ASN
1	B	267	HIS
1	B	371	GLN
1	B	386	ASN
1	B	403	GLN
1	B	487	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FBP	A	829	-	18,20,20	1.14	2 (11%)	21,32,32	1.32	3 (14%)
3	AF3	B	861	1,4	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FBP	A	829	-	-	0/13/32/32	0/1/1/1
3	AF3	B	861	1,4	-	0/0/0/0	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	829	FBP	P2-O6P	-2.00	1.47	1.54
2	A	829	FBP	P2-O4P	2.26	1.58	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	829	FBP	O6-P2-O4P	-2.14	101.70	107.14
2	A	829	FBP	O6P-P2-O5P	2.61	117.33	107.38
2	A	829	FBP	O3P-P1-O2P	3.07	119.06	107.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	551/555 (99%)	-0.07	9 (1%) 74 79	15, 31, 59, 70	0
1	B	552/555 (99%)	0.36	36 (6%) 22 29	18, 38, 73, 96	0
All	All	1103/1110 (99%)	0.15	45 (4%) 41 49	15, 35, 68, 96	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	339	PHE	7.9
1	B	342	LEU	5.9
1	B	350	ILE	5.1
1	B	3	THR	4.9
1	B	338	GLU	4.5
1	B	337	GLY	4.4
1	B	554	PHE	4.4
1	B	352	VAL	4.3
1	A	63	GLY	4.2
1	B	68	SER	4.2
1	B	341	GLY	4.1
1	B	340	LYS	3.8
1	B	54	GLY	3.8
1	B	351	PHE	3.7
1	A	340	LYS	3.7
1	B	69	PHE	3.6
1	B	332	PHE	3.5
1	B	39	ILE	3.4
1	B	491	ARG	3.1
1	B	47	GLU	3.1
1	B	45	LEU	2.9
1	A	66	SER	2.8
1	B	355	LEU	2.8
1	B	52	THR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	153	GLU	2.6
1	B	372	PHE	2.6
1	B	71	LYS	2.6
1	B	411	LYS	2.6
1	B	188	TYR	2.5
1	A	70	SER	2.5
1	B	334	LYS	2.5
1	A	492	TYR	2.5
1	A	493	GLY	2.4
1	B	226	ILE	2.4
1	B	492	TYR	2.3
1	B	336	GLU	2.3
1	A	55	LEU	2.2
1	B	67	LEU	2.2
1	B	328	LEU	2.2
1	B	368	LEU	2.2
1	B	32	TYR	2.2
1	A	48	PHE	2.2
1	B	326	LEU	2.1
1	A	344	ILE	2.1
1	B	383	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	AF3	B	861	4/4	0.85	0.12	-0.03	50,52,53,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FBP	A	829	20/20	0.99	0.10	-0.43	20,23,27,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.