



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:34 AM GMT

PDB ID : 3F6T
Title : Crystal structure of aspartate aminotransferase (E.C. 2.6.1.1) (YP_194538.1)
from Lactobacillus acidophilus NCFM at 2.15 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2008-11-06
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	533	<div> <div>5%</div> <div> <div></div> <div>83%</div> <div>14%</div> <div>• •</div> </div> </div>
1	B	533	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>13%</div> <div>• •</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	A	535	-	-	-	X
2	EDO	A	536	-	-	-	X

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8986 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate aminotransferase.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	515	Total	C	N	O	P	S	Se	0	6	0
			4176	2678	690	795	1	4	8			
1	B	517	Total	C	N	O	P	S	Se	0	13	0
			4219	2701	695	810	1	4	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	leader sequence	UNP Q5FIG7
B	0	GLY	-	leader sequence	UNP Q5FIG7

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	290	Total	O	0	5
			295	295		
3	B	273	Total	O	0	3
			276	276		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aspartate aminotransferase



• Molecule 1: Aspartate aminotransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.78Å 107.75Å 119.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.92 – 2.15 29.88 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.2 (29.92-2.15) 99.3 (29.88-2.15)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.177 , 0.238 0.181 , 0.233	Depositor DCC
R_{free} test set	3199 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 63132 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8986	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/4242	0.80	3/5740 (0.1%)
1	B	0.74	0/4311	0.85	9/5834 (0.2%)
All	All	0.75	0/8553	0.83	12/11574 (0.1%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	97	GLU	CB-CA-C	-7.04	96.33	110.40
1	B	499	ARG	NE-CZ-NH1	6.43	123.52	120.30
1	B	320[A]	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	B	320[B]	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	A	97	GLU	CB-CA-C	-6.04	98.32	110.40
1	B	65	LYS	CD-CE-NZ	5.80	125.05	111.70
1	B	207	ARG	NE-CZ-NH2	-5.62	117.49	120.30
1	B	50	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	B	340	ASP	CB-CA-C	-5.26	99.88	110.40
1	B	366	ARG	NE-CZ-NH2	-5.12	117.74	120.30
1	A	320[A]	ARG	NE-CZ-NH1	5.11	122.86	120.30
1	A	320[B]	ARG	NE-CZ-NH1	5.11	122.86	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4176	0	4141	59	0
1	B	4219	0	4189	58	0
2	A	16	0	24	2	0
2	B	4	0	6	0	0
3	A	295	0	0	4	0
3	B	276	0	0	3	0
All	All	8986	0	8360	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:LYS:HZ2	1:A:438:THR:HG23	1.21	1.02
1:A:54:VAL:HG13	1:B:54:VAL:HG13	1.05	1.01
1:A:258:LYS:NZ	1:A:438:THR:HG23	1.74	1.00
1:A:54:VAL:CG1	1:B:54:VAL:HG13	1.92	1.00
1:A:54:VAL:HG13	1:B:54:VAL:CG1	1.98	0.89
1:B:424:GLN:HG2	3:B:675:HOH:O	1.81	0.78
1:B:258:LYS:HZ2	1:B:438[A]:THR:HG23	1.53	0.73
1:A:312:LLP:H5'2	1:A:312:LLP:HZ1	1.51	0.73
1:B:258:LYS:NZ	1:B:438[A]:THR:HG23	2.05	0.71
1:B:436:THR:CG2	1:B:438[A]:THR:HG22	2.21	0.70
1:A:68:ILE:HG23	2:A:533:EDO:H21	1.73	0.70
1:A:258:LYS:NZ	1:A:438:THR:CG2	2.54	0.70
1:A:436:THR:CG2	1:A:438:THR:HG22	2.26	0.66
1:A:290:VAL:HG13	1:A:440:THR:HG22	1.77	0.66
1:B:420:GLU:OE2	1:B:424:GLN:NE2	2.29	0.65
1:B:429:MSE:HE2	1:B:514:GLY:HA2	1.78	0.65
1:A:376:LEU:HB3	1:B:173:THR:HG23	1.79	0.65
1:A:200:PRO:HG3	1:A:256:THR:HG23	1.79	0.63
1:B:436:THR:HG22	1:B:438[B]:THR:H	1.64	0.62
1:A:248:ILE:HD12	1:A:281:ILE:HD12	1.81	0.62
1:B:436:THR:HG22	1:B:438[A]:THR:H	1.64	0.61
1:A:173:THR:HG23	1:B:376:LEU:HB3	1.84	0.59
1:B:312:LLP:H5'2	1:B:312:LLP:HZ1	1.67	0.59
1:A:163:GLN:HG3	3:A:610:HOH:O	2.02	0.58
1:B:194:LYS:HG2	1:B:244:ILE:HD13	1.86	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:208:ILE:HB	1:B:211:LEU:HD13	1.86	0.57
1:B:195:ILE:HG12	1:B:216:LEU:HD23	1.87	0.56
1:A:200:PRO:HG3	1:A:256:THR:CG2	2.34	0.56
1:B:151:GLN:HA	1:B:155:TYR:CG	2.40	0.56
1:A:424:GLN:HG3	1:A:511:ALA:HB2	1.88	0.56
1:B:258:LYS:NZ	1:B:438[A]:THR:CG2	2.69	0.55
1:B:151:GLN:HA	1:B:155:TYR:CD2	2.42	0.55
1:B:436:THR:HG21	1:B:438[A]:THR:HG22	1.89	0.55
1:B:200:PRO:HG3	1:B:256[A]:THR:HG23	1.89	0.54
2:A:536:EDO:H12	3:B:694:HOH:O	2.07	0.53
1:B:107:TYR:CD1	1:B:395:MSE:HE1	2.44	0.53
1:B:424:GLN:HG3	1:B:511:ALA:HB2	1.91	0.53
1:B:83:ARG:HG3	1:B:116:ARG:HB3	1.91	0.52
1:A:65:LYS:NZ	3:A:586:HOH:O	2.43	0.51
1:A:436:THR:HG21	1:A:438:THR:HG22	1.92	0.51
1:A:258:LYS:HZ1	1:A:438:THR:CG2	2.23	0.51
1:A:464:LEU:O	1:A:468:PHE:HB2	2.11	0.51
1:B:23:ALA:O	1:B:26[A]:GLN:HG2	2.11	0.51
1:A:420:GLU:OE2	1:A:424:GLN:NE2	2.43	0.50
1:A:424:GLN:HG2	3:A:685:HOH:O	2.11	0.50
1:B:39:ARG:HD3	1:B:312:LLP:H4'2	1.93	0.50
1:B:429:MSE:HG2	1:B:451:LEU:HD21	1.93	0.50
1:B:464:LEU:O	1:B:468:PHE:HB2	2.12	0.49
1:A:39:ARG:HD3	1:A:312:LLP:H4'2	1.94	0.49
1:A:208[B]:ILE:HB	1:A:211:LEU:HD13	1.95	0.49
1:A:19:ARG:HD2	1:A:19:ARG:N	2.27	0.49
1:B:195:ILE:HG13	1:B:197:ILE:HD11	1.93	0.49
1:A:107:TYR:CD1	1:A:395:MSE:HE1	2.47	0.49
1:B:427:ASP:OD1	1:B:433:LYS:CE	2.61	0.49
1:B:290:VAL:HG13	1:B:440:THR:HG22	1.95	0.48
1:A:83:ARG:HG3	1:A:116:ARG:HB3	1.95	0.48
1:A:319:TRP:HB3	1:A:321:LEU:HG	1.96	0.48
1:B:436:THR:HG22	1:B:437:ASP:N	2.29	0.48
1:A:132:TYR:HB2	1:B:33:ILE:HD13	1.96	0.48
1:A:476:LYS:HB3	1:A:520:LEU:HD21	1.94	0.48
1:B:24:LEU:HD13	1:B:471:VAL:HG13	1.96	0.47
1:A:208[B]:ILE:HG23	1:B:373:SER:HB3	1.96	0.47
1:B:51:LEU:HB3	1:B:97:GLU:CG	2.44	0.47
1:B:19:ARG:HD2	1:B:19:ARG:N	2.27	0.47
1:B:317:THR:O	1:B:320[B]:ARG:HD2	2.13	0.47
1:B:436:THR:HG22	1:B:438[A]:THR:HG22	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:241:ASP:HB3	1:A:244:ILE:HG12	1.99	0.45
1:A:269:LYS:O	1:A:273:GLU:HG3	2.16	0.45
1:B:154:SER:HB2	1:B:295:SER:HB2	1.98	0.45
1:A:311:SER:HB3	1:A:317:THR:HG22	1.98	0.45
1:A:137:ARG:O	1:A:168:PRO:HD2	2.17	0.45
1:A:317:THR:O	1:A:320[B]:ARG:HD2	2.17	0.45
1:B:90:LEU:HB3	1:B:98:ASP:HB3	1.99	0.45
1:A:306:LEU:HB3	1:A:325:ALA:HB3	1.99	0.44
1:A:208[A]:ILE:HB	1:A:211:LEU:HD13	2.00	0.44
1:A:129:ALA:HB3	1:B:319:TRP:CD1	2.52	0.44
1:A:132:TYR:HB2	1:B:33:ILE:CD1	2.48	0.44
1:B:28:ASN:HD22	1:B:485:LEU:HB3	1.83	0.43
1:B:258:LYS:NZ	1:B:434:ASP:OD2	2.49	0.43
1:A:205:TYR:O	1:A:208[A]:ILE:HD13	2.17	0.43
1:A:104:ALA:HA	1:A:395:MSE:HE2	2.01	0.43
1:B:312:LLP:OP1	1:B:320[A]:ARG:NH2	2.52	0.43
1:A:151:GLN:HA	1:A:155:TYR:CG	2.54	0.43
1:A:376:LEU:HD12	1:B:177:VAL:HG23	2.00	0.43
1:A:436:THR:HG22	1:A:437:ASP:N	2.33	0.43
3:A:677:HOH:O	1:B:320[A]:ARG:HD3	2.18	0.42
1:A:200:PRO:CG	1:A:256:THR:HG23	2.46	0.42
1:A:17:ILE:O	1:A:18:SER:C	2.58	0.42
1:A:320[A]:ARG:HD3	3:B:577:HOH:O	2.18	0.42
1:A:336:ILE:O	1:A:344:LEU:HD21	2.20	0.42
1:B:253:THR:HG1	1:B:256[B]:THR:HB	1.85	0.42
1:A:286:TYR:OH	1:A:312:LLP:H4'1	2.19	0.42
1:A:194:LYS:HG2	1:A:244:ILE:HD13	2.01	0.41
1:B:487:ASP:O	1:B:505:LEU:HD21	2.20	0.41
1:B:68:ILE:O	1:B:74:ALA:HA	2.19	0.41
1:B:286:TYR:OH	1:B:312:LLP:H4'1	2.21	0.41
1:A:509:ASP:O	1:A:513:ILE:HG12	2.20	0.41
1:B:320[B]:ARG:HA	1:B:320[B]:ARG:NH2	2.35	0.41
1:B:320[B]:ARG:HA	1:B:320[B]:ARG:HH21	1.86	0.41
1:A:246:ALA:HA	1:A:279:MSE:O	2.21	0.41
1:A:194:LYS:HD3	1:A:217:VAL:HG21	2.02	0.41
1:B:427:ASP:OD1	1:B:433:LYS:HE3	2.21	0.41
1:B:195:ILE:HG13	1:B:197:ILE:CD1	2.51	0.41
1:A:469:GLU:HG3	1:A:472:ASP:H	1.86	0.41
1:A:294:LYS:HZ2	1:A:298:SER:CB	2.34	0.40
1:A:255:PRO:HG2	1:A:445:LEU:HD11	2.03	0.40
1:B:195:ILE:HG12	1:B:216:LEU:CD2	2.51	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:444:SER:HG	1:A:502:GLN:HE22	1.66	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	518/533 (97%)	505 (98%)	13 (2%)	0	100	100
1	B	527/533 (99%)	516 (98%)	11 (2%)	0	100	100
All	All	1045/1066 (98%)	1021 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	453/457 (99%)	447 (99%)	6 (1%)	76	82
1	B	462/457 (101%)	459 (99%)	3 (1%)	90	94
All	All	915/914 (100%)	906 (99%)	9 (1%)	82	87

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LEU
1	A	234	ASN
1	A	277	LYS
1	A	341	LYS
1	A	424	GLN
1	A	444	SER
1	B	424	GLN
1	B	444	SER
1	B	450	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	B	28	ASN
1	B	378	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	LLP	A	312	1	23,24,25	1.78	5 (21%)	28,32,34	2.20	7 (25%)
1	LLP	B	312	1	23,24,25	1.66	3 (13%)	28,32,34	2.60	9 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	312	1	-	0/15/17/19	0/1/1/1
1	LLP	B	312	1	-	0/15/17/19	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	312	LLP	O3-C3	-5.33	1.24	1.37
1	B	312	LLP	O3-C3	-4.90	1.25	1.37
1	A	312	LLP	CD-CE	2.00	1.57	1.51
1	B	312	LLP	C2-N1	2.25	1.38	1.34
1	A	312	LLP	C2-N1	2.31	1.39	1.34
1	A	312	LLP	C6-N1	2.68	1.40	1.34
1	A	312	LLP	CE-NZ	2.84	1.52	1.46
1	B	312	LLP	C6-N1	2.95	1.40	1.34

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	312	LLP	C3-C4-C4'	-7.03	111.06	120.16
1	A	312	LLP	C3-C4-C4'	-6.98	111.12	120.16
1	B	312	LLP	CE-NZ-C4'	-5.26	103.79	118.97
1	B	312	LLP	O-C-CA	-2.93	117.85	125.49
1	A	312	LLP	O-C-CA	-2.50	118.98	125.49
1	A	312	LLP	OP4-P-OP1	-2.43	100.96	107.14
1	A	312	LLP	CE-NZ-C4'	-2.23	112.54	118.97
1	B	312	LLP	C5'-C5-C6	-2.20	115.12	119.28
1	B	312	LLP	CD-CE-NZ	2.05	114.34	110.98
1	A	312	LLP	C5'-C5-C4	2.85	126.26	121.47
1	A	312	LLP	C3-C4-C5	3.08	120.41	118.11
1	B	312	LLP	C5'-C5-C4	3.61	127.54	121.47
1	B	312	LLP	C5-C4-C4'	4.04	127.33	121.52
1	B	312	LLP	OP4-C5'-C5	4.47	116.38	108.99
1	B	312	LLP	C3-C4-C5	4.68	121.61	118.11
1	A	312	LLP	C5-C4-C4'	4.83	128.46	121.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	312	LLP	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	312	LLP	4	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	533	-	3,3,3	0.31	0	2,2,2	0.52	0
2	EDO	A	534	-	3,3,3	0.55	0	2,2,2	0.20	0
2	EDO	A	535	-	3,3,3	0.39	0	2,2,2	0.30	0
2	EDO	A	536	-	3,3,3	0.40	0	2,2,2	0.38	0
2	EDO	B	533	-	3,3,3	0.53	0	2,2,2	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	533	-	-	0/1/1/1	0/0/0/0
2	EDO	A	534	-	-	0/1/1/1	0/0/0/0
2	EDO	A	535	-	-	0/1/1/1	0/0/0/0
2	EDO	A	536	-	-	0/1/1/1	0/0/0/0
2	EDO	B	533	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	533	EDO	1	0
2	A	536	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	506/533 (94%)	0.05	24 (4%)	35 46	23, 29, 42, 55	0
1	B	508/533 (95%)	0.04	13 (2%)	59 68	21, 29, 42, 55	0
All	All	1014/1066 (95%)	0.04	37 (3%)	46 57	21, 29, 42, 55	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	524	TYR	5.5
1	A	228	ASP	3.3
1	A	527	GLU	3.1
1	A	437	ASP	3.1
1	B	435	GLU	3.0
1	B	459	GLU	2.9
1	A	463	TYR	2.9
1	B	436	THR	2.9
1	B	437	ASP	2.9
1	B	467	ASN	2.8
1	A	466	ASN	2.8
1	A	528	PHE	2.8
1	A	479	GLU	2.8
1	A	467	ASN	2.8
1	B	531	ASN	2.8
1	A	306	LEU	2.7
1	B	229	TRP	2.7
1	A	226	LYS	2.7
1	A	234	ASN	2.7
1	A	459	GLU	2.6
1	A	15	PHE	2.6
1	B	226	LYS	2.5
1	A	456	TYR	2.4
1	B	80	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	462	ASP	2.3
1	B	403	ASN	2.3
1	A	227	ASN	2.2
1	A	435	GLU	2.2
1	A	307	VAL	2.2
1	A	525	TYR	2.2
1	A	526	GLU	2.1
1	B	458	LYS	2.1
1	B	489	VAL	2.1
1	A	323	VAL	2.1
1	B	232	GLU	2.1
1	A	225	GLU	2.0
1	A	324	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	LLP	B	312	24/25	0.98	0.10	-	28,30,32,33	0
1	LLP	A	312	24/25	0.98	0.14	-	28,31,32,34	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	EDO	A	536	4/4	0.87	0.29	10.95	39,40,42,44	0
2	EDO	A	535	4/4	0.91	0.15	5.39	31,31,34,37	0
2	EDO	A	534	4/4	0.93	0.13	1.81	39,40,41,42	0
2	EDO	B	533	4/4	0.92	0.10	-0.12	40,41,42,42	0
2	EDO	A	533	4/4	0.95	0.17	-	40,41,42,45	0

6.5 Other polymers ⓘ

There are no such residues in this entry.