



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:21 PM GMT

PDB ID : 1F8X
Title : CRYSTAL STRUCTURE OF NUCLEOSIDE 2-DEOXYRIBOSYLTRANSFERASE
Authors : Armstrong, S.R.; Cook, W.J.; Short, S.A.; Ealick, S.E.
Deposited on : 2000-07-05
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	157	 81% 18% ..
1	B	157	 78% 18% . ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2616 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NUCLEOSIDE 2-DEOXYRIBOSYLTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	156	Total	C	N	O	S	0	0	0
			1268	816	201	246	5			
1	B	156	Total	C	N	O	S	0	0	0
			1268	816	201	246	5			

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	49	Total	O	0	0
			49	49		
2	B	31	Total	O	0	0
			31	31		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

• Molecule 1: NUCLEOSIDE 2-DEOXYRIBOSYLTRANSFERASE

Chain A: 



• Molecule 1: NUCLEOSIDE 2-DEOXYRIBOSYLTRANSFERASE

Chain B: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	151.00Å 151.00Å 151.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	5.00 – 2.50	Depositor
% Data completeness (in resolution range)	(Not available) (5.00-2.50)	Depositor
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.166 , 0.240	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2616	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/1301	0.64	1/1762 (0.1%)
1	B	0.37	0/1301	0.70	2/1762 (0.1%)
All	All	0.39	0/2602	0.67	3/3524 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	316	GLU	CA-CB-CG	-8.69	94.29	113.40
1	B	256	PRO	N-CA-C	5.79	127.16	112.10
1	A	7	TYR	N-CA-C	-5.01	97.47	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1268	0	1216	19	0
1	B	1268	0	1216	33	0
2	A	49	0	0	3	0
2	B	31	0	0	0	0
All	All	2616	0	2432	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:206:ILE:HG22	1:B:281:ILE:HB	1.34	1.10
1:B:250:ILE:HA	1:B:255:HIS:HE1	1.33	0.93
1:B:250:ILE:HA	1:B:255:HIS:CE1	2.09	0.87
1:B:250:ILE:CA	1:B:255:HIS:HE1	1.98	0.75
1:A:4:LYS:HE2	2:A:833:HOH:O	1.90	0.71
1:B:277:LYS:HA	1:B:307:LYS:HE2	1.73	0.70
1:B:336:MET:HE1	1:B:339:LEU:HD12	1.74	0.68
1:B:250:ILE:C	1:B:255:HIS:HE1	1.97	0.67
1:B:254:GLU:HB2	1:B:255:HIS:CD2	2.37	0.60
1:B:251:ARG:N	1:B:255:HIS:CE1	2.71	0.58
1:B:250:ILE:C	1:B:255:HIS:CE1	2.78	0.56
1:B:251:ARG:N	1:B:255:HIS:HE1	2.03	0.56
1:A:50:ILE:HD13	1:A:55:HIS:HB2	1.86	0.56
1:B:297:MET:HE1	1:B:301:TYR:HB2	1.88	0.55
1:B:255:HIS:N	1:B:256:PRO:HD3	2.23	0.53
1:A:13:PHE:H	1:A:17:GLN:NE2	2.07	0.52
1:B:217:GLN:HA	1:B:287:ILE:HD11	1.92	0.51
1:B:250:ILE:HG21	1:B:258:TYR:CZ	2.46	0.50
1:B:217:GLN:HA	1:B:287:ILE:CD1	2.42	0.50
1:B:336:MET:CE	1:B:339:LEU:HD12	2.40	0.50
1:A:133:VAL:HG21	1:A:152:TYR:CD2	2.47	0.50
1:B:250:ILE:HG21	1:B:258:TYR:CE2	2.47	0.50
1:B:347:PRO:O	1:B:348:ARG:HD2	2.12	0.49
1:A:122:ILE:O	1:A:156:VAL:HA	2.12	0.49
1:B:276:ILE:O	1:B:307:LYS:HE3	2.13	0.49
1:A:95:LEU:O	1:A:99:LEU:HG	2.12	0.49
1:B:303:LEU:HD12	1:B:329:VAL:HG23	1.95	0.48
1:A:103:LEU:HA	2:A:852:HOH:O	2.14	0.48
1:A:132:ASN:ND2	1:A:150:ASP:HB3	2.29	0.48
1:A:34:ILE:HD11	1:A:144:PHE:CD1	2.49	0.47
1:B:316:GLU:H	1:B:316:GLU:HG2	1.02	0.46
1:A:19:LYS:O	1:A:23:GLU:HG3	2.15	0.46
1:B:357:TYR:C	1:B:357:TYR:CD1	2.89	0.45
1:A:31:ASN:HA	1:A:32:PRO:HD3	1.82	0.45
1:A:91:GLU:HA	2:A:807:HOH:O	2.16	0.44
1:B:251:ARG:HE	1:B:254:GLU:HG3	1.83	0.44
1:A:104:SER:CB	1:B:304:SER:HB3	2.47	0.44
1:A:156:VAL:O	1:B:259:LEU:HD22	2.17	0.44
1:B:347:PRO:C	1:B:348:ARG:HD2	2.39	0.43
1:B:255:HIS:N	1:B:255:HIS:CD2	2.86	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:TRP:HB2	1:A:18:ASN:HD21	1.82	0.43
1:B:224:ALA:HA	1:B:336:MET:HE1	2.00	0.42
1:B:252:VAL:HG23	1:B:253:ASP:N	2.35	0.42
1:A:50:ILE:HG13	1:A:50:ILE:O	2.20	0.41
1:A:94:GLY:O	1:A:98:GLU:HG3	2.20	0.41
1:B:250:ILE:CA	1:B:255:HIS:CE1	2.84	0.41
1:A:104:SER:HB3	1:B:304:SER:HB3	2.02	0.41
1:A:74:ASN:HA	1:A:77:LYS:HG2	2.02	0.41
1:B:321:PRO:HA	1:B:355:ALA:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	154/157 (98%)	149 (97%)	5 (3%)	0	100	100
1	B	154/157 (98%)	139 (90%)	14 (9%)	1 (1%)	30	50
All	All	308/314 (98%)	288 (94%)	19 (6%)	1 (0%)	46	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	256	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	136/137 (99%)	135 (99%)	1 (1%)	88	97
1	B	136/137 (99%)	127 (93%)	9 (7%)	21	38
All	All	272/274 (99%)	262 (96%)	10 (4%)	41	68

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ILE
1	B	221	TYR
1	B	241	VAL
1	B	255	HIS
1	B	262	LYS
1	B	297	MET
1	B	307	LYS
1	B	316	GLU
1	B	324	LEU
1	B	339	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	GLN
1	A	18	ASN
1	A	38	ASN
1	A	46	GLN
1	A	70	ASN
1	A	74	ASN
1	A	132	ASN
1	B	238	ASN
1	B	255	HIS
1	B	260	HIS
1	B	274	ASN
1	B	279	ASN
1	B	345	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.