



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 02:35 PM BST

PDB ID : 1F8Z  
Title : NMR STRUCTURE OF THE SIXTH LIGAND-BINDING MODULE OF  
THE LDL RECEPTOR  
Authors : Clayton, D.J.; Brereton, I.M.; Kroon, P.A.; Smith, R.  
Deposited on : 2000-07-05

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	39	<div> <div>59%</div> <div>21%</div> <div>21%</div> </div>

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 13 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:4-A:34 (31)	0.17	13

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 5, 6, 7, 8, 9, 13, 14, 18, 19, 20
2	3, 4, 12, 16
3	10, 17
4	11, 15
Single-model clusters	2

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 552 atoms, of which 252 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called LOW-DENSITY LIPOPROTEIN RECEPTOR.

Mol	Chain	Residues	Atoms							Trace
1	A	39	Total	C	H	N	O	S		0
			551	171	252	55	66	7		

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	
2	A	1	Total	Ca
			1	1




### 4.2.3 Score per residue for model 3

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR



### 4.2.4 Score per residue for model 4

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR




### 4.2.5 Score per residue for model 5

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR




### 4.2.6 Score per residue for model 6

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR




### 4.2.7 Score per residue for model 7

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR



### 4.2.8 Score per residue for model 8

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR




### 4.2.9 Score per residue for model 9

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR




### 4.2.10 Score per residue for model 10

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR



### 4.2.11 Score per residue for model 11

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR




### 4.2.12 Score per residue for model 12

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR




#### 4.2.13 Score per residue for model 13 (medoid)

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR



#### 4.2.14 Score per residue for model 14

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR




#### 4.2.15 Score per residue for model 15

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR



#### 4.2.16 Score per residue for model 16

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR



#### 4.2.17 Score per residue for model 17


- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR





### 4.2.18 Score per residue for model 18

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR




### 4.2.19 Score per residue for model 19

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR



### 4.2.20 Score per residue for model 20

- Molecule 1: LOW-DENSITY LIPOPROTEIN RECEPTOR



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *Structure determination by Torsion angle dynamics, Molecular dynamics.*

Of the 50 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	1.5
X-PLOR	structure solution	3.1
X-PLOR	refinement	3.1

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.23±0.01	0±0/253 (0.0±0.0%)	1.40±0.02	1±1/338 (0.1±0.2%)
All	All	1.23	0/5060 (0.0%)	1.40	10/6760 (0.1%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	20	ARG	NE-CZ-NH2	-6.10	117.25	120.30	6	2
1	A	4	ARG	NE-CZ-NH2	-5.46	117.57	120.30	8	5
1	A	24	ARG	NE-CZ-NH2	-5.09	117.75	120.30	1	3

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	249	205	205	2±1
All	All	5000	4100	4100	42

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:16:ILE:HD13	1:A:33:ASP:HB2	0.73	1.61	5	20
1:A:14:ASN:HB3	1:A:28:CYS:SG	0.53	2.44	20	9
1:A:10:CYS:SG	1:A:33:ASP:HB3	0.51	2.45	8	7
1:A:14:ASN:O	1:A:28:CYS:SG	0.47	2.73	17	3
1:A:25:GLU:O	1:A:26:TYR:C	0.41	2.58	15	2
1:A:21:GLN:HG2	1:A:22:CYS:SG	0.40	2.55	10	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	31/39 (79%)	26±1 (85±2%)	4±1 (12±2%)	1±0 (3±1%)	8	38
All	All	620/780 (79%)	524 (85%)	75 (12%)	21 (3%)	8	38

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	26	TYR	20
1	A	31	MET	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	29/35 (83%)	26±1 (89±4%)	3±1 (11±4%)	12	55
All	All	580/700 (83%)	515 (89%)	65 (11%)	12	55

All 15 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	29	LYS	18
1	A	23	ASP	14
1	A	24	ARG	8
1	A	20	ARG	6
1	A	12	ASP	2
1	A	25	GLU	2
1	A	33	ASP	2
1	A	14	ASN	2
1	A	30	ASP	2
1	A	9	GLN	2
1	A	31	MET	2
1	A	4	ARG	2
1	A	21	GLN	1
1	A	32	SER	1
1	A	27	ASP	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided