



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:27 PM GMT

PDB ID : 1FO4
Title : CRYSTAL STRUCTURE OF XANTHINE DEHYDROGENASE ISOLATED FROM BOVINE MILK
Authors : Enroth, C.; Eger, B.T.; Okamoto, K.; Nishino, T.; Nishino, T.; Pai, E.F.
Deposited on : 2000-08-24
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	1332	
1	B	1332	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MOS	A	3004	-	-	X	-
5	MOS	B	4004	-	-	X	-
7	FAD	A	3006	X	-	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	FAD	B	4006	X	-	-	-
8	GOL	A	3007	-	X	X	-
8	GOL	A	3008	-	X	-	-
8	GOL	B	4007	-	X	X	-
8	GOL	B	4008	-	X	X	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 22402 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called XANTHINE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1299	Total	C	N	O	S	0	0	0
			10077	6404	1728	1884	61			
1	B	1296	Total	C	N	O	S	0	0	0
			10054	6391	1724	1878	61			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Fe S 4 2 2	0	0
3	A	1	Total Fe S 4 2 2	0	0
3	B	1	Total Fe S 4 2 2	0	0
3	B	1	Total Fe S 4 2 2	0	0

- Molecule 4 is PHOSPHONIC ACIDMONO-(2-AMINO-5,6-DIMERCAPTO-4-OXO-3,7,8A, 9,10,10A-HEXAHYDRO-4H-8-OXA-1,3,9,10-TETRAAZA-ANTHRACEN-7-YLMETHYL) ESTER (three-letter code: MTE) (formula: C₁₀H₁₄N₅O₆P₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O P S 24 10 5 6 1 2	0	0
4	B	1	Total C N O P S 24 10 5 6 1 2	0	0

- Molecule 5 is DIOXOTHIO MOLYBDENUM(VI) ION (three-letter code: MOS) (formula: HMoO₂S).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	Mo	O	S	0	0
			4	1	2	1		
5	B	1	Total	Mo	O	S	0	0
			4	1	2	1		

- Molecule 6 is 2-HYDROXYBENZOIC ACID (three-letter code: SAL) (formula: $C_7H_6O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			10	7	3		
6	B	1	Total	C	O	0	0
			10	7	3		

- Molecule 7 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
7	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		
8	B	1	Total	C	O	0	0
			6	3	3		
8	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1047	Total	O	0	0
			1047	1047		
9	B	1000	Total	O	0	0
			1000	1000		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: XANTHINE DEHYDROGENASE



• Molecule 1: XANTHINE DEHYDROGENASE



L1054	G837	D691	R508	L313	GLY
K1055	G838	D691	R509	E314	GLY
P1072	B840	I695	T512	R328	ASN
V1081	A844	I698	L527	L331	ASN
S1082	K847	E699	G528	E332	PRO
I1085	K852	D700	LYS		ASN
Q1088	N705	I705	ASP	R335	CYS
A1093	S706	S706	SER	W336	MET
	F707	F707	LYS	F337	ASN
	E860		ASP	A338	GLN
			LYS	G339	LYS
	E879	K713	CYS	K340	LYS
I1097	R880	I714	GLY		ASP
			E537	A346	HIS
N1108	M885	L719	L538		THR
P1109	K720	K720		N351	VAL
	K721	K721	L548		THR
R1124	R895	G722	H552	L361	
V1125	G896	F723			
S1126		S724		I374	
T1129	R899	E725	N565	V375	N197
	I900	A726		S376	
	C901	D727	R569	R377	E200
V1133				G378	E209
R1134	T909	Q739	A586	T379	P210
G1139	F911	H741	D595	R394	
T1301	R912	F742	R598	L398	L217
P1302	G913	Y743	V599	L219	R218
	F914	L744	E600	E402	K220
V1310	G915				
D1311	Q918	H747	R606	I407	G231
K1312	K1172	T752	L607		
F1313	N1173		V608	R427	T241
	L1174	E757	T609	E428	L242
	R1175			D429	K243
L1316	S1184	E761	A615	L441	E244
C1317	S1185		K616		
V1318	L1186	F955			
A1321	P1322	V964	S619	M447	K256
G1323	P1187	P965		E451	L257
N1324	A1189	R786	V628	Q471	V258
C1325	I1190	I787	F634	K472	L264
K1326	D1191	L788		Q473	F275
	I1192		E645		
	F987	G797		K476	I279
L1330	V1195	E902	N650		C280
R1331	E1196		D651	E480	P281
V1332		V810		K461	L284
	T1207	P822	T659	L482	
	L1208			L483	L287
	T1221	M833	V670	Q484	
	Y1227	S1008	E679	D485	K310
		A1017	R835	V486	T311
			T326		L312

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	169.45Å 124.49Å 148.33Å 90.00° 90.94° 90.00°	Depositor
Resolution (Å)	25.00 – 2.10	Depositor
% Data completeness (in resolution range)	86.9 (25.00-2.10)	Depositor
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.198 , 0.238	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	22402	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SAL, MOS, CA, FES, FAD, MTE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/10298	0.64	1/13939 (0.0%)
1	B	0.37	0/10275	0.64	1/13909 (0.0%)
All	All	0.37	0/20573	0.64	2/27848 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1191	ASP	N-CA-C	-5.23	96.89	111.00
1	B	243	LYS	N-CA-C	-5.00	97.49	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10077	0	10075	203	0
1	B	10054	0	10054	181	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	8	0	0	0	0
3	B	8	0	0	0	0
4	A	24	0	10	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	24	0	10	2	0
5	A	4	0	0	2	0
5	B	4	0	0	2	0
6	A	10	0	4	0	0
6	B	10	0	4	1	0
7	A	53	0	29	3	0
7	B	53	0	30	2	0
8	A	12	0	5	7	0
8	B	12	0	7	11	0
9	A	1047	0	0	21	0
9	B	1000	0	0	11	0
All	All	22402	0	20228	386	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 386 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:3008:GOL:C1	8:A:3008:GOL:O1	1.68	1.39
8:B:4008:GOL:C1	8:B:4008:GOL:O1	1.71	1.36
1:A:3:ALA:HB1	1:A:228:ARG:H	1.19	1.07
1:B:645:GLU:HG2	1:B:650:ASN:HD22	1.19	1.07
5:B:4004:MOS:MO	5:B:4004:MOS:S	1.66	1.06

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1293/1332 (97%)	1228 (95%)	54 (4%)	11 (1%)	21 15

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	1290/1332 (97%)	1236 (96%)	45 (4%)	9 (1%)	26	21
All	All	2583/2664 (97%)	2464 (95%)	99 (4%)	20 (1%)	24	17

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1008	SER
1	A	1331	ARG
1	B	244	GLU
1	B	1008	SER
1	B	1287	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1101/1128 (98%)	1084 (98%)	17 (2%)	72	78
1	B	1098/1128 (97%)	1078 (98%)	20 (2%)	66	72
All	All	2199/2256 (98%)	2162 (98%)	37 (2%)	68	74

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1325	CYS
1	B	328	ARG
1	B	1325	CYS
1	B	100	PRO
1	B	129	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1284	GLN
1	A	1324	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	650	ASN
1	A	1212	HIS
1	B	1145	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 18 ligands modelled in this entry, 2 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FES	A	3001	1	0,4,4	0.00	-	0,4,4	0.00	-
3	FES	A	3002	1	0,4,4	0.00	-	0,4,4	0.00	-
4	MTE	A	3003	5	19,26,26	7.40	15 (78%)	19,40,40	3.01	7 (36%)
5	MOS	A	3004	4	0,3,3	0.00	-	0,3,3	0.00	-
6	SAL	A	3005	-	7,10,10	1.70	3 (42%)	10,13,13	1.21	0
7	FAD	A	3006	-	48,58,58	4.62	30 (62%)	54,89,89	2.87	19 (35%)
8	GOL	A	3007	-	5,5,5	7.28	4 (80%)	5,5,5	6.01	4 (80%)
8	GOL	A	3008	-	5,5,5	4.57	4 (80%)	5,5,5	5.95	3 (60%)
3	FES	B	4001	1	0,4,4	0.00	-	0,4,4	0.00	-
3	FES	B	4002	1	0,4,4	0.00	-	0,4,4	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MTE	B	4003	5	19,26,26	7.51	12 (63%)	19,40,40	2.97	7 (36%)
5	MOS	B	4004	4	0,3,3	0.00	-	0,3,3	0.00	-
6	SAL	B	4005	-	7,10,10	1.67	3 (42%)	10,13,13	1.33	1 (10%)
7	FAD	B	4006	-	48,58,58	4.41	32 (66%)	54,89,89	2.90	19 (35%)
8	GOL	B	4007	-	5,5,5	6.84	5 (100%)	5,5,5	5.92	4 (80%)
8	GOL	B	4008	-	5,5,5	4.43	3 (60%)	5,5,5	6.08	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FES	A	3001	1	-	0/0/4/4	0/1/1/1
3	FES	A	3002	1	-	0/0/4/4	0/1/1/1
4	MTE	A	3003	5	-	0/6/34/34	0/3/3/3
5	MOS	A	3004	4	-	0/0/0/0	0/0/0/0
6	SAL	A	3005	-	-	0/0/4/4	0/1/1/1
7	FAD	A	3006	-	2/2/9/9	0/30/50/50	0/6/6/6
8	GOL	A	3007	-	-	0/4/4/4	0/0/0/0
8	GOL	A	3008	-	-	0/4/4/4	0/0/0/0
3	FES	B	4001	1	-	0/0/4/4	0/1/1/1
3	FES	B	4002	1	-	0/0/4/4	0/1/1/1
4	MTE	B	4003	5	-	0/6/34/34	0/3/3/3
5	MOS	B	4004	4	-	0/0/0/0	0/0/0/0
6	SAL	B	4005	-	-	0/0/4/4	0/1/1/1
7	FAD	B	4006	-	2/2/9/9	0/30/50/50	0/6/6/6
8	GOL	B	4007	-	-	0/4/4/4	0/0/0/0
8	GOL	B	4008	-	-	0/4/4/4	0/0/0/0

The worst 5 of 111 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	3007	GOL	C3-C2	-13.59	1.00	1.52
8	B	4007	GOL	C3-C2	-12.68	1.03	1.52
7	A	3006	FAD	C5'-C4'	-8.62	1.38	1.51
8	A	3008	GOL	C3-C2	-7.31	1.24	1.52
7	B	4006	FAD	C5'-C4'	-7.17	1.40	1.51

The worst 5 of 67 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	4006	FAD	C4-C4X-C10	-7.92	114.87	119.94
7	A	3006	FAD	C4-C4X-C10	-7.76	114.98	119.94
7	A	3006	FAD	N3A-C2A-N1A	-7.51	123.14	128.89
7	B	4006	FAD	O5B-PA-O1A	-6.05	86.14	109.62
7	B	4006	FAD	C4X-C10-N10	-5.71	117.16	120.52

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
7	B	4006	FAD	C2'
7	B	4006	FAD	C3'
7	A	3006	FAD	C2'
7	A	3006	FAD	C3'

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 31 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	3003	MTE	1	0
5	A	3004	MOS	2	0
7	A	3006	FAD	3	0
8	A	3007	GOL	4	0
8	A	3008	GOL	3	0
4	B	4003	MTE	2	0
5	B	4004	MOS	2	0
6	B	4005	SAL	1	0
7	B	4006	FAD	2	0
8	B	4007	GOL	7	0
8	B	4008	GOL	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.