



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:27 PM GMT

PDB ID : 1FOR
Title : STRUCTURE DETERMINATION OF AN FAB FRAGMENT THAT NEUTRALIZES HUMAN RHINOVIRUS AND ANALYSIS OF THE FAB-VIRUS COMPLEX
Authors : Smith, T.J.; Liu, H.
Deposited on : 1994-05-24
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	210	 80% 16% •
2	H	219	 62% 35% •

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3416 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGG2A-KAPPA 17-IA FAB (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	210	Total	C	N	O	S	0	0	0
			1619	1011	271	329	8			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	14	PHE	SER	CONFLICT	EMBL X79906
L	15	PRO	LEU	CONFLICT	EMBL X79906
L	18	LYS	ARG	CONFLICT	EMBL X79906
L	21	ILE	MET	CONFLICT	EMBL X79906
L	24	SER	THR	CONFLICT	EMBL X79906
L	26	THR	SER	CONFLICT	EMBL X79906
L	?	-	SER	DELETION	EMBL X79906
L	?	-	SER	DELETION	EMBL X79906
L	30	ASN	SER	CONFLICT	EMBL X79906
L	32	MET	LEU	CONFLICT	EMBL X79906
L	35	PHE	TYR	CONFLICT	EMBL X79906
L	41	THR	SER	CONFLICT	EMBL X79906
L	50	SER	THR	CONFLICT	EMBL X79906
L	76	ARG	SER	CONFLICT	EMBL X79906
L	88	GLN	HIS	CONFLICT	EMBL X79906
L	90	ARG	TYR	CONFLICT	EMBL X79906
L	91	SER	HIS	CONFLICT	EMBL X79906
L	92	SER	ARG	CONFLICT	EMBL X79906
L	93	TYR	PHE	CONFLICT	EMBL X79906
L	95	ILE	HIS	CONFLICT	EMBL X79906
L	99	SER	GLY	CONFLICT	EMBL X79906

- Molecule 2 is a protein called IGG2A-KAPPA 17-IA FAB (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	219	Total	C	N	O	S	0	0	0
			1640	1037	271	325	7			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	2	GLY	ILE	CONFLICT	PIR S38950
H	9	ALA	PRO	CONFLICT	PIR S38950
H	16	SER	ALA	CONFLICT	PIR S38950
H	28	ALA	THR	CONFLICT	PIR S38950
H	30	SER	THR	CONFLICT	PIR S38950
H	31	SER	ASP	CONFLICT	PIR S38950
H	32	PHE	TYR	CONFLICT	PIR S38950
H	33	TRP	TYR	CONFLICT	PIR S38950
H	34	VAL	ILE	CONFLICT	PIR S38950
H	35	ASN	HIS	CONFLICT	PIR S38950
H	43	GLN	GLU	CONFLICT	PIR S38950
H	50	GLN	TRP	CONFLICT	PIR S38950
H	55	ASP	SER	CONFLICT	PIR S38950
H	57	ASP	ASN	CONFLICT	PIR S38950
H	58	ASN	THR	CONFLICT	PIR S38950
H	62	GLY	GLU	CONFLICT	PIR S38950
H	72	ALA	VAL	CONFLICT	PIR S38950
H	74	LYS	THR	CONFLICT	PIR S38950
H	77	THR	SER	CONFLICT	PIR S38950
H	84	TYR	SER	CONFLICT	PIR S38950
H	99	SER	GLY	CONFLICT	PIR S38950
H	101	ASN	-	INSERTION	PIR S38950
H	102	TYR	-	INSERTION	PIR S38950
H	103	PRO	LYS	CONFLICT	PIR S38950
H	104	TYR	PHE	CONFLICT	PIR S38950
H	136	GLY	ASP	CONFLICT	PIR S38950
H	179	GLY	ASP	CONFLICT	PIR S38950
H	198	THR	SER	CONFLICT	PIR S38950

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	82	Total	O	0	0
			82	82		
3	L	75	Total	O	0	0
			75	75		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: IGG2A-KAPPA 17-IA FAB (LIGHT CHAIN)



• Molecule 2: IGG2A-KAPPA 17-IA FAB (HEAVY CHAIN)



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	37.80 Å 97.30 Å 129.50 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.75	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-2.75)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ, X-PLOR	Depositor
R, R_{free}	0.174 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3416	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.39	0/1658	0.67	0/2251
2	H	0.43	0/1684	0.73	2/2296 (0.1%)
All	All	0.41	0/3342	0.70	2/4547 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	H	150	GLY	N-CA-C	5.29	126.33	113.10
2	H	194	TRP	N-CA-C	5.01	124.54	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1619	0	1557	32	0
2	H	1640	0	1593	63	0
3	H	82	0	0	10	0
3	L	75	0	0	2	0
All	All	3416	0	3150	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 88 close contacts within the same asymmetric unit are listed below, sorted by their

clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:51:ILE:HG22	2:H:58:ASN:HB3	1.52	0.90
2:H:154:GLU:HG2	3:H:360:HOH:O	1.82	0.78
2:H:30:SER:HA	2:H:53:PRO:HG2	1.65	0.76
2:H:37:VAL:HG11	2:H:109:TRP:HZ3	1.54	0.73
1:L:135:LEU:HD23	1:L:174:MET:HG2	1.71	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	208/210 (99%)	196 (94%)	11 (5%)	1 (0%)	34	67
2	H	217/219 (99%)	188 (87%)	21 (10%)	8 (4%)	4	12
All	All	425/429 (99%)	384 (90%)	32 (8%)	9 (2%)	9	25

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	102	TYR
2	H	194	TRP
2	H	134	CYS
1	L	139	TYR
2	H	3	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	185/185 (100%)	172 (93%)	13 (7%)	19	44
2	H	182/182 (100%)	167 (92%)	15 (8%)	14	35
All	All	367/367 (100%)	339 (92%)	28 (8%)	16	39

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	210	ARG
2	H	20	ILE
2	H	190	THR
2	H	7	SER
2	H	12	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	L	144	ASN
2	H	6	GLN
1	L	189	ASN
1	L	136	ASN
1	L	160	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.