



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 11:16 PM GMT

PDB ID : 5FPY
Title : Structure of hepatitis C virus (HCV) full-length NS3 complex with small-molecule ligand 5-bromo-1-methyl-1H-indole-2-carboxylic acid (AT21457) in an alternate binding site.
Authors : Davies, T.G.; Jhoti, H.; Ludlow, R.F.; Saini, H.K.; Tickle, I.J.; Verdonk, M.
Deposited on : 2015-12-03
Resolution : 2.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4241 (2.54-2.50)
Clashscore	102246	4968 (2.54-2.50)
Ramachandran outliers	100387	4873 (2.54-2.50)
Sidechain outliers	100360	4875 (2.54-2.50)
RSRZ outliers	91569	4253 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	666	
1	B	666	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	R2N	A	1721	-	-	-	X
2	R2N	B	1721	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10186 atoms, of which 14 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERINE PROTEASE NS3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	645	Total	C	N	O	S	0	0	1
			4700	2969	800	901	30			
1	B	642	Total	C	N	O	S	0	0	0
			4716	2984	802	900	30			

There are 82 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	684	MET	-	EXPRESSION TAG	UNP P26663
A	685	GLY	-	EXPRESSION TAG	UNP P26663
A	686	SER	-	EXPRESSION TAG	UNP P26663
A	687	SER	-	EXPRESSION TAG	UNP P26663
A	688	HIS	-	EXPRESSION TAG	UNP P26663
A	689	HIS	-	EXPRESSION TAG	UNP P26663
A	690	HIS	-	EXPRESSION TAG	UNP P26663
A	691	HIS	-	EXPRESSION TAG	UNP P26663
A	692	HIS	-	EXPRESSION TAG	UNP P26663
A	693	HIS	-	EXPRESSION TAG	UNP P26663
A	694	SER	-	EXPRESSION TAG	UNP P26663
A	695	SER	-	EXPRESSION TAG	UNP P26663
A	696	GLY	-	EXPRESSION TAG	UNP P26663
A	697	LEU	-	EXPRESSION TAG	UNP P26663
A	698	VAL	-	EXPRESSION TAG	UNP P26663
A	699	PRO	-	EXPRESSION TAG	UNP P26663
A	700	ARG	-	EXPRESSION TAG	UNP P26663
A	701	GLY	-	EXPRESSION TAG	UNP P26663
A	702	SER	-	EXPRESSION TAG	UNP P26663
A	703	HIS	-	EXPRESSION TAG	UNP P26663
A	704	MET	-	EXPRESSION TAG	UNP P26663
A	705	GLY	-	EXPRESSION TAG	UNP P26663
A	706	SER	-	EXPRESSION TAG	UNP P26663
A	707	VAL	-	EXPRESSION TAG	UNP P26663
A	708	VAL	-	EXPRESSION TAG	UNP P26663

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	709	ILE	-	EXPRESSION TAG	UNP P26663
A	710	VAL	-	EXPRESSION TAG	UNP P26663
A	711	GLY	-	EXPRESSION TAG	UNP P26663
A	712	ARG	-	EXPRESSION TAG	UNP P26663
A	713	ILE	-	EXPRESSION TAG	UNP P26663
A	714	ILE	-	EXPRESSION TAG	UNP P26663
A	715	LEU	-	EXPRESSION TAG	UNP P26663
A	716	SER	-	EXPRESSION TAG	UNP P26663
A	717	GLY	-	EXPRESSION TAG	UNP P26663
A	718	SER	-	EXPRESSION TAG	UNP P26663
A	719	GLY	-	EXPRESSION TAG	UNP P26663
A	720	SER	-	EXPRESSION TAG	UNP P26663
A	66	GLY	ALA	CONFLICT	UNP P26663
A	86	GLN	PRO	CONFLICT	UNP P26663
A	87	ALA	LYS	CONFLICT	UNP P26663
A	147	SER	PHE	CONFLICT	UNP P26663
B	684	MET	-	EXPRESSION TAG	UNP P26663
B	685	GLY	-	EXPRESSION TAG	UNP P26663
B	686	SER	-	EXPRESSION TAG	UNP P26663
B	687	SER	-	EXPRESSION TAG	UNP P26663
B	688	HIS	-	EXPRESSION TAG	UNP P26663
B	689	HIS	-	EXPRESSION TAG	UNP P26663
B	690	HIS	-	EXPRESSION TAG	UNP P26663
B	691	HIS	-	EXPRESSION TAG	UNP P26663
B	692	HIS	-	EXPRESSION TAG	UNP P26663
B	693	HIS	-	EXPRESSION TAG	UNP P26663
B	694	SER	-	EXPRESSION TAG	UNP P26663
B	695	SER	-	EXPRESSION TAG	UNP P26663
B	696	GLY	-	EXPRESSION TAG	UNP P26663
B	697	LEU	-	EXPRESSION TAG	UNP P26663
B	698	VAL	-	EXPRESSION TAG	UNP P26663
B	699	PRO	-	EXPRESSION TAG	UNP P26663
B	700	ARG	-	EXPRESSION TAG	UNP P26663
B	701	GLY	-	EXPRESSION TAG	UNP P26663
B	702	SER	-	EXPRESSION TAG	UNP P26663
B	703	HIS	-	EXPRESSION TAG	UNP P26663
B	704	MET	-	EXPRESSION TAG	UNP P26663
B	705	GLY	-	EXPRESSION TAG	UNP P26663
B	706	SER	-	EXPRESSION TAG	UNP P26663
B	707	VAL	-	EXPRESSION TAG	UNP P26663
B	708	VAL	-	EXPRESSION TAG	UNP P26663
B	709	ILE	-	EXPRESSION TAG	UNP P26663

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	710	VAL	-	EXPRESSION TAG	UNP P26663
B	711	GLY	-	EXPRESSION TAG	UNP P26663
B	712	ARG	-	EXPRESSION TAG	UNP P26663
B	713	ILE	-	EXPRESSION TAG	UNP P26663
B	714	ILE	-	EXPRESSION TAG	UNP P26663
B	715	LEU	-	EXPRESSION TAG	UNP P26663
B	716	SER	-	EXPRESSION TAG	UNP P26663
B	717	GLY	-	EXPRESSION TAG	UNP P26663
B	718	SER	-	EXPRESSION TAG	UNP P26663
B	719	GLY	-	EXPRESSION TAG	UNP P26663
B	720	SER	-	EXPRESSION TAG	UNP P26663
B	66	GLY	ALA	CONFLICT	UNP P26663
B	86	GLN	PRO	CONFLICT	UNP P26663
B	87	ALA	LYS	CONFLICT	UNP P26663
B	147	SER	PHE	CONFLICT	UNP P26663

- Molecule 2 is 5-BROMO-1-METHYL-1H-INDOLE-2-CARBOXYLIC ACID (three-letter code: R2N) (formula: C₁₀H₈BrNO₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	Br	C	H	N	O	0	0
			21	1	10	7	1	2		
2	B	1	Total	Br	C	H	N	O	0	0
			21	1	10	7	1	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	403	Total 403	O 403	0	0
3	B	325	Total 325	O 325	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SERINE PROTEASE NS3



• Molecule 1: SERINE PROTEASE NS3



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.90Å 109.77Å 142.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.99 – 2.52 46.99 – 2.52	Depositor EDS
% Data completeness (in resolution range)	95.3 (46.99-2.52) 95.3 (46.99-2.52)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.163 , 0.241 0.181 , 0.246	Depositor DCC
R_{free} test set	2354 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	38.6	Xtriage
Anisotropy	0.772	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 72.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 46396 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10186	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: R2N

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/4807	0.68	0/6579
1	B	0.52	0/4823	0.67	0/6594
All	All	0.53	0/9630	0.68	0/13173

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4700	0	4558	24	0
1	B	4716	0	4613	22	0
2	A	14	7	7	0	0
2	B	14	7	7	0	0
3	A	403	0	0	0	0
3	B	325	0	0	0	0
All	All	10172	14	9185	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:596:THR:HG22	1:A:607:ASN:HD22	1.36	0.90
1:A:82:LEU:HG	1:A:170:VAL:HG11	1.63	0.80
1:B:333:HIS:HD2	1:B:335:ASN:H	1.41	0.68
1:A:596:THR:HG22	1:A:607:ASN:ND2	2.16	0.57
1:A:251:ASN:HD21	1:A:262:GLY:H	1.52	0.57
1:B:82:LEU:HG	1:B:170:VAL:HG11	1.87	0.57
1:B:294:SER:HB3	1:B:299:THR:HG21	1.87	0.56
1:B:229:ASN:HD22	1:B:231:SER:H	1.55	0.54
1:B:336:ILE:HD13	1:B:462:ARG:HG2	1.90	0.54
1:A:365:LEU:HD13	1:A:408:VAL:HG23	1.89	0.53
1:A:363:ARG:HG2	1:A:406:VAL:HG12	1.89	0.53
1:A:574:PRO:O	1:A:596:THR:HB	2.09	0.51
1:A:248:ILE:O	1:A:250:PRO:HD3	2.11	0.51
1:A:210:LYS:HA	1:A:214:VAL:HG13	1.92	0.51
1:A:104:LEU:HD22	1:A:118:ARG:HB2	1.93	0.50
1:B:194:PRO:HG3	1:B:198:GLN:HB2	1.92	0.50
1:B:341:LEU:HD13	1:B:474:ARG:HB3	1.94	0.50
1:B:536:PHE:HA	1:B:539:LEU:HD22	1.93	0.49
1:A:171:PRO:HB2	1:A:173:GLU:HG2	1.93	0.49
1:A:251:ASN:HD22	1:A:260:THR:HA	1.78	0.49
1:B:414:LEU:HD22	1:B:418:TYR:HB3	1.94	0.48
1:A:14:LEU:HD22	1:B:21:LEU:HD23	1.96	0.48
1:B:456:VAL:HG23	1:B:483:SER:HB2	1.96	0.48
1:B:574:PRO:HG2	1:B:607:ASN:ND2	2.29	0.47
1:B:63:THR:HG22	1:B:72:THR:HA	1.96	0.47
1:A:158:VAL:HB	1:A:166:ALA:HB3	1.95	0.47
1:A:598:LEU:HD22	1:A:609:VAL:HG11	1.98	0.46
1:A:3:ILE:HG22	1:A:113:VAL:HG11	1.98	0.46
1:A:456:VAL:HG23	1:A:483:SER:HB2	1.97	0.46
1:B:333:HIS:CD2	1:B:335:ASN:H	2.29	0.46
1:A:236:LEU:HD22	1:A:257:ARG:HD2	1.98	0.45
1:A:565:ALA:HB1	1:A:585:LEU:HD21	1.99	0.45
1:B:580:GLN:HA	1:B:580:GLN:HE21	1.81	0.45
1:A:132:VAL:HG22	1:A:159:CYS:SG	2.57	0.44
1:B:82:LEU:HD11	1:B:175:MET:HG2	2.00	0.43
1:A:712:ARG:HG3	1:A:8:GLN:HB2	2.00	0.43
1:A:576:PRO:HA	1:A:596:THR:HG21	2.01	0.43
1:B:575:PRO:HB3	1:B:585:LEU:HD23	2.00	0.42
1:B:236:LEU:HD23	1:B:252:ILE:HG21	2.01	0.42
1:B:104:LEU:HB2	1:B:116:VAL:O	2.20	0.42
1:A:23:GLY:HA3	1:A:70:PRO:HG3	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:436:VAL:HG11	1:A:488:SER:HB3	2.02	0.42
1:B:171:PRO:HB2	1:B:173:GLU:HG2	2.03	0.41
1:B:339:VAL:O	1:B:474:ARG:HA	2.21	0.40
1:B:710:VAL:HA	1:B:11:ARG:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	641/666 (96%)	613 (96%)	26 (4%)	2 (0%)	46	67
1	B	638/666 (96%)	616 (97%)	19 (3%)	3 (0%)	34	55
All	All	1279/1332 (96%)	1229 (96%)	45 (4%)	5 (0%)	39	60

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	ASP
1	B	100	GLY
1	B	99	CYS
1	A	245	ALA
1	B	281	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/544 (91%)	464 (94%)	29 (6%)	24	42
1	B	498/544 (92%)	462 (93%)	36 (7%)	18	32
All	All	991/1088 (91%)	926 (93%)	65 (7%)	21	36

All (65) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	712	ARG
1	A	64	LEU
1	A	82	LEU
1	A	104	LEU
1	A	109	ARG
1	A	126	LEU
1	A	173	GLU
1	A	189	SER
1	A	211	SER
1	A	214	VAL
1	A	225	VAL
1	A	231	SER
1	A	232	VAL
1	A	249	ASP
1	A	329	VAL
1	A	341	LEU
1	A	344	THR
1	A	397	VAL
1	A	405	ASP
1	A	419	THR
1	A	447	GLU
1	A	449	THR
1	A	529	LEU
1	A	585	LEU
1	A	588	LEU
1	A	596	THR
1	A	600	TYR
1	A	611	LEU
1	A	627	LEU
1	B	21	LEU
1	B	27	ASN
1	B	64	LEU
1	B	72	THR
1	B	82	LEU
1	B	95	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	126	LEU
1	B	139	SER
1	B	155	ARG
1	B	172	VAL
1	B	173	GLU
1	B	176	GLU
1	B	210	LYS
1	B	341	LEU
1	B	344	THR
1	B	373	LYS
1	B	395	LEU
1	B	397	VAL
1	B	403	ILE
1	B	414	LEU
1	B	419	THR
1	B	433	THR
1	B	453	GLN
1	B	477	THR
1	B	529	LEU
1	B	535	VAL
1	B	539	LEU
1	B	580	GLN
1	B	585	LEU
1	B	591	THR
1	B	592	LEU
1	B	600	TYR
1	B	609	VAL
1	B	611	LEU
1	B	627	LEU
1	B	631	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	251	ASN
1	A	333	HIS
1	A	606	GLN
1	A	607	ASN
1	B	27	ASN
1	B	221	GLN
1	B	229	ASN
1	B	251	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	333	HIS
1	B	580	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	R2N	A	1721	-	10,15,15	0.71	0	13,22,22	0.80	0
2	R2N	B	1721	-	10,15,15	0.81	1 (10%)	13,22,22	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	R2N	A	1721	-	-	0/0/4/4	0/2/2/2
2	R2N	B	1721	-	-	0/0/4/4	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1721	R2N	C6-C7	2.10	1.39	1.36

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	645/666 (96%)	-0.09	10 (1%) 74 78	21, 44, 78, 108	0
1	B	642/666 (96%)	-0.16	6 (0%) 85 88	25, 44, 68, 90	0
All	All	1287/1332 (96%)	-0.13	16 (1%) 81 84	21, 44, 74, 108	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	388	ALA	5.0
1	A	99	CYS	4.6
1	A	100	GLY	3.7
1	B	3	ILE	3.3
1	B	104	LEU	3.2
1	B	100	GLY	3.1
1	B	196	SER	3.1
1	A	468	GLY	3.0
1	A	386	ILE	2.9
1	A	473	TYR	2.6
1	A	361	GLY	2.3
1	A	127	LEU	2.2
1	B	593	HIS	2.2
1	A	472	ILE	2.1
1	B	4	THR	2.1
1	A	339	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	R2N	A	1721	14/14	0.89	0.24	2.91	77,78,82,82	21
2	R2N	B	1721	14/14	0.92	0.21	2.58	52,63,67,73	21

6.5 Other polymers [i](#)

There are no such residues in this entry.