



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:59 PM GMT

PDB ID : 4GUR
Title : Crystal structure of LSD2-NPAC with H3 in space group P21
Authors : Chen, F.; Dong, Z.; Fang, J.; Yang, Y.; Li, Z.; Xu, Y.; Yang, H.; Wang, P.;
Fang, R.; Shi, Y.; Xu, Y.
Deposited on : 2012-08-29
Resolution : 2.51 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	776	<div> <div>3%</div> <div>75%</div> <div>17%</div> <div>5%</div> </div>
2	B	124	<div> <div>2%</div> <div>7%</div> <div>90%</div> </div>
3	C	21	<div> <div>29%</div> <div>67%</div> <div>29%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	902	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6234 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific histone demethylase 1B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	734	Total	C	N	O	S	0	0	0
			5818	3716	990	1071	41			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	PRO	-	EXPRESSION TAG	UNP Q8NB78
A	48	LEU	-	EXPRESSION TAG	UNP Q8NB78
A	49	GLY	-	EXPRESSION TAG	UNP Q8NB78
A	50	SER	-	EXPRESSION TAG	UNP Q8NB78

- Molecule 2 is a protein called Putative oxidoreductase GLYR1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	0	0	0
			105	69	19	17			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	145	PRO	-	EXPRESSION TAG	UNP Q49A26
B	146	LEU	-	EXPRESSION TAG	UNP Q49A26
B	147	GLY	-	EXPRESSION TAG	UNP Q49A26
B	148	SER	-	EXPRESSION TAG	UNP Q49A26
B	149	PRO	-	EXPRESSION TAG	UNP Q49A26
B	150	GLU	-	EXPRESSION TAG	UNP Q49A26
B	151	PHE	-	EXPRESSION TAG	UNP Q49A26


- Molecule 3 is a protein called Histone H3.3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	20	Total	C	N	O	S	0	0	0
			151	90	34	26	1			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	4	MET	LYS	ENGINEERED MUTATION	UNP P84243

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	3	Total	Zn	0	0
			3	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	90	Total	O	0	0
			90	90		
7	B	3	Total	O	0	0
			3	3		
7	C	5	Total	O	0	0
			5	5		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:
-
- 3% 75% 17% 5%
- VAL PRD PRD GLY SER R51 R52 C53 T59 A60 T61 S70 E71 R72 N76 Y83 H84 L85 N93 E94 Y99 S102 R103 R104 D105 G106 Y107 D108 K109 Y110 T111 K114 W117 P125 K128 Q134 Q135 L136 P137 Y138 W139 T143 K149 R151 Q152
- L375 L376 L377 K378 K379 Y380 H381 H382 K383 K384 V385 I388 L401 K407 V410 L411 E412 W422 V430 P446 E452 Q453 L454 G455 I456 S457 L467 E470 G471 G472 R473 V493 V494 S495 R498 V507 K512 K515 I516 Y517 I521 K522 E523 S524
- Q527 L536 Q537 L543 E544 Y545 A546 C547 L551 W559 L574 P577 G578 V581 K585 V598 Q609 S619 V625 T626 A630 V634 L664 M671 T680 P685 P686 R691 G692 L693 Q703 K704 K705 V712 A719 S720 T721 R722
- VAL PRD PRD G264 W265 R266 R267 F272 Y273 E277 C283 P286 L284 P288 E299 R302 L308 T327 P328 Q329 K330 C331 P332 P333 R338 G339 R344 C345 V346 V349 L353 M356 T357 R358 K359 V362 N363 V366 V369 G370 D371 D372 D373 V374

- Chain B:
-
- | Amino Acid | Frequency (%) |
|------------|---------------|
| THR | 2% |
| ALA | 2% |
| SER | 2% |
| GLU | 2% |
| PRO | 2% |
| VAL | 2% |
| LYS | 2% |
| ASP | 2% |
| ALA | 2% |
| D14 | 2% |
| P215 | 2% |
| E218 | 2% |
| S223 | 2% |
| Q224 | 2% |
| T225 | 2% |
| GLU | 2% |
| LYS | 2% |
| PRO | 2% |
| ALA | 2% |
| VAL | 2% |
| CYS | 2% |
| TYR | 2% |
| GLN | 2% |
| ALA | 2% |
| ILE | 2% |
| THR | 2% |
| LYS | 2% |
| LYS | 2% |
| LEU | 2% |
| LYS | 2% |
| ILE | 2% |
| CYS | 2% |
| GLU | 2% |
| GLU | 2% |
| GLU | 2% |
| THR | 2% |
| GLY | 2% |
| LYS | 2% |
| SER | 2% |
| THR | 2% |
| SER | 2% |
| LYS | 2% |
| GLN | 2% |
| VAL | 2% |
| ASP | 2% |
| SER | 2% |
| THR | 2% |
| ASN | 2% |
| VAL | 2% |
| GLY | 2% |
| MET | 2% |
| LYS | 2% |
| VAL | 2% |
| LYS | 2% |
| THR | 2% |
| THR | 2% |
| THR | 2% |
| ASP | 2% |
| LYS | 2% |

- Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.24Å 89.03Å 88.78Å 90.00° 103.30° 90.00°	Depositor
Resolution (Å)	38.87 – 2.51 38.87 – 2.51	Depositor EDS
% Data completeness (in resolution range)	88.6 (38.87-2.51) 94.3 (38.87-2.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.60 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.1_353)	Depositor
R, R_{free}	0.188 , 0.204 0.187 , 0.200	Depositor DCC
R_{free} test set	1563 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	42.6	Xtriage
Anisotropy	0.435	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 30672 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6234	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	7/5962 (0.1%)	0.65	0/8076
2	B	0.74	0/110	0.59	0/149
3	C	0.38	0/151	0.58	0/198
All	All	0.55	7/6223 (0.1%)	0.64	0/8423

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	559	TRP	CD2-CE2	5.53	1.48	1.41
1	A	139	TRP	CD2-CE2	5.44	1.47	1.41
1	A	671	TRP	CD2-CE2	5.38	1.47	1.41
1	A	117	TRP	CD2-CE2	5.25	1.47	1.41
1	A	200	TRP	CD2-CE2	5.19	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5818	0	5730	83	0
2	B	105	0	91	1	0
3	C	151	0	170	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	53	0	30	3	0
5	A	6	0	8	2	0
6	A	3	0	0	0	0
7	A	90	0	0	2	0
7	B	3	0	0	0	0
7	C	5	0	0	0	0
All	All	6234	0	6029	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 84 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:ARG:HG2	1:A:72:ARG:HH11	0.94	1.05
1:A:72:ARG:HG2	1:A:72:ARG:NH1	1.70	0.94
1:A:72:ARG:CG	1:A:72:ARG:HH11	1.84	0.89
1:A:473:ARG:HH11	1:A:473:ARG:HG3	1.46	0.80
1:A:266:ASN:HD22	1:A:452:GLU:HG2	1.50	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	728/776 (94%)	684 (94%)	40 (6%)	4 (0%)	34	55
2	B	10/124 (8%)	9 (90%)	1 (10%)	0	100	100
3	C	18/21 (86%)	16 (89%)	2 (11%)	0	100	100
All	All	756/921 (82%)	709 (94%)	43 (6%)	4 (0%)	34	55

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	155	LYS
1	A	372	ASP
1	A	523	GLU
1	A	803	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	630/662 (95%)	577 (92%)	53 (8%)	14	25
2	B	12/106 (11%)	11 (92%)	1 (8%)	14	26
3	C	15/15 (100%)	12 (80%)	3 (20%)	1	3
All	All	657/783 (84%)	600 (91%)	57 (9%)	13	24

5 of 57 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	366	VAL
1	A	430	VAL
1	A	775	SER
1	A	369	VAL
1	A	385	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	440	ASN
1	A	527	GLN
1	A	732	GLN
1	A	400	GLN
1	A	609	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	FAD	A	901	-	48,58,58	2.34	23 (47%)	54,89,89	2.15	13 (24%)
5	GOL	A	902	-	5,5,5	0.36	0	5,5,5	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FAD	A	901	-	-	0/30/50/50	0/6/6/6
5	GOL	A	902	-	-	0/4/4/4	0/0/0/0

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	901	FAD	C10-N10	-5.07	1.33	1.39
4	A	901	FAD	PA-O5B	-4.36	1.39	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	901	FAD	O4B-C1B	-4.22	1.35	1.41
4	A	901	FAD	P-O2P	-4.10	1.37	1.54
4	A	901	FAD	PA-O1A	-3.90	1.36	1.51

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	901	FAD	N3A-C2A-N1A	-8.07	122.72	128.89
4	A	901	FAD	C4A-C5A-N7A	-2.98	106.74	109.48
4	A	901	FAD	C4X-C4-N3	-2.57	120.07	123.59
4	A	901	FAD	O4'-C4'-C5'	-2.48	104.79	110.19
4	A	901	FAD	C4B-O4B-C1B	-2.31	107.18	109.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	901	FAD	3	0
5	A	902	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	734/776 (94%)	0.05	22 (2%) 54 59	26, 47, 83, 102	0
2	B	12/124 (9%)	0.76	3 (25%) 1 1	46, 66, 82, 90	0
3	C	20/21 (95%)	0.85	6 (30%) 1 0	47, 55, 90, 95	0
All	All	766/921 (83%)	0.09	31 (4%) 42 47	26, 47, 84, 102	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	822	PHE	6.5
1	A	527	GLN	4.8
1	A	200	TRP	3.7
3	C	18	LYS	3.6
2	B	225	THR	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	GOL	A	902	6/6	0.85	0.28	4.11	48,51,54,60	0
4	FAD	A	901	53/53	0.96	0.17	-0.20	26,32,38,43	0
6	ZN	A	905	1/1	0.99	0.07	-1.45	55,55,55,55	0
6	ZN	A	903	1/1	0.97	0.07	-1.59	63,63,63,63	0
6	ZN	A	904	1/1	0.99	0.10	-2.17	54,54,54,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.