



Full wwPDB X-ray Structure Validation Report ⓘ

May 30, 2016 – 05:42 AM EDT

PDB ID : 5HGA
Title : HLA*A2402 complex with HIV nef138 Y2F-8mer mutant epitope
Authors : Shi, Y.; Qi, J.; Gao, G.F.
Deposited on : 2016-01-08
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>%</div> <div>81% 17% .</div> </div>
1	D	275	<div> <div>5%</div> <div>83% 16% .</div> </div>
2	B	100	<div> <div>5%</div> <div>82% 16% ..</div> </div>
2	E	100	<div> <div>8%</div> <div>83% 12% . .</div> </div>
3	C	8	<div> <div>13%</div> <div>50% 38% 13%</div> </div>
3	F	8	<div> <div>38%</div> <div>63% 25% 13%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6416 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-24 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	274	Total	C	N	O	S	0	0	0
			2221	1382	403	426	10			
1	D	274	Total	C	N	O	S	0	0	0
			2221	1382	403	426	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P05534
D	0	MET	-	initiating methionine	UNP P05534

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			836	533	141	158	4			
2	E	100	Total	C	N	O	S	0	0	0
			836	533	141	158	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
E	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called 8-mer from Protein Nef.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	8	Total	C	N	O	0	0	0
			74	52	12	10			
3	F	8	Total	C	N	O	0	0	0
			74	52	12	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	2	PHE	TYR	engineered mutation	UNP P18801
F	2	PHE	TYR	engineered mutation	UNP P18801


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	62	Total O 62 62	0	0
4	B	14	Total O 14 14	0	0
4	C	8	Total O 8 8	0	0
4	D	53	Total O 53 53	0	0
4	E	14	Total O 14 14	0	0
4	F	3	Total O 3 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, A-24 alpha chain




- Molecule 1: HLA class I histocompatibility antigen, A-24 alpha chain



- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin




- Molecule 3: 8-mer from Protein Nef





- Molecule 3: 8-mer from Protein Nef



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.84Å 86.91Å 151.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.86 – 2.20 37.86 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.86-2.20) 99.9 (37.86-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.205 , 0.250 0.205 , 0.251	Depositor DCC
R_{free} test set	2223 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	43.5	Xtriage
Anisotropy	0.555	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6416	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.50 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.8216e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.32	0/2281	0.53	0/3092
1	D	0.32	0/2281	0.53	0/3092
2	B	0.31	0/859	0.52	0/1162
2	E	0.32	0/859	0.53	0/1162
3	C	0.33	0/78	0.58	0/104
3	F	0.33	0/78	0.55	0/104
All	All	0.32	0/6436	0.53	0/8716

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2221	0	2082	27	0
1	D	2221	0	2082	23	1
2	B	836	0	803	9	0
2	E	836	0	803	11	0
3	C	74	0	71	4	0
3	F	74	0	71	2	0
4	A	62	0	0	3	0
4	B	14	0	0	2	1
4	C	8	0	0	2	0
4	D	53	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	14	0	0	1	0
4	F	3	0	0	0	0
All	All	6416	0	5912	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:GLY:O	4:A:301:HOH:O	1.99	0.79
2:B:42:ASN:O	4:B:101:HOH:O	2.07	0.72
2:B:19:LYS:O	4:B:102:HOH:O	2.08	0.71
1:D:211:ALA:O	4:D:301:HOH:O	2.12	0.68
1:A:177:GLU:HG2	1:A:178:THR:HG23	1.78	0.66
1:D:154:GLU:OE2	4:D:302:HOH:O	2.15	0.64
1:D:144:LYS:NZ	1:D:148:GLU:OE2	2.31	0.64
3:C:1:ARG:NH1	4:C:101:HOH:O	2.25	0.64
2:B:96:ASP:O	2:B:98:ASP:N	2.31	0.63
1:A:222:GLU:O	1:A:224:GLN:NE2	2.31	0.61
1:A:106:ASP:OD1	4:A:302:HOH:O	2.17	0.58
1:D:111:ARG:NH2	4:D:303:HOH:O	2.27	0.57
2:E:96:ASP:O	2:E:98:ASP:N	2.38	0.56
2:E:99:MET:HB2	4:E:104:HOH:O	2.04	0.56
1:D:44:ARG:NH2	1:D:61:ASP:OD1	2.38	0.55
1:A:255:GLN:O	1:A:273:ARG:NH1	2.34	0.55
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.42	0.55
1:D:268:LYS:HG3	1:D:269:PRO:HD2	1.90	0.54
1:A:230:LEU:HD13	1:A:245:ALA:HB2	1.89	0.53
1:D:127:LYS:NZ	1:D:134:THR:OG1	2.30	0.53
1:A:268:LYS:HG3	1:A:269:PRO:HD2	1.91	0.52
1:A:106:ASP:OD1	1:A:108:ARG:HD3	2.12	0.51
2:B:16:GLU:OE1	2:E:59:ASP:HB3	2.12	0.50
1:D:119:ASP:HB3	2:E:0:MET:HA	1.95	0.49
1:A:106:ASP:OD2	1:A:108:ARG:NH1	2.44	0.48
3:C:1:ARG:NH1	4:C:102:HOH:O	2.32	0.48
1:D:202:ARG:HD3	1:D:244:TRP:CE3	2.48	0.48
2:E:25:CYS:HB2	2:E:39:LEU:HD21	1.96	0.48
1:D:202:ARG:HG2	1:D:204:TRP:NE1	2.29	0.47
1:A:224:GLN:O	1:A:228:THR:HG23	2.15	0.46
1:A:19:GLU:OE1	1:A:75:ARG:HD3	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.50	0.46
1:A:5:MET:HB2	1:A:168:LEU:HD13	1.97	0.46
1:A:202:ARG:HD3	1:A:244:TRP:CE3	2.50	0.46
1:A:70:HIS:ND1	3:C:5:THR:HG21	2.31	0.46
2:B:16:GLU:HG3	2:B:19:LYS:HB2	1.98	0.46
1:D:224:GLN:O	1:D:228:THR:HG23	2.16	0.46
1:D:223:ASP:HB3	1:D:225:THR:HG23	1.98	0.46
2:B:74:GLU:OE2	2:B:75:LYS:N	2.49	0.45
1:D:234:ARG:HH12	2:E:99:MET:HE2	1.81	0.45
2:E:58:LYS:HE3	2:E:58:LYS:H	1.82	0.45
2:E:74:GLU:OE2	2:E:75:LYS:N	2.49	0.44
1:A:55:GLU:OE1	1:A:170:ARG:NH2	2.50	0.44
1:A:273:ARG:HG3	1:A:274:TRP:N	2.33	0.44
1:D:49:ALA:O	1:D:52:ILE:HG22	2.18	0.44
3:F:2:PHE:CG	3:F:3:PRO:HD2	2.54	0.43
1:D:44:ARG:NH2	1:D:61:ASP:HA	2.34	0.43
1:A:154:GLU:OE2	4:A:303:HOH:O	2.21	0.43
1:A:187:THR:HG21	1:A:261:VAL:HG21	2.01	0.43
1:A:154:GLU:HG2	1:A:154:GLU:H	1.59	0.42
1:D:35:ARG:HD3	2:E:53:ASP:OD2	2.19	0.42
1:D:154:GLU:HB3	4:D:308:HOH:O	2.18	0.42
1:D:250:PRO:HB2	1:D:253:GLU:HG3	2.01	0.42
1:A:202:ARG:HG2	1:A:204:TRP:NE1	2.35	0.42
1:D:182:THR:HG21	1:D:265:GLY:HA2	2.01	0.42
2:E:29:GLY:HA2	2:E:61:SER:OG	2.20	0.42
1:D:70:HIS:ND1	3:F:5:THR:HG21	2.35	0.42
1:A:250:PRO:HB2	1:A:253:GLU:HG3	2.01	0.42
3:C:2:PHE:CG	3:C:3:PRO:HD2	2.54	0.42
2:B:37:VAL:HG22	2:B:82:VAL:HG22	2.01	0.41
1:A:49:ALA:O	1:A:52:ILE:HG22	2.20	0.41
1:A:44:ARG:NH2	1:A:61:ASP:OD1	2.47	0.41
1:A:249:VAL:HG21	1:A:254:GLU:HG3	2.02	0.41
1:D:127:LYS:HD2	1:D:132:SER:OG	2.21	0.41
1:A:110:LEU:HD12	1:A:110:LEU:HA	1.88	0.41
1:D:266:LEU:HA	1:D:267:PRO:HD2	1.91	0.41
2:B:51:HIS:HB3	2:B:66:TYR:CD2	2.56	0.41
1:A:201:LEU:HA	1:A:201:LEU:HD23	1.80	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:166:ASP:OD1	4:B:101:HOH:O[4_445]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/275 (99%)	264 (97%)	6 (2%)	2 (1%)	26	25
1	D	272/275 (99%)	264 (97%)	6 (2%)	2 (1%)	26	25
2	B	98/100 (98%)	95 (97%)	1 (1%)	2 (2%)	9	5
2	E	98/100 (98%)	95 (97%)	1 (1%)	2 (2%)	9	5
3	C	6/8 (75%)	6 (100%)	0	0	100	100
3	F	6/8 (75%)	6 (100%)	0	0	100	100
All	All	752/766 (98%)	730 (97%)	14 (2%)	8 (1%)	17	14

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	98	ASP
2	E	98	ASP
1	A	225	THR
2	B	97	ARG
1	D	225	THR
1	A	221	GLY
1	D	221	GLY
2	E	97	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/231 (100%)	214 (93%)	16 (7%)	19	19
1	D	230/231 (100%)	216 (94%)	14 (6%)	23	26
2	B	95/95 (100%)	88 (93%)	7 (7%)	17	17
2	E	95/95 (100%)	88 (93%)	7 (7%)	17	17
3	C	7/7 (100%)	6 (86%)	1 (14%)	4	3
3	F	7/7 (100%)	6 (86%)	1 (14%)	4	3
All	All	664/666 (100%)	618 (93%)	46 (7%)	19	20

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	THR
1	A	74	ASP
1	A	79	ARG
1	A	82	LEU
1	A	110	LEU
1	A	141	GLN
1	A	154	GLU
1	A	170	ARG
1	A	180	GLN
1	A	182	THR
1	A	186	LYS
1	A	197	HIS
1	A	222	GLU
1	A	226	GLN
1	A	256	ARG
1	A	273	ARG
2	B	0	MET
2	B	4	THR
2	B	58	LYS
2	B	70	PHE
2	B	74	GLU
2	B	98	ASP
2	B	99	MET
3	C	5	THR
1	D	73	THR
1	D	74	ASP
1	D	79	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	82	LEU
1	D	110	LEU
1	D	141	GLN
1	D	154	GLU
1	D	170	ARG
1	D	182	THR
1	D	183	ASP
1	D	186	LYS
1	D	222	GLU
1	D	226	GLN
1	D	256	ARG
2	E	0	MET
2	E	4	THR
2	E	58	LYS
2	E	70	PHE
2	E	74	GLU
2	E	98	ASP
2	E	99	MET
3	F	5	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/275 (99%)	0.24	4 (1%) 76 75	15, 31, 68, 106	0
1	D	274/275 (99%)	0.38	13 (4%) 35 34	17, 33, 64, 85	0
2	B	100/100 (100%)	0.42	5 (5%) 32 32	18, 37, 67, 92	0
2	E	100/100 (100%)	0.53	8 (8%) 15 14	17, 35, 68, 83	0
3	C	8/8 (100%)	1.08	1 (12%) 5 5	20, 22, 24, 38	0
3	F	8/8 (100%)	1.33	3 (37%) 0 0	21, 25, 29, 42	0
All	All	764/766 (99%)	0.37	34 (4%) 37 36	15, 32, 68, 106	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	273	ARG	5.1
2	E	99	MET	5.0
2	E	0	MET	4.0
1	A	226	GLN	3.7
1	D	16	GLY	3.5
1	D	110	LEU	3.3
1	D	225	THR	3.1
1	D	222	GLU	3.1
2	B	99	MET	3.1
2	B	98	ASP	2.9
2	E	98	ASP	2.8
2	B	19	LYS	2.8
3	F	6	PHE	2.8
2	E	44	GLU	2.6
1	A	197	HIS	2.6
1	A	227	ASP	2.6
1	D	197	HIS	2.6
2	B	0	MET	2.6
2	E	75	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	74	GLU	2.4
1	D	17	ARG	2.4
3	C	6	PHE	2.4
3	F	4	LEU	2.3
1	D	256	ARG	2.3
1	D	177	GLU	2.2
2	E	74	GLU	2.2
1	D	107	GLY	2.2
2	E	1	ILE	2.2
1	D	226	GLN	2.2
3	F	5	THR	2.2
2	E	92	ILE	2.1
1	D	105	SER	2.1
1	D	25	VAL	2.0
1	D	166	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.