



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2016 – 11:24 PM EDT

PDB ID : 5HHM
Title : Crystal Structure of the JM22 TCR in complex with HLA-A*0201 in complex with M1-F5L
Authors : Gras, S.; Josephs, T.M.; Rossjohn, J.
Deposited on : 2016-01-11
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027257
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027257


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	<div> <div>0%</div> <div> <div></div> <div>67%</div> <div>9%</div> <div>23%</div> </div> </div>
1	F	276	<div> <div>2%</div> <div> <div></div> <div>67%</div> <div>9%</div> <div>23%</div> </div> </div>
2	B	100	<div> <div>10%</div> <div> <div></div> <div>79%</div> <div>18%</div> <div>3%</div> </div> </div>
2	G	100	<div> <div>3%</div> <div> <div></div> <div>81%</div> <div>15%</div> <div>3%</div> </div> </div>
3	C	9	<div> <div></div> <div>100%</div> </div>
3	H	9	<div> <div></div> <div>78%</div> <div>22%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	200	<div><div></div><div>4%</div><div>84%</div><div>12%</div><div></div><div></div></div>
4	I	200	<div><div></div><div>10%</div><div>87%</div><div>11%</div><div></div><div></div></div>
5	E	241	<div><div></div><div>%</div><div>84%</div><div>15%</div><div></div><div></div></div>
5	J	241	<div><div></div><div>%</div><div>86%</div><div>12%</div><div></div><div></div></div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12678 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-2 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	0	0	0
			1727	1083	312	326	6			
1	F	212	Total	C	N	O	S	0	0	0
			1727	1083	312	326	6			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	98	Total	C	N	O	S	0	0	0
			820	524	139	154	3			
2	G	98	Total	C	N	O	S	0	0	0
			820	524	139	154	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
G	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called M1-F5L, GILGLVFTL.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			66	46	9	11			
3	H	9	Total	C	N	O	0	0	0
			66	46	9	11			

- Molecule 4 is a protein called JM22 TCR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	194	Total	C	N	O	S	0	0	0
			1495	940	249	300	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	I	196	Total	C	N	O	S	0	0	0
			1507	946	251	304	6			

- Molecule 5 is a protein called JM22 TCR beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	240	Total	C	N	O	S	0	1	0
			1931	1218	334	374	5			
5	J	240	Total	C	N	O	S	0	1	0
			1931	1218	334	374	5			


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	109	Total	O	0	0
			109	109		
6	B	37	Total	O	0	0
			37	37		
6	C	3	Total	O	0	0
			3	3		
6	D	57	Total	O	0	0
			57	57		
6	E	107	Total	O	0	0
			107	107		
6	F	82	Total	O	0	0
			82	82		
6	G	22	Total	O	0	0
			22	22		
6	H	2	Total	O	0	0
			2	2		
6	I	68	Total	O	0	0
			68	68		
6	J	101	Total	O	0	0
			101	101		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain




- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain



- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin




- Molecule 3: M1-F5L, GILGLVFTL

Chain C:  100%


There are no outlier residues recorded for this chain.


- Molecule 3: M1-F5L, GILGLVFTL

Chain H:  78% 22%





- Molecule 4: JM22 TCR alpha chain

Chain D:  4% 84% 12% ..





- Molecule 4: JM22 TCR alpha chain

Chain I:  10% 87% 11% ..





- Molecule 5: JM22 TCR beta chain


Chain E:  % 84% 15% .



- Molecule 5: JM22 TCR beta chain

Chain J:  % 86% 12% .





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	242.88Å 47.30Å 185.94Å 90.00° 115.49° 90.00°	Depositor
Resolution (Å)	43.58 – 2.50 43.58 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (43.58-2.50) 99.8 (43.58-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.51Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.10.0	Depositor
R, R_{free}	0.216 , 0.289 0.227 , 0.296	Depositor DCC
R_{free} test set	3250 reflections (5.10%)	DCC
Wilson B-factor (Å ²)	34.9	Xtriage
Anisotropy	0.415	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 61.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12678	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.01 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.9098e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/1774	0.73	1/2399 (0.0%)
1	F	0.49	0/1774	0.73	0/2399
2	B	0.53	0/843	0.74	0/1141
2	G	0.47	0/843	0.74	0/1141
3	C	0.74	0/66	0.90	0/87
3	H	0.64	0/66	0.71	0/87
4	D	0.51	0/1524	0.74	0/2064
4	I	0.50	0/1536	0.73	0/2080
5	E	0.54	0/1987	0.76	0/2702
5	J	0.50	0/1987	0.73	0/2702
All	All	0.51	0/12400	0.74	1/16802 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	183	ASP	C-N-CA	7.15	139.56	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1727	0	1618	15	0
1	F	1727	0	1618	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	820	0	792	8	0
2	G	820	0	792	9	0
3	C	66	0	77	0	0
3	H	66	0	77	0	0
4	D	1495	0	1449	11	0
4	I	1507	0	1459	10	0
5	E	1931	0	1835	19	0
5	J	1931	0	1835	16	0
6	A	109	0	0	0	0
6	B	37	0	0	0	0
6	C	3	0	0	0	0
6	D	57	0	0	0	0
6	E	107	0	0	0	0
6	F	82	0	0	0	0
6	G	22	0	0	0	0
6	H	2	0	0	0	0
6	I	68	0	0	1	0
6	J	101	0	0	0	0
All	All	12678	0	11552	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:101:CYS:HG	1:F:164:CYS:HG	1.24	0.84
2:G:17:ASN:OD1	2:G:97:ARG:NH2	2.12	0.83
1:A:234:ARG:HE	1:A:242:GLN:HE21	1.28	0.80
2:B:25:CYS:HG	2:B:80:CYS:HG	0.81	0.78
5:E:148:THR:HG22	5:E:189:SER:OG	1.96	0.65

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/276 (75%)	198 (96%)	7 (3%)	1 (0%)	34	55
1	F	206/276 (75%)	194 (94%)	12 (6%)	0	100	100
2	B	96/100 (96%)	92 (96%)	2 (2%)	2 (2%)	9	14
2	G	96/100 (96%)	93 (97%)	3 (3%)	0	100	100
3	C	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	H	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
4	D	190/200 (95%)	181 (95%)	9 (5%)	0	100	100
4	I	192/200 (96%)	185 (96%)	7 (4%)	0	100	100
5	E	239/241 (99%)	230 (96%)	9 (4%)	0	100	100
5	J	239/241 (99%)	229 (96%)	10 (4%)	0	100	100
All	All	1478/1652 (90%)	1414 (96%)	61 (4%)	3 (0%)	52	75

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	184	ALA
2	B	73	THR
2	B	72	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/232 (75%)	168 (97%)	6 (3%)	44	72
1	F	174/232 (75%)	168 (97%)	6 (3%)	44	72
2	B	93/95 (98%)	87 (94%)	6 (6%)	21	39
2	G	93/95 (98%)	86 (92%)	7 (8%)	17	31
3	C	7/7 (100%)	7 (100%)	0	100	100
3	H	7/7 (100%)	5 (71%)	2 (29%)	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	D	170/176 (97%)	161 (95%)	9 (5%)	28	50
4	I	172/176 (98%)	164 (95%)	8 (5%)	32	56
5	E	212/212 (100%)	201 (95%)	11 (5%)	29	51
5	J	212/212 (100%)	202 (95%)	10 (5%)	32	56
All	All	1314/1444 (91%)	1249 (95%)	65 (5%)	30	55

5 of 65 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	E	193	ARG
1	F	176	LYS
5	J	107	PRO
5	E	208	PHE
1	F	34	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
5	E	233	GLN
1	F	180	GLN
5	J	43	GLN
1	F	72	GLN
2	B	51	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	212/276 (76%)	-0.11	2 (0%) 85 88	17, 37, 66, 87	1 (0%)
1	F	212/276 (76%)	-0.06	6 (2%) 56 61	13, 38, 66, 96	1 (0%)
2	B	98/100 (98%)	0.50	10 (10%) 9 9	21, 47, 81, 96	0
2	G	98/100 (98%)	0.25	3 (3%) 52 57	23, 45, 80, 88	0
3	C	9/9 (100%)	-0.02	0 100 100	18, 24, 29, 31	0
3	H	9/9 (100%)	-0.40	0 100 100	15, 23, 27, 27	0
4	D	194/200 (97%)	0.08	8 (4%) 41 46	12, 42, 95, 120	0
4	I	196/200 (98%)	0.28	19 (9%) 10 10	14, 43, 92, 116	0
5	E	240/241 (99%)	-0.16	2 (0%) 87 89	11, 36, 68, 79	0
5	J	240/241 (99%)	-0.18	3 (1%) 79 82	9, 36, 76, 98	0
All	All	1508/1652 (91%)	0.01	53 (3%) 48 53	9, 38, 83, 120	2 (0%)

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	70	PHE	5.4
1	A	244	TRP	5.1
4	I	126	SER	5.0
4	I	186	ASN	4.4
2	B	20	SER	4.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.