



# Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2016 – 07:52 PM EDT

PDB ID : 1HJA  
Title : LYS 18 VARIANT OF TURKEY OVOMUCOID INHIBITOR THIRD DO-  
MAIN COMPLEXED WITH ALPHA-CHYMOTRYPSIN  
Authors : Ding, J.-H.; James, M.N.G.  
Deposited on : 1997-07-09  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027674  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027674


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	13	<div> <div>54%</div> <div>23%</div> <div>23%</div> </div>
2	B	131	<div> <div>78%</div> <div>21%</div> <div>.</div> </div>
3	C	97	<div> <div>76%</div> <div>23%</div> <div>.</div> </div>
4	I	51	<div> <div>2%</div> <div>71%</div> <div>29%</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-CHYMOTRYPSIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	10	Total	C	N	O	S	0	0	0
			68	45	11	11	1			

- Molecule 2 is a protein called ALPHA-CHYMOTRYPSIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	131	Total	C	N	O	S	0	0	0
			980	618	162	196	4			

- Molecule 3 is a protein called ALPHA-CHYMOTRYPSIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	97	Total	C	N	O	S	0	0	0
			702	436	123	136	7			

- Molecule 4 is a protein called OVOMUCOID INHIBITOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	I	51	Total	C	N	O	S	0	0	0
			388	238	66	78	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	18	LYS	LEU	CONFLICT	UNP P68390

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	10	Total	O	0	0
			10	10		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	44	Total 44	O 44	0	0
5	C	24	Total 24	O 24	0	0
5	I	21	Total 21	O 21	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: ALPHA-CHYMOTRYPSIN

Chain A: 




#### • Molecule 2: ALPHA-CHYMOTRYPSIN

Chain B: 



#### • Molecule 3: ALPHA-CHYMOTRYPSIN

Chain C: 



#### • Molecule 4: OVOMUCOID INHIBITOR

Chain I: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.47Å 103.47Å 47.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 2.30 24.85 – 2.16	Depositor EDS
% Data completeness (in resolution range)	86.9 (10.00-2.30) 85.2 (24.85-2.16)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.56 (at 2.17Å)	Xtriage
Refinement program	X-PLOR 3.843	Depositor
R, $R_{free}$	0.193 , 0.246 0.194 , 0.235	Depositor DCC
$R_{free}$ test set	1214 reflections (10.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.6	Xtriage
Anisotropy	0.277	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 52.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.064 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2237	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/69	0.83	0/95
2	B	0.46	0/1000	0.71	0/1361
3	C	0.40	0/715	0.70	0/973
4	I	0.44	0/396	0.71	0/533
All	All	0.44	0/2180	0.71	0/2962

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	68	0	76	2	0
2	B	980	0	951	19	0
3	C	702	0	698	14	0
4	I	388	0	360	9	0
5	A	10	0	0	0	0
5	B	44	0	0	0	0
5	C	24	0	0	1	0
5	I	21	0	0	0	0
All	All	2237	0	2085	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:I:12:PRO:HG3	4:I:42:VAL:HG21	1.49	0.95
2:B:79:LYS:HA	2:B:79:LYS:HE3	1.63	0.81
2:B:66:VAL:HG23	2:B:85:ILE:HD11	1.63	0.79
3:C:217:SER:HB2	3:C:224:THR:HG21	1.70	0.74
2:B:144:THR:HG23	3:C:152:PRO:HD3	1.72	0.72
2:B:48:ASN:ND2	2:B:50:ASN:H	1.94	0.66
2:B:48:ASN:HD22	2:B:50:ASN:H	1.45	0.65
3:C:236:ASN:O	3:C:240:GLN:HG3	1.99	0.62
2:B:79:LYS:CE	2:B:79:LYS:HA	2.30	0.59
3:C:201:CYS:SG	3:C:210:VAL:HG21	2.42	0.59
4:I:39:ASN:O	4:I:42:VAL:HG22	2.03	0.58
4:I:23:LEU:HD23	4:I:50:LEU:HD13	1.85	0.58
3:C:202:LYS:HE3	3:C:207:TRP:CE2	2.40	0.56
3:C:165:ASN:O	3:C:169:LYS:HG3	2.06	0.56
2:B:86:ALA:HB2	2:B:109:SER:HB3	1.88	0.55
2:B:46:LEU:HD23	2:B:52:VAL:HG22	1.92	0.52
1:A:9:VAL:HG23	2:B:23:VAL:HG21	1.93	0.51
2:B:29:TRP:CG	2:B:121:VAL:HB	2.45	0.51
3:C:197:GLY:HA3	5:C:3:HOH:O	2.10	0.51
2:B:51:TRP:CZ2	2:B:107:LYS:HD3	2.46	0.50
3:C:172:TRP:HB3	3:C:175:LYS:HB2	1.95	0.48
2:B:146:TYR:CD2	3:C:219:THR:HA	2.50	0.47
2:B:93:LYS:HE2	2:B:93:LYS:HA	1.97	0.47
4:I:44:SER:HB2	4:I:47:THR:OG1	2.15	0.46
3:C:202:LYS:HE3	3:C:207:TRP:NE1	2.29	0.46
3:C:187:GLY:C	3:C:222:THR:HB	2.36	0.46
2:B:95:ASN:HD21	2:B:97:LEU:HB3	1.81	0.45
4:I:11:TYR:HB3	4:I:12:PRO:HA	1.99	0.45
4:I:37:PHE:O	4:I:41:VAL:HG23	2.17	0.45
4:I:51:SER:O	4:I:52:HIS:HB3	2.15	0.45
2:B:36:LYS:HG2	2:B:63:SER:O	2.17	0.44
2:B:146:TYR:HD2	3:C:219:THR:HA	1.83	0.43
3:C:184:GLY:O	3:C:185:ALA:HB3	2.19	0.43
2:B:124:PRO:HD3	3:C:209:LEU:O	2.19	0.42
4:I:22:PRO:HA	4:I:31:TYR:O	2.21	0.41
4:I:12:PRO:HG3	4:I:42:VAL:CG2	2.35	0.41
1:A:7:GLN:NE2	1:A:8:PRO:HD2	2.35	0.41
2:B:145:ARG:HG2	2:B:146:TYR:N	2.36	0.40
2:B:39:PHE:N	2:B:39:PHE:CD1	2.89	0.40



There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	8/13 (62%)	8 (100%)	0	0	100	100
2	B	129/131 (98%)	123 (95%)	4 (3%)	2 (2%)	12	11
3	C	95/97 (98%)	90 (95%)	5 (5%)	0	100	100
4	I	49/51 (96%)	48 (98%)	1 (2%)	0	100	100
All	All	281/292 (96%)	269 (96%)	10 (4%)	2 (1%)	26	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	76	SER
2	B	99	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	8/10 (80%)	8 (100%)	0	100	100
2	B	109/109 (100%)	104 (95%)	5 (5%)	33	44
3	C	77/77 (100%)	73 (95%)	4 (5%)	29	38
4	I	45/45 (100%)	44 (98%)	1 (2%)	60	77
All	All	239/241 (99%)	229 (96%)	10 (4%)	36	49

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	62	THR
2	B	79	LYS
2	B	93	LYS
2	B	110	THR
2	B	116	GLN
3	C	160	LEU
3	C	163	LEU
3	C	165	ASN
3	C	203	LYS
4	I	43	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	18	ASN
2	B	48	ASN
2	B	95	ASN
3	C	150	ASN
3	C	165	ASN
3	C	167	ASN
4	I	28	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	10/13 (76%)	-0.59	0 <a href="#">100</a> <a href="#">100</a>	11, 14, 20, 22	0
2	B	131/131 (100%)	-0.45	0 <a href="#">100</a> <a href="#">100</a>	3, 12, 24, 29	0
3	C	97/97 (100%)	-0.22	0 <a href="#">100</a> <a href="#">100</a>	4, 13, 25, 28	0
4	I	51/51 (100%)	-0.30	1 (1%) <a href="#">68</a> <a href="#">75</a>	5, 14, 29, 36	0
All	All	289/292 (98%)	-0.35	1 (0%) <a href="#">94</a> <a href="#">96</a>	3, 13, 25, 36	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	I	45	ASN	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.