



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:51 PM GMT

PDB ID : 1HLM
Title : AMINO ACID SEQUENCE OF A GLOBIN FROM THE SEA CUCUMBER
CAUDINA (MOLPADIA) ARENICOLA
Authors : Hackert, M.L.; Mitchell, D.T.; Ernst, S.R.
Deposited on : 1994-08-26
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	<div> <div>4%</div> <div>45%</div> <div>39%</div> <div>13%</div> <div>.</div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 1297 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMOGLOBIN (CYANO MET).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	159	Total	C	N	O	S	0	0	0
			1247	790	221	230	6			

- Molecule 2 is CYANIDE ION (three-letter code: CYN) (formula: CN).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			2	1	1		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total	O	0	0
			5	5		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HEMOGLOBIN (CYANO MET)



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	77.00Å 77.00Å 61.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	5.00 – 2.90 9.98 – 2.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (5.00-2.90) 70.3 (9.98-2.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.190 , (Not available) 0.184 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	45.5	Xtriage
Anisotropy	0.025	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 130.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ¹	$\langle L \rangle = 0.35$, $\langle L^2 \rangle = 0.18$	Xtriage
Outliers	0 of 2748 reflections	Xtriage
F_o, F_c correlation	0.74	EDS
Total number of atoms	1297	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, CYN, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/1271	1.77	26/1719 (1.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	102	ARG	NE-CZ-NH1	11.09	125.84	120.30
1	A	28	MET	CA-CB-CG	-9.24	97.59	113.30
1	A	141	TRP	CD1-CG-CD2	7.77	112.52	106.30
1	A	76	ARG	NE-CZ-NH1	7.60	124.10	120.30
1	A	137	VAL	CG1-CB-CG2	-7.48	98.93	110.90
1	A	141	TRP	CE2-CD2-CG	-7.38	101.40	107.30
1	A	38	VAL	CG1-CB-CG2	-7.15	99.47	110.90
1	A	24	TRP	CD1-CG-CD2	6.91	111.83	106.30
1	A	123	GLU	CA-CB-CG	6.84	128.44	113.40
1	A	24	TRP	CE2-CD2-CG	-6.70	101.94	107.30
1	A	108	HIS	CA-CB-CG	6.45	124.56	113.60
1	A	131	VAL	CA-CB-CG2	-6.35	101.37	110.90
1	A	3	THR	N-CA-C	-6.31	93.96	111.00
1	A	23	THR	CA-CB-OG1	-5.97	96.46	109.00
1	A	131	VAL	CA-CB-CG1	5.85	119.67	110.90
1	A	21	ARG	NE-CZ-NH2	-5.48	117.56	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	102	ARG	CD-NE-CZ	5.48	131.27	123.60
1	A	23	THR	CA-CB-CG2	5.47	120.06	112.40
1	A	9	VAL	CA-CB-CG2	-5.43	102.76	110.90
1	A	130	GLY	N-CA-C	-5.35	99.72	113.10
1	A	102	ARG	CA-CB-CG	5.29	125.04	113.40
1	A	65	ARG	N-CA-C	-5.23	96.88	111.00
1	A	97	LEU	CA-CB-CG	5.21	127.29	115.30
1	A	151	VAL	CG1-CB-CG2	-5.20	102.58	110.90
1	A	92	VAL	CA-CB-CG2	-5.16	103.16	110.90
1	A	141	TRP	CG-CD2-CE3	5.02	138.42	133.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	84	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1247	0	1251	26	0
2	A	2	0	0	0	0
3	A	43	0	30	3	0
4	A	5	0	0	1	0
All	All	1297	0	1281	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:GLN:HB2	4:A:163:HOH:O	1.90	0.72
1:A:60:SER:HB3	1:A:63:GLU:HG2	1.77	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:THR:HA	1:A:74:ALA:HB1	1.82	0.62
1:A:154:THR:HG23	1:A:155:LYS:H	1.68	0.56
1:A:55:GLN:HG2	1:A:56:MET:SD	2.45	0.56
3:A:159:HEM:HBC2	3:A:159:HEM:HHO	1.86	0.56
1:A:19:LEU:HD23	1:A:136:GLN:HE22	1.73	0.54
1:A:92:VAL:O	1:A:96:LEU:HB2	2.10	0.52
1:A:50:GLN:CD	1:A:61:PRO:HA	2.30	0.52
1:A:50:GLN:NE2	1:A:61:PRO:HA	2.25	0.52
1:A:43:PHE:CD1	1:A:50:GLN:HG2	2.46	0.51
1:A:20:ILE:HD11	1:A:88:MET:SD	2.52	0.50
1:A:16:GLU:HA	1:A:136:GLN:NE2	2.29	0.47
1:A:30:HIS:HB3	1:A:33:GLY:HA3	1.96	0.47
1:A:40:ILE:HA	1:A:43:PHE:HD2	1.81	0.46
1:A:99:THR:O	1:A:102:ARG:HB2	2.15	0.46
1:A:42:ILE:HG23	1:A:116:LEU:HB3	1.98	0.46
1:A:12:LEU:HB3	1:A:16:GLU:HB2	1.99	0.45
1:A:31:ARG:O	1:A:35:VAL:HG23	2.16	0.45
1:A:153:ILE:HA	1:A:156:HIS:HB2	1.99	0.45
1:A:90:THR:O	1:A:92:VAL:N	2.49	0.45
1:A:11:ASP:O	1:A:12:LEU:HB2	2.18	0.44
1:A:138:HIS:CG	1:A:138:HIS:O	2.71	0.43
1:A:109:VAL:HG22	3:A:159:HEM:CBH	2.51	0.41
1:A:75:ILE:O	1:A:78:SER:N	2.53	0.41
1:A:100:LEU:HD11	3:A:159:HEM:C3A	2.56	0.41
1:A:19:LEU:HD13	1:A:19:LEU:HA	1.85	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	157/159 (99%)	113 (72%)	27 (17%)	17 (11%)	<div>01</div>

All (17) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	VAL
1	A	13	THR
1	A	48	THR
1	A	91	GLU
1	A	108	HIS
1	A	131	VAL
1	A	61	PRO
1	A	65	ARG
1	A	67	SER
1	A	149	GLN
1	A	47	PRO
1	A	154	THR
1	A	57	ALA
1	A	12	LEU
1	A	33	GLY
1	A	132	GLY
1	A	151	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	133/133 (100%)	91 (68%)	42 (32%)	0 1

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	THR
1	A	11	ASP
1	A	12	LEU
1	A	14	PRO
1	A	18	ASP
1	A	19	LEU
1	A	21	ARG
1	A	27	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	34	PHE
1	A	35	VAL
1	A	44	HIS
1	A	45	ASN
1	A	51	ARG
1	A	54	PRO
1	A	55	GLN
1	A	56	MET
1	A	61	PRO
1	A	63	GLU
1	A	64	LEU
1	A	65	ARG
1	A	66	THR
1	A	68	ARG
1	A	69	GLN
1	A	70	MET
1	A	76	ARG
1	A	78	SER
1	A	80	LEU
1	A	81	MET
1	A	87	GLU
1	A	89	ASP
1	A	92	VAL
1	A	97	LEU
1	A	102	ARG
1	A	112	LYS
1	A	115	ASP
1	A	116	LEU
1	A	121	LEU
1	A	125	ILE
1	A	126	LYS
1	A	134	THR
1	A	144	THR
1	A	155	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	ASN
1	A	71	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	HEM	A	159	1,2	30,50,50	4.15	12 (40%)	24,82,82	2.40	8 (33%)
2	CYN	A	160	3	0,1,1	0.00	-	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	159	1,2	-	0/10/54/54	0/0/8/8
2	CYN	A	160	3	-	0/0/0/0	0/0/0/0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	159	HEM	C3B-C4B	-13.86	1.39	1.51
3	A	159	HEM	C3D-C4D	-9.49	1.39	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	159	HEM	C2C-C1C	-7.56	1.38	1.52
3	A	159	HEM	C3B-CAB	-6.49	1.39	1.51
3	A	159	HEM	C3C-CAC	-6.02	1.40	1.51
3	A	159	HEM	C2D-C3D	-4.80	1.40	1.54
3	A	159	HEM	C2D-C1D	-4.07	1.38	1.51
3	A	159	HEM	C2B-C1B	-3.63	1.39	1.51
3	A	159	HEM	CHD-C4C	2.11	1.41	1.36
3	A	159	HEM	CBB-CAB	2.78	1.45	1.29
3	A	159	HEM	CBC-CAC	2.91	1.46	1.29
3	A	159	HEM	FE-NC	3.12	2.08	1.95

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	159	HEM	C1D-CHD-C4C	2.16	129.43	125.82
3	A	159	HEM	CAA-CBA-CGA	2.34	117.03	112.75
3	A	159	HEM	CAA-C2A-C1A	2.59	129.82	127.01
3	A	159	HEM	CMD-C2D-C3D	3.04	127.82	114.35
3	A	159	HEM	CAD-C3D-C4D	4.19	127.25	112.47
3	A	159	HEM	CMC-C2C-C3C	4.48	127.71	116.53
3	A	159	HEM	CMB-C2B-C3B	4.78	128.45	116.53
3	A	159	HEM	CAD-C3D-C2D	5.38	128.69	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	159	HEM	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	158/159 (99%)	0.28	7 (4%) 38 32	2, 21, 41, 48	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	9	VAL	5.2
1	A	130	GLY	4.4
1	A	3	THR	3.8
1	A	158	SER	3.2
1	A	156	HIS	3.1
1	A	127	ALA	2.8
1	A	14	PRO	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	HEM	A	159	43/43	0.94	0.17	-0.48	7,20,31,47	0
2	CYN	A	160	2/2	0.99	0.12	-1.18	11,11,11,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.