



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 08:56 PM BST

PDB ID : 2HWT
Title : NMR solution structure of the Master-Rep protein nuclease domain (2-95)
from the Faba Bean Necrotic Yellows Virus
Authors : Vega-Rocha, S.; Gronenborn, B.; Gronenborn, A.M.; Campos-Olivas, R.
Deposited on : 2006-08-02

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment is 91%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	94	 71% 26% ..

2 Ensemble composition and analysis

This entry contains 31 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:95 (93)	0.32	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15, 19, 20, 21, 22, 23, 24, 25, 26, 30, 31
2	2, 17, 27
3	16, 29
Single-model clusters	12; 18; 28

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 1502 atoms, of which 743 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Putative replicase-associated protein.


Mol	Chain	Residues	Atoms						Trace
1	A	94	Total	C	H	N	O	S	0
			1502	479	743	135	139	6	

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Putative replicase-associated protein




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Putative replicase-associated protein



4.2.2 Score per residue for model 2


- Molecule 1: Putative replicase-associated protein






4.2.8 Score per residue for model 8

- Molecule 1: Putative replicase-associated protein




4.2.9 Score per residue for model 9

- Molecule 1: Putative replicase-associated protein




4.2.10 Score per residue for model 10

- Molecule 1: Putative replicase-associated protein



4.2.11 Score per residue for model 11

- Molecule 1: Putative replicase-associated protein



4.2.12 Score per residue for model 12

- Molecule 1: Putative replicase-associated protein






4.2.13 Score per residue for model 13

- Molecule 1: Putative replicase-associated protein

Chain A: 60% 35% ..



4.2.14 Score per residue for model 14

- Molecule 1: Putative replicase-associated protein


Chain A: 66% 30% ..



4.2.15 Score per residue for model 15

- Molecule 1: Putative replicase-associated protein


Chain A: 63% 33% ...



4.2.16 Score per residue for model 16


- Molecule 1: Putative replicase-associated protein

Chain A: 62% 34% ...




4.2.17 Score per residue for model 17

- Molecule 1: Putative replicase-associated protein




4.2.18 Score per residue for model 18

- Molecule 1: Putative replicase-associated protein




4.2.19 Score per residue for model 19

- Molecule 1: Putative replicase-associated protein




4.2.20 Score per residue for model 20

- Molecule 1: Putative replicase-associated protein




4.2.21 Score per residue for model 21

- Molecule 1: Putative replicase-associated protein




4.2.22 Score per residue for model 22

- Molecule 1: Putative replicase-associated protein




4.2.23 Score per residue for model 23

- Molecule 1: Putative replicase-associated protein




4.2.24 Score per residue for model 24

- Molecule 1: Putative replicase-associated protein




4.2.25 Score per residue for model 25

- Molecule 1: Putative replicase-associated protein




4.2.26 Score per residue for model 26

- Molecule 1: Putative replicase-associated protein




4.2.27 Score per residue for model 27

- Molecule 1: Putative replicase-associated protein




4.2.28 Score per residue for model 28

- Molecule 1: Putative replicase-associated protein




4.2.29 Score per residue for model 29

- Molecule 1: Putative replicase-associated protein




4.2.30 Score per residue for model 30

- Molecule 1: Putative replicase-associated protein



4.2.31 Score per residue for model 31

- Molecule 1: Putative replicase-associated protein



5 Refinement protocol and experimental data overview

The models were refined using the following method: *30 conformers calculated using simulated annealing in torsion angle space with CYANA 2.1 (Models 2-31) and average structure after CYANA variable target function minimisation (Model 1).*

Of the 300 calculated structures, 31 were deposited, based on the following criterion: *target function.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.1
CYANA	refinement	2.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 7112
Number of chemical shift lists	1
Total number of shifts	1179
Number of shifts mapped to atoms	1179
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	91%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	754	738	738	12±3
All	All	23374	22878	22878	371

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:72:THR:HG23	1:A:76:ALA:HB2	0.84	1.49	15	1
1:A:32:THR:HG21	1:A:40:ILE:HD12	0.73	1.59	28	7
1:A:12:LEU:HD11	1:A:61:ILE:HG21	0.71	1.59	31	1
1:A:22:HIS:CE1	1:A:60:LEU:HD11	0.67	2.23	8	24
1:A:22:HIS:HB2	1:A:60:LEU:HD21	0.66	1.67	19	31
1:A:72:THR:CG2	1:A:76:ALA:HB2	0.65	2.22	20	3
1:A:32:THR:CG2	1:A:40:ILE:HD12	0.64	2.22	13	11
1:A:22:HIS:CG	1:A:60:LEU:HD21	0.63	2.28	8	19
1:A:5:VAL:HG23	1:A:47:GLU:OE1	0.63	1.93	16	1
1:A:21:LEU:HD22	1:A:94:GLY:HA2	0.63	1.70	16	29
1:A:72:THR:HB	1:A:76:ALA:HB2	0.62	1.70	18	2
1:A:11:THR:HG23	1:A:43:GLN:CG	0.62	2.25	29	12
1:A:29:VAL:HG11	1:A:45:TYR:CZ	0.62	2.30	8	13
1:A:5:VAL:HG11	1:A:50:LYS:HA	0.62	1.71	21	28

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:6:ILE:HG21	1:A:70:ARG:HB3	0.61	1.73	9	1
1:A:72:THR:CB	1:A:76:ALA:HB2	0.61	2.26	18	2
1:A:72:THR:O	1:A:76:ALA:HB3	0.60	1.96	17	2
1:A:12:LEU:CD1	1:A:61:ILE:HG21	0.59	2.26	31	1
1:A:11:THR:HG23	1:A:43:GLN:CD	0.59	2.18	13	17
1:A:11:THR:HG23	1:A:43:GLN:HG3	0.58	1.73	21	11
1:A:6:ILE:HG23	1:A:73:GLN:HG3	0.57	1.76	15	2
1:A:28:LEU:HD12	1:A:29:VAL:N	0.57	2.15	22	24
1:A:6:ILE:HG23	1:A:73:GLN:CG	0.56	2.31	2	2
1:A:72:THR:HG22	1:A:75:GLU:CB	0.55	2.31	5	2
1:A:45:TYR:CE2	1:A:76:ALA:HB1	0.55	2.35	15	1
1:A:29:VAL:HG11	1:A:45:TYR:CE2	0.54	2.37	10	4
1:A:12:LEU:HD12	1:A:13:ASN:N	0.54	2.18	13	7
1:A:12:LEU:HD13	1:A:61:ILE:CG2	0.54	2.33	8	1
1:A:60:LEU:C	1:A:61:ILE:HD13	0.54	2.23	22	2
1:A:34:GLN:HG3	1:A:87:LEU:HD11	0.53	1.79	25	2
1:A:12:LEU:HD22	1:A:15:PRO:HB3	0.53	1.81	31	1
1:A:6:ILE:HG21	1:A:70:ARG:HG2	0.53	1.80	16	7
1:A:21:LEU:HD22	1:A:94:GLY:CA	0.52	2.34	31	4
1:A:29:VAL:HG11	1:A:45:TYR:CE1	0.52	2.40	25	5
1:A:19:LEU:HD12	1:A:30:TYR:CB	0.52	2.35	7	2
1:A:29:VAL:HG21	1:A:45:TYR:CZ	0.51	2.40	29	1
1:A:35:GLY:O	1:A:36:GLU:C	0.51	2.49	14	7
1:A:34:GLN:O	1:A:85:THR:HG22	0.50	2.06	19	2
1:A:22:HIS:HB2	1:A:25:MET:HE3	0.50	1.84	22	8
1:A:46:ILE:HD13	1:A:57:MET:HE1	0.49	1.84	30	1
1:A:11:THR:HG23	1:A:43:GLN:NE2	0.48	2.24	30	7
1:A:34:GLN:CD	1:A:87:LEU:HD11	0.48	2.28	19	3
1:A:11:THR:HG23	1:A:43:GLN:HG2	0.48	1.86	20	3
1:A:22:HIS:CB	1:A:60:LEU:HD21	0.48	2.39	29	10
1:A:22:HIS:CG	1:A:60:LEU:HD11	0.47	2.44	14	2
1:A:60:LEU:CB	1:A:61:ILE:HD12	0.47	2.40	12	8
1:A:16:LEU:HD23	1:A:16:LEU:O	0.46	2.10	27	1
1:A:21:LEU:HD13	1:A:94:GLY:HA3	0.46	1.87	7	2
1:A:31:GLN:HG3	1:A:81:MET:HE3	0.46	1.87	8	2
1:A:19:LEU:HD12	1:A:30:TYR:HB3	0.46	1.86	7	1
1:A:72:THR:OG1	1:A:76:ALA:HB2	0.45	2.10	31	2
1:A:46:ILE:HG21	1:A:57:MET:HE1	0.45	1.87	31	1
1:A:21:LEU:HD13	1:A:94:GLY:CA	0.45	2.41	8	1
1:A:60:LEU:O	1:A:61:ILE:HD13	0.44	2.12	22	1
1:A:45:TYR:CD2	1:A:76:ALA:HB1	0.44	2.47	5	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:31:GLN:HE22	1:A:86:ARG:NE	0.44	2.10	2	1
1:A:22:HIS:ND1	1:A:60:LEU:HD11	0.44	2.27	17	6
1:A:29:VAL:HG21	1:A:45:TYR:CE1	0.43	2.48	8	1
1:A:22:HIS:CD2	1:A:60:LEU:HD11	0.43	2.49	14	3
1:A:12:LEU:CD1	1:A:64:ALA:HB2	0.42	2.45	27	1
1:A:19:LEU:HD22	1:A:61:ILE:HD11	0.42	1.91	19	1
1:A:72:THR:HG22	1:A:75:GLU:HB3	0.42	1.92	5	1
1:A:72:THR:HG22	1:A:75:GLU:HB2	0.42	1.91	8	1
1:A:32:THR:HG23	1:A:40:ILE:HD12	0.42	1.92	31	1
1:A:29:VAL:HG22	1:A:92:GLU:HG3	0.42	1.91	20	1
1:A:29:VAL:HG13	1:A:91:TRP:O	0.41	2.15	21	1
1:A:6:ILE:HG21	1:A:70:ARG:CG	0.41	2.44	16	1
1:A:71:GLY:O	1:A:72:THR:HG23	0.41	2.14	16	1
1:A:19:LEU:HD12	1:A:30:TYR:CG	0.41	2.50	8	1
1:A:79:TYR:O	1:A:81:MET:N	0.41	2.54	6	1
1:A:10:PHE:CE2	1:A:61:ILE:HG21	0.41	2.51	7	1
1:A:25:MET:CE	1:A:46:ILE:HG21	0.40	2.46	21	1
1:A:33:GLU:O	1:A:40:ILE:HG22	0.40	2.17	5	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	92/94 (98%)	79±2 (86±2%)	9±2 (10±2%)	4±1 (4±1%)	6	30
All	All	2852/2914 (98%)	2442 (86%)	284 (10%)	126 (4%)	6	30

All 11 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	50	LYS	31
1	A	80	SER	23
1	A	73	GLN	20
1	A	36	GLU	19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	37	ALA	14
1	A	70	ARG	8
1	A	72	THR	6
1	A	81	MET	2
1	A	82	LYS	1
1	A	3	ARG	1
1	A	39	ASN	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	79/79 (100%)	62±3 (78±4%)	17±3 (22±4%)	4	32
All	All	2449/2449 (100%)	1916 (78%)	533 (22%)	4	32

All 50 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	41	HIS	31
1	A	34	GLN	31
1	A	17	SER	27
1	A	49	LYS	25
1	A	52	THR	25
1	A	26	LYS	24
1	A	87	LEU	24
1	A	43	GLN	23
1	A	85	THR	15
1	A	50	LYS	14
1	A	73	GLN	14
1	A	70	ARG	14
1	A	23	ASP	14
1	A	72	THR	14
1	A	68	LYS	12
1	A	58	LYS	12
1	A	32	THR	11
1	A	67	GLU	11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	81	MET	11
1	A	84	ASP	11
1	A	75	GLU	11
1	A	3	ARG	11
1	A	82	LYS	11
1	A	31	GLN	11
1	A	77	ARG	10
1	A	51	ARG	9
1	A	80	SER	9
1	A	59	LYS	8
1	A	57	MET	8
1	A	69	ARG	8
1	A	83	GLU	8
1	A	33	GLU	7
1	A	79	TYR	7
1	A	24	SER	6
1	A	4	GLN	6
1	A	28	LEU	5
1	A	86	ARG	5
1	A	16	LEU	5
1	A	36	GLU	4
1	A	39	ASN	4
1	A	95	GLU	3
1	A	92	GLU	3
1	A	40	ILE	2
1	A	11	THR	2
1	A	48	MET	2
1	A	13	ASN	1
1	A	12	LEU	1
1	A	25	MET	1
1	A	14	ASN	1
1	A	9	CYS	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 91% for the well-defined parts and 90% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 7112

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1179
Number of shifts mapped to atoms	1179
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	4

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	94	-0.02 ± 0.15	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	85	-0.55 ± 0.23	Should be applied
$^{13}\text{C}'$	93	0.19 ± 0.21	None needed (< 0.5 ppm)
^{15}N	83	0.04 ± 0.45	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 91%, i.e. 1078 atoms were assigned a chemical shift out of a possible 1191. 4 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	444/457 (97%)	176/182 (97%)	185/186 (99%)	83/89 (93%)
Sidechain	529/619 (85%)	334/369 (91%)	187/217 (86%)	8/33 (24%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	105/115 (91%)	55/59 (93%)	42/48 (88%)	8/8 (100%)
Overall	1078/1191 (91%)	565/610 (93%)	414/451 (92%)	99/130 (76%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 90%, i.e. 1083 atoms were assigned a chemical shift out of a possible 1198. 4 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	447/462 (97%)	177/184 (96%)	187/188 (99%)	83/90 (92%)
Sidechain	531/621 (86%)	335/370 (91%)	188/218 (86%)	8/33 (24%)
Aromatic	105/115 (91%)	55/59 (93%)	42/48 (88%)	8/8 (100%)
Overall	1083/1198 (90%)	567/613 (92%)	417/454 (92%)	99/131 (76%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	57	MET	HG2	-0.41	4.23 – 0.63	-7.9
1	A	57	MET	HB3	-0.44	3.70 – 0.30	-7.2
1	A	89	GLY	HA3	1.44	5.80 – 2.00	-6.5
1	A	15	PRO	HG2	0.29	3.48 – 0.38	-5.3

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

