



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 02:49 PM BST

PDB ID : 1I4C
Title : THE SOLUTION STRUCTURE OF THE MINOR FAMILY OF THE MUTANT STEM LOOP C 5'UUA3' TRILOOP OF BROME MOSAIC VIRUS (+) STRAND RNA
Authors : Tinoco Jr., I.; Kim, C.-H.
Deposited on : 2001-02-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
RNA backbone	3027	600

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	13	

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 415 atoms, of which 142 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called RNA (5'-R(*GP*GP*UP*GP*CP*UP*UP*AP*GP*CP*AP*CP*C)-3').


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	13	415	123	142	48	90	12	0

4 Residue-property plots

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: RNA (5'-R(*GP*GP*UP*GP*CP*UP*UP*AP*GP*CP*AP*CP*C)-3')

Chain A: 



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.1
X-PLOR	refinement	3.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5023
Number of chemical shift lists	1
Total number of shifts	10
Number of shifts mapped to atoms	0
Number of unparsed shifts	0
Number of shifts with mapping errors	10
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	0%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	227.47	8/304 (2.6%)	6.20	16/472 (3.4%)
All	All	227.47	8/304 (2.6%)	6.20	16/472 (3.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	5	1
All	All	5	1

All bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	G	C2'-C1'	1711.16	20.35	1.53
1	A	1	G	C4'-C3'	1691.69	20.14	1.53
1	A	1	G	C2'-O2'	1450.94	20.27	1.41
1	A	13	C	C2'-O2'	1404.69	19.67	1.41
1	A	13	C	C3'-O3'	1358.20	20.43	1.42
1	A	1	G	C1'-N9	1262.27	20.42	1.48
1	A	1	G	O5'-C5'	1136.00	19.62	1.44
1	A	10	C	P-OP2	1063.99	19.57	1.49

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	G	N9-C1'-C2'	-82.30	7.01	114.00
1	A	1	G	C8-N9-C1'	-41.86	72.58	127.00
1	A	1	G	C4-N9-C1'	-39.90	74.64	126.50
1	A	13	C	C4'-C3'-O3'	-39.44	26.59	109.40
1	A	1	G	C1'-C2'-O2'	-34.58	6.86	110.60
1	A	13	C	C1'-C2'-O2'	-32.82	12.14	110.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	10	C	O5'-P-OP2	-30.48	74.12	110.70
1	A	9	G	OP2-P-O3'	-27.44	44.84	105.20
1	A	1	G	C3'-C2'-C1'	-26.51	80.29	101.50
1	A	10	C	OP1-P-OP2	-22.92	85.22	119.60
1	A	1	G	C5'-C4'-C3'	-21.05	82.33	116.00
1	A	1	G	C3'-C2'-O2'	-11.23	80.75	113.30
1	A	13	C	C2'-C3'-O3'	-9.84	87.86	109.50
1	A	1	G	C4'-C3'-O3'	-6.41	95.95	109.40
1	A	13	C	C3'-C2'-O2'	-6.35	94.90	113.30
1	A	1	G	O5'-C5'-C4'	-5.60	101.07	111.70

All chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms
1	A	1	G	C4',C2',C3',C1'
1	A	13	C	C3'

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	A	1	G	Sidechain

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	273	142	142	23
All	All	273	142	142	23

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 55.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:1:G:C2'	1:A:1:G:N9	0.74	2.49
1:A:1:G:C1'	1:A:7:U:OP2	0.73	2.36
1:A:5:C:O2'	1:A:6:U:H2'	0.73	1.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:7:U:H3'	1:A:8:A:O4'	0.72	1.84
1:A:4:G:H2'	1:A:5:C:O4'	0.72	1.83
1:A:1:G:C1'	1:A:7:U:H5'	0.71	2.15
1:A:1:G:O2'	1:A:1:G:C1'	0.67	2.43
1:A:10:C:N4	1:A:11:A:N6	0.56	2.53
1:A:7:U:O5'	1:A:7:U:H6	0.55	1.84
1:A:12:C:C4	1:A:13:C:C4	0.54	2.96
1:A:6:U:O2'	1:A:9:G:N2	0.52	2.42
1:A:8:A:O2'	1:A:9:G:N7	0.52	2.42
1:A:9:G:O2'	1:A:10:C:C5	0.51	2.61
1:A:7:U:O5'	1:A:7:U:C6	0.48	2.66
1:A:7:U:O2'	1:A:8:A:H5'	0.48	2.09
1:A:12:C:H2'	1:A:13:C:O4'	0.48	2.08
1:A:10:C:C4	1:A:11:A:N7	0.48	2.82
1:A:5:C:O2'	1:A:6:U:C2'	0.46	2.60
1:A:4:G:C6	1:A:5:C:N3	0.45	2.84
1:A:6:U:O2'	1:A:10:C:O2	0.42	2.36
1:A:9:G:C4	1:A:10:C:C4	0.41	3.07
1:A:7:U:H5''	1:A:8:A:N3	0.41	2.31
1:A:9:G:O2'	1:A:10:C:OP1	0.40	2.39

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	11/13 (85%)	7 (64%)	3 (27%)	0.13
All	All	11/13 (85%)	7 (64%)	3 (27%)	0.13

The overall RNA backbone suiteness is 0.12.

All RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	5	C
1	A	8	A
1	A	3	U
1	A	7	U
1	A	6	U
1	A	9	G
1	A	10	C

All RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	5	C
1	A	8	A
1	A	1	G

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 0% for the well-defined parts and 0% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 5023

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	10
Number of shifts mapped to atoms	0
Number of unparsed shifts	0
Number of shifts with mapping errors	10
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Chain not found in structure. All 10 occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
UNMAPPED	3	U	H2'	3.9	-1.0	1
UNMAPPED	1	G	H3'	4.353	-1.0	1
UNMAPPED	4	G	H3'	3.798	-1.0	1
UNMAPPED	5	C	H3'	4.387	-1.0	1
UNMAPPED	1	G	H2'	4.501	-1.0	1
UNMAPPED	2	G	H2'	4.134	-1.0	1
UNMAPPED	5	C	H2'	4.157	-1.0	1
UNMAPPED	4	G	H2'	4.086	-1.0	1
UNMAPPED	2	G	H3'	4.23	-1.0	1
UNMAPPED	3	U	H3'	3.705	-1.0	1

7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 245. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	0/245 (0%)	0/141 (0%)	0/87 (0%)	0/17 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 245. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	0/245 (0%)	0/141 (0%)	0/87 (0%)	0/17 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned_chem_shift_list_1). RCI is only applicable to proteins.