



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2016 – 01:14 AM EDT

PDB ID : 4ICG  
Title : N-terminal dimerization domain of H-NS in complex with Hha (Salmonella Typhimurium)  
Authors : Ali, S.S.; Whitney, J.C.; Stevenson, J.; Robinson, H.; Howell, P.L.; Navarre, W.W.  
Deposited on : 2012-12-10  
Resolution : 2.92 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027457
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1643 (2.94-2.90)
Clashscore	102246	1871 (2.94-2.90)
Ramachandran outliers	100387	1824 (2.94-2.90)
Sidechain outliers	100360	1826 (2.94-2.90)
RSRZ outliers	91569	1650 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	46	<div> <div>2%</div> <div>61% 35% .</div> </div>
1	B	46	<div> <div>2%</div> <div>63% 30% . .</div> </div>
2	C	75	<div> <div>%</div> <div>52% 35% . 11%</div> </div>
2	D	75	<div> <div>68% 21% . 9%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1688 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-binding protein H-NS.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	44	Total	C	N	O	S	Se	0	0	0
			337	204	60	71	1	1			
1	A	44	Total	C	N	O	S	Se	0	0	0
			342	209	64	67	1	1			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MSE	-	EXPRESSION TAG	UNP P0A1S2
B	2	GLY	-	EXPRESSION TAG	UNP P0A1S2
A	1	MSE	-	EXPRESSION TAG	UNP P0A1S2
A	2	GLY	-	EXPRESSION TAG	UNP P0A1S2

- Molecule 2 is a protein called Hemolysin expression modulating protein (Involved in environmental regulation of virulence factors).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	67	Total	C	N	O	S	Se	0	0	0
			506	324	86	93	1	2			
2	D	68	Total	C	N	O	S	Se	0	0	0
			503	319	88	93	1	2			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	EXPRESSION TAG	UNP Q7CR17
C	-1	SER	-	EXPRESSION TAG	UNP Q7CR17
C	0	HIS	-	EXPRESSION TAG	UNP Q7CR17
C	1	MSE	-	EXPRESSION TAG	UNP Q7CR17
D	-2	GLY	-	EXPRESSION TAG	UNP Q7CR17
D	-1	SER	-	EXPRESSION TAG	UNP Q7CR17
D	0	HIS	-	EXPRESSION TAG	UNP Q7CR17

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	1	MSE	-	EXPRESSION TAG	UNP Q7CR17


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-binding protein H-NS



- Molecule 1: DNA-binding protein H-NS



- Molecule 2: Hemolysin expression modulating protein (Involved in environmental regulation of virulence factors)



- Molecule 2: Hemolysin expression modulating protein (Involved in environmental regulation of virulence factors)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.88Å 82.90Å 47.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.13 – 2.92 47.13 – 2.92	Depositor EDS
% Data completeness (in resolution range)	88.0 (47.13-2.92) 87.2 (47.13-2.92)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	10.59 (at 2.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.277 , 0.332 0.273 , 0.332	Depositor DCC
$R_{free}$ test set	620 reflections (10.13%)	DCC
Wilson B-factor (Å <sup>2</sup> )	66.1	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 54.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	1688	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/340	0.58	0/454
1	B	0.35	0/335	0.57	0/448
2	C	0.35	0/513	0.55	0/696
2	D	0.31	0/510	0.51	0/690
All	All	0.34	0/1698	0.55	0/2288

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	342	0	345	13	0
1	B	337	0	319	15	0
2	C	506	0	450	18	0
2	D	503	0	439	11	0
All	All	1688	0	1553	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 47 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:24:LEU:HA	2:D:27:VAL:HG12	1.69	0.72
2:C:8:LYS:NZ	1:A:28:GLU:OE2	2.23	0.72
2:D:20:THR:O	2:D:22:ASP:N	2.21	0.72
2:C:20:THR:O	2:C:22:ASP:N	2.22	0.72
1:B:21:CYS:O	1:A:40:ARG:NH2	2.29	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	42/46 (91%)	38 (90%)	3 (7%)	1 (2%)	7	28
1	B	42/46 (91%)	38 (90%)	4 (10%)	0	100	100
2	C	65/75 (87%)	62 (95%)	2 (3%)	1 (2%)	13	41
2	D	66/75 (88%)	63 (96%)	2 (3%)	1 (2%)	13	41
All	All	215/242 (89%)	201 (94%)	11 (5%)	3 (1%)	14	43

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	21	ILE
2	D	21	ILE
1	A	45	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	35/39 (90%)	34 (97%)	1 (3%)	50	83
1	B	33/39 (85%)	32 (97%)	1 (3%)	48	82
2	C	46/67 (69%)	43 (94%)	3 (6%)	21	51
2	D	43/67 (64%)	41 (95%)	2 (5%)	32	68
All	All	157/212 (74%)	150 (96%)	7 (4%)	34	69

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	44	TYR
2	D	44	TYR
1	A	12	ARG
2	C	16	ARG
2	D	14	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	43/46 (93%)	0.02	1 (2%) 64 59	37, 68, 98, 127	0
1	B	43/46 (93%)	0.17	1 (2%) 64 59	40, 65, 101, 111	0
2	C	65/75 (86%)	0.01	1 (1%) 76 74	43, 72, 95, 108	0
2	D	66/75 (88%)	0.01	0 100 100	60, 89, 116, 124	0
All	All	217/242 (89%)	0.04	3 (1%) 78 76	37, 75, 109, 127	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	44	GLU	6.2
2	C	60	TYR	3.1
1	A	46	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.