



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:28 PM GMT

PDB ID : 4IEF  
Title : Complex of Porphyromonas gingivalis RgpB pro- and mature domains  
Authors : de Diego, I.; Veillard, F.T.; Guevara, T.; Potempa, B.; Sztukowska, M.;  
Potempa, J.; Gomis-Ruth, F.X.  
Deposited on : 2012-12-13  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	<div> <div>5%</div> <div>82% 9% 9%</div> </div>
1	C	210	<div> <div>8%</div> <div>85% 8% 7%</div> </div>
1	E	210	<div> <div>5%</div> <div>82% 10% 8%</div> </div>
1	G	210	<div> <div>10%</div> <div>84% 6% 10%</div> </div>
2	B	439	<div> <div>2%</div> <div>88% 8% .</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	D	439	
2	F	439	
2	H	439	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	TRS	B	706	-	-	-	X
9	GOL	H	706	-	-	-	X

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 20046 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gingipain R2 Pro-Domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	192	Total	C	N	O	S	0	0	0
			1504	958	250	292	4			
1	C	195	Total	C	N	O	S	0	0	0
			1527	972	256	295	4			
1	E	194	Total	C	N	O	S	0	0	0
			1516	966	252	294	4			
1	G	190	Total	C	N	O	S	0	0	0
			1492	951	248	289	4			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	GLY	-	EXPRESSION TAG	UNP P95493
A	21	PRO	-	EXPRESSION TAG	UNP P95493
A	22	LEU	-	EXPRESSION TAG	UNP P95493
A	23	GLY	-	EXPRESSION TAG	UNP P95493
A	24	SER	-	EXPRESSION TAG	UNP P95493
C	20	GLY	-	EXPRESSION TAG	UNP P95493
C	21	PRO	-	EXPRESSION TAG	UNP P95493
C	22	LEU	-	EXPRESSION TAG	UNP P95493
C	23	GLY	-	EXPRESSION TAG	UNP P95493
C	24	SER	-	EXPRESSION TAG	UNP P95493
E	20	GLY	-	EXPRESSION TAG	UNP P95493
E	21	PRO	-	EXPRESSION TAG	UNP P95493
E	22	LEU	-	EXPRESSION TAG	UNP P95493
E	23	GLY	-	EXPRESSION TAG	UNP P95493
E	24	SER	-	EXPRESSION TAG	UNP P95493
G	20	GLY	-	EXPRESSION TAG	UNP P95493
G	21	PRO	-	EXPRESSION TAG	UNP P95493
G	22	LEU	-	EXPRESSION TAG	UNP P95493
G	23	GLY	-	EXPRESSION TAG	UNP P95493
G	24	SER	-	EXPRESSION TAG	UNP P95493

- Molecule 2 is a protein called Gingipain R2 Mature Domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	423	Total	C	N	O	S	0	0	0
			3272	2058	548	649	17			
2	D	424	Total	C	N	O	S	0	0	0
			3280	2062	550	651	17			
2	F	424	Total	C	N	O	S	0	0	0
			3280	2062	550	651	17			
2	H	424	Total	C	N	O	S	0	0	0
			3280	2062	550	651	17			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	663	HIS	-	EXPRESSION TAG	UNP P95493
B	664	HIS	-	EXPRESSION TAG	UNP P95493
B	665	HIS	-	EXPRESSION TAG	UNP P95493
B	666	HIS	-	EXPRESSION TAG	UNP P95493
B	667	HIS	-	EXPRESSION TAG	UNP P95493
B	668	HIS	-	EXPRESSION TAG	UNP P95493
D	663	HIS	-	EXPRESSION TAG	UNP P95493
D	664	HIS	-	EXPRESSION TAG	UNP P95493
D	665	HIS	-	EXPRESSION TAG	UNP P95493
D	666	HIS	-	EXPRESSION TAG	UNP P95493
D	667	HIS	-	EXPRESSION TAG	UNP P95493
D	668	HIS	-	EXPRESSION TAG	UNP P95493
F	663	HIS	-	EXPRESSION TAG	UNP P95493
F	664	HIS	-	EXPRESSION TAG	UNP P95493
F	665	HIS	-	EXPRESSION TAG	UNP P95493
F	666	HIS	-	EXPRESSION TAG	UNP P95493
F	667	HIS	-	EXPRESSION TAG	UNP P95493
F	668	HIS	-	EXPRESSION TAG	UNP P95493
H	663	HIS	-	EXPRESSION TAG	UNP P95493
H	664	HIS	-	EXPRESSION TAG	UNP P95493
H	665	HIS	-	EXPRESSION TAG	UNP P95493
H	666	HIS	-	EXPRESSION TAG	UNP P95493
H	667	HIS	-	EXPRESSION TAG	UNP P95493
H	668	HIS	-	EXPRESSION TAG	UNP P95493

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0

- Molecule 4 is BARIUM ION (three-letter code: BA) (formula: Ba).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	1	Total Ba 1 1	0	0
4	B	1	Total Ba 1 1	0	0
4	D	1	Total Ba 1 1	0	0
4	F	1	Total Ba 1 1	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	H	3	Total Ca 3 3	0	0
5	B	4	Total Ca 4 4	0	0
5	D	3	Total Ca 3 3	0	0
5	F	4	Total Ca 4 4	0	0

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	F	3	Total	Na	0	0
			3	3		

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	H	1	Total	Mg	0	0
			1	1		

- Molecule 9 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	68	Total	O	0	0
			68	68		
10	B	179	Total	O	0	0
			179	179		
10	C	62	Total	O	0	0
			62	62		
10	D	139	Total	O	0	0
			139	139		
10	E	69	Total	O	0	0
			69	69		
10	F	156	Total	O	0	0
			156	156		
10	G	43	Total	O	0	0
			43	43		
10	H	142	Total	O	0	0
			142	142		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gingipain R2 Pro-Domain




- Molecule 1: Gingipain R2 Pro-Domain




- Molecule 1: Gingipain R2 Pro-Domain





- Molecule 1: Gingipain R2 Pro-Domain









• Molecule 2: Gingipain R2 Mature Domain





• Molecule 2: Gingipain R2 Mature Domain



• Molecule 2: Gingipain R2 Mature Domain



• Molecule 2: Gingipain R2 Mature Domain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.42Å 133.14Å 109.83Å 90.00° 90.52° 90.00°	Depositor
Resolution (Å)	33.67 – 2.30 33.61 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.2 (33.67-2.30) 99.3 (33.61-2.30)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.68 (at 2.29Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, $R_{free}$	0.189 , 0.225 0.192 , 0.232	Depositor DCC
$R_{free}$ test set	5323 reflections (5.25%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.5	Xtriage
Anisotropy	0.266	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 47.7	EDS
Estimated twinning fraction	0.027 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 106784 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	20046	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, BA, CL, CSD, CA, NA, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1531	0.60	0/2075
1	C	0.40	0/1554	0.60	0/2105
1	E	0.39	0/1543	0.59	0/2091
1	G	0.39	0/1519	0.59	0/2058
2	B	0.41	0/3324	0.62	0/4508
2	D	0.40	0/3332	0.63	0/4519
2	F	0.41	0/3332	0.62	0/4519
2	H	0.40	0/3332	0.62	0/4519
All	All	0.40	0/19467	0.61	0/26394

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1504	0	1513	10	0
1	C	1527	0	1540	9	0
1	E	1516	0	1527	13	0
1	G	1492	0	1501	7	0
2	B	3272	0	3219	18	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	3280	0	3225	22	0
2	F	3280	0	3225	26	0
2	H	3280	0	3225	17	0
3	A	1	0	0	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
4	F	1	0	0	0	0
4	H	1	0	0	0	0
5	B	4	0	0	0	0
5	D	3	0	0	0	0
5	F	4	0	0	0	0
5	H	3	0	0	0	0
6	B	8	0	12	0	0
7	F	3	0	0	0	0
8	H	1	0	0	0	0
9	H	6	0	6	0	0
10	A	68	0	0	3	0
10	B	179	0	0	1	0
10	C	62	0	0	1	0
10	D	139	0	0	3	0
10	E	69	0	0	5	0
10	F	156	0	0	3	0
10	G	43	0	0	1	0
10	H	142	0	0	3	0
All	All	20046	0	18993	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 105 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:217:ILE:HD11	2:D:535:ILE:HG21	1.59	0.85
1:G:103:GLU:HG3	10:G:319:HOH:O	1.86	0.75
2:F:452:GLY:H	2:F:455:HIS:HD2	1.39	0.69
2:F:537:ARG:HD2	2:F:649:TYR:HB2	1.74	0.68
2:B:452:GLY:H	2:B:455:HIS:HD2	1.40	0.68

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	188/210 (90%)	182 (97%)	6 (3%)	0	100	100
1	C	191/210 (91%)	186 (97%)	5 (3%)	0	100	100
1	E	190/210 (90%)	183 (96%)	7 (4%)	0	100	100
1	G	186/210 (89%)	180 (97%)	6 (3%)	0	100	100
2	B	420/439 (96%)	413 (98%)	5 (1%)	2 (0%)	34	41
2	D	421/439 (96%)	412 (98%)	6 (1%)	3 (1%)	26	31
2	F	421/439 (96%)	410 (97%)	9 (2%)	2 (0%)	34	41
2	H	421/439 (96%)	413 (98%)	5 (1%)	3 (1%)	26	31
All	All	2438/2596 (94%)	2379 (98%)	49 (2%)	10 (0%)	39	48

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	449	SER
2	B	474	VAL
2	D	449	SER
2	F	449	SER
2	F	474	VAL

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/184 (92%)	164 (96%)	6 (4%)	43	58

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	172/184 (94%)	167 (97%)	5 (3%)	50	66
1	E	171/184 (93%)	164 (96%)	7 (4%)	37	50
1	G	169/184 (92%)	166 (98%)	3 (2%)	66	82
2	B	357/372 (96%)	351 (98%)	6 (2%)	68	83
2	D	358/372 (96%)	351 (98%)	7 (2%)	63	79
2	F	358/372 (96%)	351 (98%)	7 (2%)	63	79
2	H	358/372 (96%)	350 (98%)	8 (2%)	60	77
All	All	2113/2224 (95%)	2064 (98%)	49 (2%)	58	75

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	616	ASP
1	E	60	GLN
2	H	481	ASN
1	E	31	ARG
1	E	131	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	590	ASN
2	F	464	GLN
2	H	475	ASN
2	F	282	ASN
2	F	455	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	CSD	B	473	2	3,7,8	0.67	0	3,8,10	2.78	1 (33%)
2	CSD	D	473	2	3,7,8	0.72	0	3,8,10	2.94	1 (33%)
2	CSD	F	473	2	3,7,8	0.77	0	3,8,10	2.74	1 (33%)
2	CSD	H	473	2	3,7,8	0.74	0	3,8,10	2.83	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSD	B	473	2	-	0/2/6/8	0/0/0/0
2	CSD	D	473	2	-	0/2/6/8	0/0/0/0
2	CSD	F	473	2	-	0/2/6/8	0/0/0/0
2	CSD	H	473	2	-	0/2/6/8	0/0/0/0

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	473	CSD	CB-CA-C	-4.71	98.56	111.46
2	H	473	CSD	CB-CA-C	-4.47	99.21	111.46
2	F	473	CSD	CB-CA-C	-4.38	99.46	111.46
2	B	473	CSD	CB-CA-C	-4.36	99.51	111.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	473	CSD	1	0
2	F	473	CSD	1	0
2	H	473	CSD	1	0



## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 25 ligands modelled in this entry, 23 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	TRS	B	706	-	7,7,7	0.84	1 (14%)	9,9,9	1.73	3 (33%)
9	GOL	H	706	5	5,5,5	0.33	0	5,5,5	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	TRS	B	706	-	-	0/9/9/9	0/0/0/0
9	GOL	H	706	5	-	0/4/4/4	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	706	TRS	C-N	-2.09	1.47	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	706	TRS	C3-C-N	-2.57	103.41	108.09
6	B	706	TRS	C3-C-C2	2.09	115.31	110.78
6	B	706	TRS	O2-C2-C	2.59	116.43	111.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	192/210 (91%)	0.14	11 (5%) 27 36	26, 38, 59, 97	0
1	C	195/210 (92%)	0.27	16 (8%) 14 20	23, 42, 68, 103	0
1	E	194/210 (92%)	0.33	11 (5%) 27 36	28, 43, 70, 101	0
1	G	190/210 (90%)	0.49	20 (10%) 8 12	31, 45, 67, 109	0
2	B	422/439 (96%)	0.03	8 (1%) 70 76	19, 32, 55, 80	0
2	D	423/439 (96%)	0.22	22 (5%) 31 39	19, 35, 64, 87	0
2	F	423/439 (96%)	0.22	21 (4%) 32 41	21, 35, 56, 92	0
2	H	423/439 (96%)	0.05	13 (3%) 52 62	20, 34, 57, 76	0
All	All	2462/2596 (94%)	0.19	122 (4%) 32 41	19, 37, 63, 109	0

The worst 5 of 122 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	196	ALA	7.2
1	A	227	ALA	6.0
2	D	594	SER	5.6
1	G	197	GLU	5.5
2	F	532	PRO	5.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CSD	D	473	8/9	0.94	0.11	-	28,30,39,41	0
2	CSD	F	473	8/9	0.98	0.08	-	30,32,36,38	0
2	CSD	H	473	8/9	0.95	0.11	-	29,31,36,38	0
2	CSD	B	473	8/9	0.95	0.09	-	28,29,36,36	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
6	TRS	B	706	8/8	0.65	0.26	5.39	54,55,56,56	0
9	GOL	H	706	6/6	0.91	0.16	4.10	39,42,46,49	0
5	CA	B	702	1/1	0.99	0.11	-0.27	27,27,27,27	0
5	CA	D	703	1/1	0.99	0.12	-0.52	23,23,23,23	0
8	MG	H	705	1/1	0.93	0.13	-0.84	50,50,50,50	0
5	CA	B	703	1/1	0.99	0.10	-1.08	25,25,25,25	0
5	CA	H	703	1/1	0.98	0.09	-1.32	31,31,31,31	0
5	CA	F	702	1/1	0.99	0.09	-1.86	32,32,32,32	0
5	CA	F	703	1/1	0.99	0.05	-2.02	36,36,36,36	0
3	CL	A	301	1/1	0.99	0.04	-3.07	34,34,34,34	0
5	CA	D	702	1/1	0.99	0.07	-3.34	28,28,28,28	0
4	BA	F	701	1/1	0.99	0.06	-3.76	59,59,59,59	0
4	BA	H	701	1/1	0.98	0.05	-4.24	67,67,67,67	0
4	BA	B	701	1/1	1.00	0.06	-4.69	55,55,55,55	0
5	CA	H	702	1/1	0.99	0.03	-5.60	35,35,35,35	0
7	NA	F	707	1/1	0.94	0.11	-	63,63,63,63	0
5	CA	H	704	1/1	0.99	0.16	-	44,44,44,44	0
5	CA	F	704	1/1	0.98	0.18	-	54,54,54,54	0
5	CA	B	705	1/1	0.97	0.24	-	44,44,44,44	0
5	CA	F	705	1/1	0.97	0.25	-	55,55,55,55	0
5	CA	D	704	1/1	0.96	0.12	-	42,42,42,42	0
5	CA	B	704	1/1	0.78	0.13	-	51,51,51,51	0
7	NA	F	708	1/1	0.94	0.34	-	42,42,42,42	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	BA	D	701	1/1	1.00	0.05	-	63,63,63,63	0
7	NA	F	706	1/1	0.94	0.23	-	38,38,38,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.