



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:44 AM GMT

PDB ID : 2IH8  
Title : A low-dose crystal structure of a recombinant Melanocarpus albomyces laccase  
Authors : Hakulinen, N.; Rouvinen, J.  
Deposited on : 2006-09-26  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	559	<div> <div></div> <div>97%</div> </div>
1	B	559	<div> <div></div> <div>95%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	SO4	A	801	-	-	-	X
2	NAG	A	700	-	-	-	X
2	NAG	A	760	-	-	-	X
2	NAG	B	700	-	-	-	X
2	NAG	B	750	-	-	-	X
2	NAG	B	760	-	-	-	X
4	NAG	B	731	-	-	-	X
6	NDG	A	770	-	-	-	X

## 2 Entry composition [i](#)


There are 12 unique types of molecules in this entry. The entry contains 10333 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Laccase-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	559	Total	C	N	O	S	0	0	0
			4369	2764	759	831	15			
1	B	559	Total	C	N	O	S	0	0	0
			4369	2764	759	831	15			

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	5	Total	C	N	O	0	0
			61	34	2	25		


- Molecule 4 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	3	Total	C	N	O	0	0
			39	22	2	15		
4	A	3	Total	C	N	O	0	0
			39	22	2	15		
4	B	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	2	Total	C	N	O	0	0
			28	16	2	10		
5	B	2	Total	C	N	O	0	0
			28	16	2	10		
5	B	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 6 is SUGAR (2-(ACETYLAMINO)-2-DEOXY-A-D-GLUCOPYRANOSE) (three-letter code: NDG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	4	Total	C	N	O	0	0
			50	28	2	20		


- Molecule 8 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	4	Total	Cu	0	0
			4	4		
8	A	4	Total	Cu	0	0
			4	4		

- Molecule 9 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	1	Total	Cl	0	0
			1	1		
9	A	1	Total	Cl	0	0
			1	1		

- Molecule 10 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	O	S	0	0
			5	4	1		
10	A	1	Total	O	S	0	0
			5	4	1		
10	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 11 is OXYGEN MOLECULE (three-letter code: OXY) (formula: O<sub>2</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	1	Total	O	0	0
			2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	B	1	Total	O	0	0
			2	2		

- Molecule 12 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	600	Total	O	0	0
			600	600		
12	B	556	Total	O	0	0
			556	556		




### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Laccase-1



#### • Molecule 1: Laccase-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	173.06 Å   61.81 Å   123.91 Å 90.00°   96.36°   90.00°	Depositor
Resolution (Å)	25.00 – 2.00 19.59 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (25.00-2.00) 100.0 (19.59-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.30 (at 2.01 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.183   ,   0.213 0.202   ,   0.227	Depositor DCC
$R_{free}$ test set	4409 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.9	Xtriage
Anisotropy	0.159	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 40.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 88169 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	10333	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, CL, NDG, OXY, SO4, CU, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.62	0/4506	0.74	0/6191
1	B	0.62	0/4506	0.74	2/6191 (0.0%)
All	All	0.62	0/9012	0.74	2/12382 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	40	LEU	CA-CB-CG	5.20	127.27	115.30
1	B	288	PHE	N-CA-C	-5.03	97.42	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	546	TYR	Sidechain
1	B	546	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4369	0	4110	6	0
1	B	4369	0	4111	10	0
2	A	42	0	39	0	0
2	B	42	0	39	0	0
3	A	61	0	52	0	0
4	A	78	0	68	2	0
4	B	39	0	34	1	0
5	A	28	0	25	0	0
5	B	56	0	50	1	0
6	A	14	0	13	1	0
7	B	50	0	43	0	0
8	A	4	0	0	0	0
8	B	4	0	0	0	0
9	A	1	0	0	0	0
9	B	1	0	0	0	0
10	A	10	0	0	0	0
10	B	5	0	0	0	0
11	A	2	0	0	0	0
11	B	2	0	0	0	0
12	A	600	0	0	2	0
12	B	556	0	0	2	0
All	All	10333	0	8584	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:770:NDG:H6C2	12:A:1387:HOH:O	1.97	0.61
1:B:380:ILE:HG21	1:B:514:SER:HB3	1.87	0.55
1:A:251:ALA:HB3	1:A:274:ASP:HB2	1.89	0.55
1:A:311:HIS:CD2	4:A:720:NAG:H5	2.42	0.54
1:B:323:GLU:OE2	4:B:731:NAG:H4	2.07	0.54
1:A:40:LEU:HD22	1:A:60:ILE:HG12	1.90	0.53

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:311:HIS:CD2	5:B:720:NAG:H5	2.44	0.51
1:A:54:LYS:HD2	1:A:177:TYR:HA	1.93	0.51
1:A:380:ILE:HG21	1:A:514:SER:HB3	1.93	0.50
1:B:40:LEU:HD22	1:B:60:ILE:HG12	1.94	0.50
1:A:48:GLY:HA3	12:A:802:HOH:O	2.14	0.47
1:B:251:ALA:HB3	1:B:274:ASP:HB2	1.96	0.46
1:B:426:PRO:HD2	12:B:1049:HOH:O	2.18	0.43
1:B:48:GLY:HA3	12:B:956:HOH:O	2.18	0.43
1:B:137:TYR:CZ	1:B:151:GLY:HA3	2.54	0.42
1:B:198:VAL:HG11	1:B:290:VAL:HG11	2.01	0.42
4:A:721:NAG:H4	4:A:722:BMA:H2	1.89	0.41
1:B:173:THR:HB	1:B:232:THR:OG1	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	557/559 (100%)	543 (98%)	14 (2%)	0	100	100
1	B	557/559 (100%)	542 (97%)	15 (3%)	0	100	100
All	All	1114/1118 (100%)	1085 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	478/478 (100%)	471 (98%)	7 (2%)	72	75
1	B	478/478 (100%)	469 (98%)	9 (2%)	65	67
All	All	956/956 (100%)	940 (98%)	16 (2%)	68	71

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	PRO
1	A	88	ASN
1	A	109	ASN
1	A	123	ARG
1	A	140	HIS
1	A	274	ASP
1	A	513	LEU
1	B	7	PRO
1	B	40	LEU
1	B	88	ASN
1	B	104	LEU
1	B	109	ASN
1	B	123	ARG
1	B	140	HIS
1	B	465	LEU
1	B	513	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	GLN
1	A	144	GLN
1	A	213	GLN
1	B	72	ASN
1	B	99	GLN
1	B	144	GLN
1	B	213	GLN
1	B	236	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

24 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	710	1,3	14,14,15	0.44	0	15,19,21	0.69	0
3	NAG	A	711	3	14,14,15	0.49	0	15,19,21	0.54	0
3	BMA	A	712	3	11,11,12	0.58	0	14,15,17	0.43	0
3	MAN	A	713	3	11,11,12	0.52	0	14,15,17	0.63	1 (7%)
3	MAN	A	714	3	11,11,12	0.51	0	14,15,17	0.61	1 (7%)
4	NAG	A	720	1,4	14,14,15	0.45	0	15,19,21	0.64	0
4	NAG	A	721	4	14,14,15	0.56	0	15,19,21	0.61	0
4	BMA	A	722	4	11,11,12	0.50	0	14,15,17	0.40	0
4	NAG	A	730	1,4	14,14,15	0.65	0	15,19,21	0.57	0
4	NAG	A	731	4	14,14,15	0.49	0	15,19,21	0.71	0
4	BMA	A	732	4	11,11,12	0.55	0	14,15,17	0.26	0
5	NAG	A	740	1,5	14,14,15	0.52	0	15,19,21	0.75	0
5	NAG	A	741	5	14,14,15	0.47	0	15,19,21	0.81	1 (6%)
7	NAG	B	710	1,7	14,14,15	0.44	0	15,19,21	0.66	0
7	NAG	B	711	7	14,14,15	0.45	0	15,19,21	0.64	0
7	BMA	B	712	7	11,11,12	0.52	0	14,15,17	0.23	0
7	MAN	B	714	7	11,11,12	0.52	0	14,15,17	0.60	1 (7%)
5	NAG	B	720	1,5	14,14,15	0.49	0	15,19,21	0.64	0
5	NAG	B	721	5	14,14,15	0.45	0	15,19,21	0.62	0
4	NAG	B	730	1,4	14,14,15	0.61	0	15,19,21	0.61	0
4	NAG	B	731	4	14,14,15	0.85	1 (7%)	15,19,21	1.52	2 (13%)
4	BMA	B	732	4	11,11,12	0.44	0	14,15,17	0.64	0
5	NAG	B	740	1,5	14,14,15	0.50	0	15,19,21	0.71	0
5	NAG	B	741	5	14,14,15	0.50	0	15,19,21	0.68	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	710	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	711	3	-	0/6/23/26	0/1/1/1
3	BMA	A	712	3	-	0/2/19/22	0/1/1/1
3	MAN	A	713	3	-	0/2/19/22	0/1/1/1
3	MAN	A	714	3	-	0/2/19/22	0/1/1/1
4	NAG	A	720	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	721	4	-	0/6/23/26	0/1/1/1
4	BMA	A	722	4	-	0/2/19/22	0/1/1/1
4	NAG	A	730	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	731	4	-	0/6/23/26	0/1/1/1
4	BMA	A	732	4	-	0/2/19/22	0/1/1/1
5	NAG	A	740	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	741	5	-	0/6/23/26	0/1/1/1
7	NAG	B	710	1,7	-	0/6/23/26	0/1/1/1
7	NAG	B	711	7	-	0/6/23/26	0/1/1/1
7	BMA	B	712	7	-	0/2/19/22	0/1/1/1
7	MAN	B	714	7	-	0/2/19/22	0/1/1/1
5	NAG	B	720	1,5	-	0/6/23/26	0/1/1/1
5	NAG	B	721	5	-	0/6/23/26	0/1/1/1
4	NAG	B	730	1,4	-	0/6/23/26	0/1/1/1
4	NAG	B	731	4	-	0/6/23/26	0/1/1/1
4	BMA	B	732	4	-	0/2/19/22	0/1/1/1
5	NAG	B	740	1,5	-	0/6/23/26	0/1/1/1
5	NAG	B	741	5	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	731	NAG	C1-C2	2.52	1.55	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	731	NAG	C4-C3-C2	-4.03	104.96	111.23
5	A	741	NAG	C2-N2-C7	-2.48	119.85	123.04
5	B	741	NAG	C2-N2-C7	-2.03	120.43	123.04
3	A	714	MAN	C1-O5-C5	2.03	114.83	112.25
7	B	714	MAN	C1-O5-C5	2.06	114.86	112.25
4	B	731	NAG	O4-C4-C3	2.17	115.23	110.34
3	A	713	MAN	C1-O5-C5	2.20	115.04	112.25



There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	720	NAG	1	0
4	A	721	NAG	1	0
4	A	722	BMA	1	0
5	B	720	NAG	1	0
4	B	731	NAG	1	0

## 5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 10 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
11	OXY	A	620	8	1,1,1	0.10	0	0,0,0	0.00	-
2	NAG	A	700	1	14,14,15	0.51	0	15,19,21	0.62	0
2	NAG	A	750	1	14,14,15	0.45	0	15,19,21	0.79	1 (6%)
2	NAG	A	760	1	14,14,15	0.56	0	15,19,21	0.66	0
6	NDG	A	770	1	14,14,15	0.99	1 (7%)	15,19,21	1.01	1 (6%)
10	SO4	A	800	-	4,4,4	0.18	0	6,6,6	0.09	0
10	SO4	A	801	-	4,4,4	0.15	0	6,6,6	0.12	0
11	OXY	B	620	8	1,1,1	0.09	0	0,0,0	0.00	-
2	NAG	B	700	1	14,14,15	0.54	0	15,19,21	0.64	0
2	NAG	B	750	1	14,14,15	0.56	0	15,19,21	0.63	0
2	NAG	B	760	1	14,14,15	0.49	0	15,19,21	0.62	0
10	SO4	B	800	-	4,4,4	0.19	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	OXY	A	620	8	-	0/0/0/0	0/0/0/0
2	NAG	A	700	1	-	0/6/23/26	0/1/1/1
2	NAG	A	750	1	-	0/6/23/26	0/1/1/1
2	NAG	A	760	1	-	0/6/23/26	0/1/1/1
6	NDG	A	770	1	-	0/6/23/26	0/1/1/1
10	SO4	A	800	-	-	0/0/0/0	0/0/0/0
10	SO4	A	801	-	-	0/0/0/0	0/0/0/0
11	OXY	B	620	8	-	0/0/0/0	0/0/0/0
2	NAG	B	700	1	-	0/6/23/26	0/1/1/1
2	NAG	B	750	1	-	0/6/23/26	0/1/1/1
2	NAG	B	760	1	-	0/6/23/26	0/1/1/1
10	SO4	B	800	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	770	NDG	C1-C2	3.21	1.56	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	750	NAG	C2-N2-C7	-2.49	119.83	123.04
6	A	770	NDG	C3-C4-C5	-2.42	105.98	110.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	770	NDG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	559/559 (100%)	-0.36	6 (1%) 82 83	7, 15, 27, 47	0
1	B	559/559 (100%)	-0.24	7 (1%) 79 80	8, 17, 31, 50	0
All	All	1118/1118 (100%)	-0.30	13 (1%) 81 81	7, 16, 29, 50	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	462	ALA	4.4
1	A	462	ALA	3.6
1	B	1	GLU	3.5
1	A	1	GLU	3.3
1	A	3	THR	3.3
1	B	314	GLY	3.2
1	A	463	VAL	3.2
1	B	463	VAL	2.8
1	A	368	THR	2.6
1	B	3	THR	2.5
1	A	2	PRO	2.4
1	B	354	ARG	2.2
1	B	368	THR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	B	731	14/15	0.79	0.23	8.89	28,33,37,46	0
5	NAG	A	740	14/15	0.90	0.12	1.64	21,24,34,36	0
5	NAG	B	740	14/15	0.84	0.15	1.58	23,29,35,39	0
4	NAG	A	721	14/15	0.86	0.18	1.49	30,39,44,51	0
4	NAG	A	731	14/15	0.89	0.14	1.46	19,23,28,36	0
4	NAG	A	720	14/15	0.89	0.14	0.92	26,29,34,36	0
5	NAG	B	720	14/15	0.82	0.18	0.91	28,31,36,36	0
5	NAG	B	721	14/15	0.84	0.20	0.76	34,38,40,42	0
3	NAG	A	711	14/15	0.92	0.11	0.22	15,23,30,37	0
7	NAG	B	711	14/15	0.90	0.12	-0.06	22,28,34,41	0
4	NAG	A	730	14/15	0.94	0.10	-0.44	14,16,17,19	0
7	NAG	B	710	14/15	0.94	0.10	-0.48	13,16,18,23	0
3	NAG	A	710	14/15	0.97	0.09	-0.65	10,13,17,18	0
4	NAG	B	730	14/15	0.95	0.09	-0.82	17,18,21,24	0
5	NAG	B	741	14/15	0.79	0.33	-	44,48,52,52	0
3	MAN	A	714	11/12	0.64	0.26	-	64,65,66,68	0
3	MAN	A	713	11/12	0.72	0.53	-	66,69,70,70	0
3	BMA	A	712	11/12	0.61	0.32	-	47,56,61,63	0
7	MAN	B	714	11/12	0.53	0.29	-	66,67,68,69	0
4	BMA	A	732	11/12	0.77	0.35	-	41,46,47,48	0
5	NAG	A	741	14/15	0.81	0.29	-	40,45,48,49	0
4	BMA	B	732	11/12	0.70	0.42	-	51,56,58,58	0
4	BMA	A	722	11/12	0.70	0.41	-	57,61,62,62	0
7	BMA	B	712	11/12	0.64	0.29	-	48,53,59,63	0

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	760	14/15	0.74	0.22	5.63	32,35,38,40	0
6	NDG	A	770	14/15	0.70	0.28	5.23	43,46,48,49	0
2	NAG	A	760	14/15	0.76	0.22	5.12	30,32,33,36	0
2	NAG	A	700	14/15	0.84	0.17	3.78	26,28,31,33	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
10	SO4	A	801	5/5	0.84	0.18	3.43	74,74,74,74	0
2	NAG	B	700	14/15	0.87	0.17	2.91	30,32,34,35	0
2	NAG	B	750	14/15	0.87	0.16	2.90	22,24,26,26	0
11	OXY	A	620	2/2	0.98	0.10	1.96	32,32,32,33	0
2	NAG	A	750	14/15	0.85	0.21	1.93	34,38,43,44	0
11	OXY	B	620	2/2	0.98	0.10	1.02	27,27,27,28	0
9	CL	A	610	1/1	0.99	0.07	-0.91	22,22,22,22	0
9	CL	B	610	1/1	0.99	0.07	-1.71	27,27,27,27	0
8	CU	A	601	1/1	0.99	0.04	-2.59	15,15,15,15	0
8	CU	B	602	1/1	0.97	0.04	-2.98	22,22,22,22	0
8	CU	A	604	1/1	1.00	0.03	-3.28	15,15,15,15	0
8	CU	B	601	1/1	0.99	0.03	-3.38	15,15,15,15	0
8	CU	A	603	1/1	0.99	0.03	-4.17	15,15,15,15	0
8	CU	B	603	1/1	0.99	0.04	-5.27	17,17,17,17	0
8	CU	A	602	1/1	0.99	0.04	-5.97	18,18,18,18	0
8	CU	B	604	1/1	1.00	0.02	-7.45	20,20,20,20	0
10	SO4	A	800	5/5	0.88	0.14	-	62,62,63,63	0
10	SO4	B	800	5/5	0.92	0.09	-	71,71,71,71	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.