



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:48 AM GMT

PDB ID : 2IRF
Title : CRYSTAL STRUCTURE OF AN IRF-2/DNA COMPLEX.
Authors : Fujii, Y.; Shimizu, T.; Kusumoto, M.; Kyogoku, Y.; Taniguchi, T.; Hakoshima, T.
Deposited on : 1999-05-30
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	12	
1	B	12	
1	C	12	
2	D	13	
2	E	13	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	13	<div><div></div><div>62%38%</div></div>
3	G	113	<div><div></div><div>85%11%••</div></div>
3	H	113	<div><div></div><div>7%79%16%••</div></div>
3	I	113	<div><div></div><div>2%83%12%••</div></div>
3	J	113	<div><div></div><div>6%77%17%••</div></div>
3	K	113	<div><div></div><div>2%86%9%••</div></div>
3	L	113	<div><div></div><div>7%86%8%••</div></div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7397 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(P*AP*AP*GP*TP*GP*AP*AP*AP*GP*(5IU)P*GP*A)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	12	Total	C	I	N	O	P	0	0	0
			254	119	1	54	68	12			
1	B	12	Total	C	I	N	O	P	0	0	0
			254	119	1	54	68	12			
1	C	12	Total	C	I	N	O	P	0	0	0
			254	119	1	54	68	12			

- Molecule 2 is a DNA chain called DNA (5'-D(*TP*TP*CP*AP*CP*TP*TP*TP*CP*AP*CP*(5IU)P*T)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	D	13	Total	C	I	N	O	P	0	0	0
			255	125	1	36	81	12			
2	E	12	Total	C	I	N	O	P	0	0	0
			238	115	1	34	76	12			
2	F	13	Total	C	I	N	O	P	0	0	0
			255	125	1	36	81	12			

- Molecule 3 is a protein called INTERFERON REGULATORY FACTOR 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	109	Total	C	N	O	S	0	0	0
			911	583	170	152	6			
3	H	109	Total	C	N	O	S	0	0	0
			917	586	173	152	6			
3	I	109	Total	C	N	O	S	0	0	0
			911	583	170	152	6			
3	J	109	Total	C	N	O	S	0	0	0
			917	586	173	152	6			
3	K	109	Total	C	N	O	S	0	0	0
			917	586	173	152	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	109	Total	C	N	O	S	0	0	0
			911	583	170	152	6			

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	1	Total	K	0	0
			1	1		
4	J	1	Total	K	0	0
			1	1		
4	K	1	Total	K	0	0
			1	1		
4	H	1	Total	K	0	0
			1	1		
4	I	1	Total	K	0	0
			1	1		
4	L	1	Total	K	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	21	Total	O	0	0
			21	21		
5	B	20	Total	O	0	0
			20	20		
5	C	15	Total	O	0	0
			15	15		
5	D	26	Total	O	0	0
			26	26		
5	E	24	Total	O	0	0
			24	24		
5	F	29	Total	O	0	0
			29	29		
5	G	63	Total	O	0	0
			63	63		
5	H	47	Total	O	0	0
			47	47		
5	I	47	Total	O	0	0
			47	47		
5	J	14	Total	O	0	0
			14	14		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	K	55	Total	O	0	0
			55	55		
5	L	36	Total	O	0	0
			36	36		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(P*AP*AP*GP*TP*GP*AP*AP*AP*GP*(5IU)P*GP*A)-3')

Chain A: 




- Molecule 1: DNA (5'-D(P*AP*AP*GP*TP*GP*AP*AP*AP*GP*(5IU)P*GP*A)-3')

Chain B: 



- Molecule 1: DNA (5'-D(P*AP*AP*GP*TP*GP*AP*AP*AP*GP*(5IU)P*GP*A)-3')

Chain C: 



- Molecule 2: DNA (5'-D(*TP*TP*CP*AP*CP*TP*TP*TP*CP*AP*CP*(5IU)P*T)-3')

Chain D: 




- Molecule 2: DNA (5'-D(*TP*TP*CP*AP*CP*TP*TP*TP*CP*AP*CP*(5IU)P*T)-3')

Chain E: 



- Molecule 2: DNA (5'-D(*TP*TP*CP*AP*CP*TP*TP*TP*CP*AP*CP*(5IU)P*T)-3')

Chain F: 



• Molecule 3: INTERFERON REGULATORY FACTOR 2

Chain G: 85% 11% . .



• Molecule 3: INTERFERON REGULATORY FACTOR 2

Chain H: 7% 79% 16% . .




• Molecule 3: INTERFERON REGULATORY FACTOR 2

Chain I: 2% 83% 12% . .




• Molecule 3: INTERFERON REGULATORY FACTOR 2

Chain J: 6% 77% 17% . .



• Molecule 3: INTERFERON REGULATORY FACTOR 2

Chain K: 2% 86% 9% . .



• Molecule 3: INTERFERON REGULATORY FACTOR 2

Chain L: 7% 86% 8% . .



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	132.30 Å 132.30 Å 296.50 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 2.20 56.25 – 2.20	Depositor EDS
% Data completeness (in resolution range)	88.4 (10.00-2.20) 90.5 (56.25-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.89 (at 2.20 Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.202 , 0.243 0.203 , 0.242	Depositor DCC
R_{free} test set	3552 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.200	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 40.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	4 of 70905 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7397	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, 5IU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/264	0.73	0/404
1	B	0.34	0/264	0.74	0/404
1	C	0.38	0/264	0.71	0/404
2	D	0.36	0/259	0.74	0/394
2	E	0.38	0/240	0.74	0/364
2	F	0.34	0/259	0.72	0/394
3	G	0.30	0/940	0.55	0/1268
3	H	0.30	0/946	0.55	0/1275
3	I	0.31	0/940	0.54	0/1268
3	J	0.31	0/946	0.53	0/1275
3	K	0.31	0/946	0.57	0/1275
3	L	0.30	0/940	0.55	0/1268
All	All	0.32	0/7208	0.60	0/9993

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1
2	E	0	1
2	F	0	1
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	1105	DT	Sidechain
2	E	1117	DT	Sidechain
2	F	1129	DT	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	254	0	132	9	0
1	B	254	0	132	10	0
1	C	254	0	132	5	0
2	D	255	0	149	4	0
2	E	238	0	136	3	0
2	F	255	0	149	3	0
3	G	911	0	906	6	0
3	H	917	0	917	13	0
3	I	911	0	906	8	0
3	J	917	0	917	13	0
3	K	917	0	917	9	0
3	L	911	0	906	10	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
4	I	1	0	0	0	0
4	J	1	0	0	0	0
4	K	1	0	0	0	0
4	L	1	0	0	0	0
5	A	21	0	0	0	0
5	B	20	0	0	0	0
5	C	15	0	0	0	0
5	D	26	0	0	0	0
5	E	24	0	0	0	0
5	F	29	0	0	1	0
5	G	63	0	0	1	0
5	H	47	0	0	2	0
5	I	47	0	0	1	0
5	J	14	0	0	0	0
5	K	55	0	0	1	0
5	L	36	0	0	0	0
All	All	7397	0	6299	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:2100:LYS:HA	3:L:2105:ALA:HA	1.42	1.00
2:F:1124:DT:H4'	3:J:701:LYS:O	1.81	0.81
1:A:1001:DA:H2''	1:A:1002:DA:C8	2.24	0.71
1:B:1013:DA:H2''	1:B:1014:DA:C8	2.27	0.69
3:G:39:MET:HE2	3:G:104:ASN:H	1.57	0.68
3:K:888:LEU:HD23	3:K:891:ILE:HD12	1.77	0.67
3:L:2088:LEU:HD23	3:L:2091:ILE:HD12	1.77	0.66
1:A:1012:DA:O3'	1:B:1013:DA:H5'	1.96	0.65
2:D:1103:DA:H5''	3:L:2007:ARG:CZ	2.27	0.65
3:H:238:TRP:CD2	3:H:307:ARG:HD3	2.32	0.63
3:J:698:SER:HB3	3:J:706:PHE:O	2.00	0.61
1:B:1024:DA:O3'	1:C:1025:DA:P	2.60	0.59
1:C:1025:DA:H2''	1:C:1026:DA:C8	2.39	0.58
1:B:1022:5IU:H2'	3:K:882:ARG:CZ	2.36	0.55
1:C:1026:DA:C8	3:L:2075:LYS:HE2	2.40	0.55
3:J:699:ILE:O	3:J:705:ALA:HA	2.07	0.55
3:J:647:ASP:HB3	3:J:650:LYS:HB3	1.91	0.53
3:G:52:ALA:N	3:G:53:PRO:HD2	2.24	0.53
2:F:1134:DC:H2'	2:F:1135:5IU:I5	2.80	0.52
1:B:1016:DT:H2''	1:B:1017:DG:C8	2.44	0.52
1:A:1010:5IU:H2'	3:I:482:ARG:CZ	2.39	0.52
3:L:2052:ALA:N	3:L:2053:PRO:HD2	2.25	0.52
1:C:1028:DT:H2'	3:L:2082:ARG:CZ	2.40	0.51
1:B:1016:DT:H2'	3:J:682:ARG:CZ	2.41	0.51
3:J:652:ALA:N	3:J:653:PRO:CD	2.74	0.51
2:E:1124:DT:H2'	5:I:3159:HOH:O	2.11	0.50
3:K:852:ALA:N	3:K:853:PRO:HD2	2.26	0.50
1:A:1010:5IU:H2'	3:I:482:ARG:NH1	2.27	0.50
3:L:2052:ALA:N	3:L:2053:PRO:CD	2.75	0.50
3:H:300:LYS:HB3	5:H:3223:HOH:O	2.12	0.49
1:B:1016:DT:H2'	3:J:682:ARG:NH1	2.28	0.49
1:C:1028:DT:H2'	3:L:2082:ARG:NH1	2.29	0.48
3:G:52:ALA:N	3:G:53:PRO:CD	2.76	0.48
3:K:852:ALA:N	3:K:853:PRO:CD	2.76	0.48
3:J:612:LEU:O	3:J:616:ILE:HG13	2.14	0.47
1:A:1012:DA:O3'	1:B:1013:DA:P	2.73	0.46
3:L:2012:LEU:O	3:L:2016:ILE:HG13	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:452:ALA:N	3:I:453:PRO:CD	2.78	0.46
3:J:628:ASN:OD1	3:J:631:LYS:HB2	2.16	0.46
2:F:1136:DT:H2'	5:G:3077:HOH:O	2.16	0.46
1:B:1022:5IU:H2'	3:K:882:ARG:NH1	2.30	0.46
3:I:433:ILE:HG12	3:I:510:ARG:HG3	1.99	0.45
2:D:1112:DT:O3'	2:E:1113:DT:H5'	2.16	0.45
1:A:1004:DT:H2'	3:H:282:ARG:CZ	2.47	0.45
3:J:622:PRO:O	3:J:637:PRO:HG2	2.17	0.45
3:I:422:PRO:O	3:I:437:PRO:HG2	2.17	0.45
2:D:1110:DC:H2'	2:D:1111:5IU:I5	2.87	0.44
5:F:3147:HOH:O	3:H:287:SER:HB3	2.18	0.44
3:I:412:LEU:O	3:I:416:ILE:HG13	2.19	0.43
3:H:252:ALA:N	3:H:253:PRO:CD	2.82	0.43
3:J:652:ALA:N	3:J:653:PRO:HD2	2.34	0.43
3:H:307:ARG:HD2	5:H:3223:HOH:O	2.19	0.42
3:J:691:ILE:HG12	3:J:711:MET:HG2	2.00	0.42
3:K:832:LYS:HB3	3:K:911:MET:HB2	2.01	0.42
1:A:1001:DA:H2''	1:A:1002:DA:H8	1.80	0.42
1:A:1002:DA:H2'	3:H:275:LYS:HG3	2.01	0.42
3:I:441:ALA:HA	3:I:446:TRP:CG	2.54	0.42
3:K:833:ILE:HG12	3:K:910:ARG:HG3	2.02	0.42
2:D:1102:DC:H2''	2:D:1103:DA:C8	2.54	0.42
3:J:700:LYS:HA	3:J:705:ALA:HB1	2.02	0.42
3:H:266:GLN:HB2	3:H:269:ILE:HB	2.01	0.41
3:H:243:ARG:HH22	3:H:301:LYS:NZ	2.18	0.41
3:L:2104:ASN:H	3:L:2104:ASN:ND2	2.18	0.41
3:K:891:ILE:HG12	3:K:911:MET:HG2	2.02	0.41
3:K:815:GLN:HG2	5:K:3056:HOH:O	2.21	0.41
1:A:1002:DA:C8	3:H:275:LYS:HE3	2.56	0.41
3:H:284:ALA:O	3:H:288:LEU:HG	2.20	0.41
3:G:88:LEU:HD12	3:G:91:ILE:HD12	2.03	0.40
1:B:1022:5IU:H2''	1:B:1023:DG:C8	2.56	0.40
3:H:291:ILE:HG12	3:H:311:MET:HG2	2.03	0.40
2:E:1122:DC:H2'	2:E:1123:5IU:I5	2.91	0.40
3:G:21:ILE:HG21	3:G:54:LEU:HB2	2.03	0.40
3:H:212:LEU:O	3:H:216:ILE:HG13	2.21	0.40
3:I:472:PRO:C	3:I:474:PRO:HD3	2.42	0.40
3:G:100:LYS:HA	3:G:105:ALA:HB1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	G	107/113 (95%)	102 (95%)	4 (4%)	1 (1%)	21	19
3	H	107/113 (95%)	104 (97%)	1 (1%)	2 (2%)	10	6
3	I	107/113 (95%)	103 (96%)	2 (2%)	2 (2%)	10	6
3	J	107/113 (95%)	99 (92%)	4 (4%)	4 (4%)	4	1
3	K	107/113 (95%)	102 (95%)	4 (4%)	1 (1%)	21	19
3	L	107/113 (95%)	99 (92%)	7 (6%)	1 (1%)	21	19
All	All	642/678 (95%)	609 (95%)	22 (3%)	11 (2%)	11	7

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	J	703	ASN
3	J	696	ASP
3	H	252	ALA
3	I	452	ALA
3	J	652	ALA
3	H	299	ILE
3	G	52	ALA
3	J	699	ILE
3	K	852	ALA
3	L	2052	ALA
3	I	499	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	G	96/101 (95%)	93 (97%)	3 (3%)	47	59
3	H	97/101 (96%)	95 (98%)	2 (2%)	61	74
3	I	96/101 (95%)	95 (99%)	1 (1%)	82	91
3	J	97/101 (96%)	95 (98%)	2 (2%)	61	74
3	K	97/101 (96%)	94 (97%)	3 (3%)	47	59
3	L	96/101 (95%)	94 (98%)	2 (2%)	61	74
All	All	579/606 (96%)	566 (98%)	13 (2%)	60	72

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	7	ARG
3	G	9	ARG
3	G	12	LEU
3	H	212	LEU
3	H	247	ASP
3	I	409	ARG
3	J	607	ARG
3	J	612	LEU
3	K	812	LEU
3	K	888	LEU
3	K	894	VAL
3	L	2012	LEU
3	L	2104	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
3	G	15	GLN
3	G	57	ASN
3	H	215	GLN
3	H	257	ASN
3	I	444	HIS
3	J	665	HIS
3	K	815	GLN
3	K	857	ASN
3	L	2015	GLN
3	L	2057	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	5IU	A	1010	1,2	12,21,22	1.26	2 (16%)	14,30,33	3.68	1 (7%)
1	5IU	B	1022	1,2	12,21,22	1.20	2 (16%)	14,30,33	3.75	1 (7%)
1	5IU	C	1034	1,2	12,21,22	1.26	2 (16%)	14,30,33	3.72	1 (7%)
2	5IU	D	1111	1,2	12,21,22	1.36	2 (16%)	14,30,33	3.77	2 (14%)
2	5IU	E	1123	1,2	12,21,22	1.26	2 (16%)	14,30,33	3.79	2 (14%)
2	5IU	F	1135	1,2	12,21,22	1.19	2 (16%)	14,30,33	3.76	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	5IU	A	1010	1,2	-	0/3/21/22	0/2/2/2
1	5IU	B	1022	1,2	-	0/3/21/22	0/2/2/2
1	5IU	C	1034	1,2	-	0/3/21/22	0/2/2/2
2	5IU	D	1111	1,2	-	0/3/21/22	0/2/2/2
2	5IU	E	1123	1,2	-	0/3/21/22	0/2/2/2
2	5IU	F	1135	1,2	-	0/3/21/22	0/2/2/2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1010	5IU	C6-N1	2.26	1.38	1.35
1	C	1034	5IU	C6-N1	2.37	1.38	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1022	5IU	C6-N1	2.43	1.38	1.35
2	F	1135	5IU	C6-N1	2.61	1.38	1.35
2	F	1135	5IU	C4-N3	2.66	1.38	1.33
1	B	1022	5IU	C4-N3	2.72	1.38	1.33
2	E	1123	5IU	C4-N3	2.85	1.38	1.33
2	D	1111	5IU	C6-N1	2.86	1.39	1.35
2	E	1123	5IU	C6-N1	2.88	1.39	1.35
1	C	1034	5IU	C4-N3	2.92	1.38	1.33
2	D	1111	5IU	C4-N3	2.94	1.38	1.33
1	A	1010	5IU	C4-N3	3.25	1.39	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1135	5IU	C2'-C1'-N1	2.07	119.19	114.16
2	D	1111	5IU	C2'-C1'-N1	2.19	119.47	114.16
2	E	1123	5IU	C2'-C1'-N1	2.32	119.81	114.16
1	A	1010	5IU	C4-N3-C2	13.70	127.09	115.25
1	C	1034	5IU	C4-N3-C2	13.83	127.20	115.25
2	D	1111	5IU	C4-N3-C2	13.86	127.22	115.25
2	F	1135	5IU	C4-N3-C2	13.86	127.23	115.25
2	E	1123	5IU	C4-N3-C2	13.93	127.28	115.25
1	B	1022	5IU	C4-N3-C2	13.95	127.30	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1010	5IU	2	0
1	B	1022	5IU	3	0
2	D	1111	5IU	1	0
2	E	1123	5IU	1	0
2	F	1135	5IU	1	0

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	11/12 (91%)	-0.77	0 100 100	9, 12, 25, 26	0
1	B	11/12 (91%)	-0.68	0 100 100	11, 15, 24, 29	0
1	C	11/12 (91%)	-0.73	0 100 100	11, 14, 24, 28	0
2	D	12/13 (92%)	-0.61	0 100 100	12, 15, 21, 46	0
2	E	11/13 (84%)	-0.78	0 100 100	12, 15, 22, 23	0
2	F	12/13 (92%)	-0.76	0 100 100	10, 13, 23, 45	0
3	G	109/113 (96%)	-0.38	0 100 100	9, 20, 44, 52	0
3	H	109/113 (96%)	0.00	8 (7%) 18 17	9, 23, 75, 90	0
3	I	109/113 (96%)	-0.31	2 (1%) 71 70	11, 25, 44, 58	0
3	J	109/113 (96%)	0.23	7 (6%) 23 22	13, 35, 70, 80	0
3	K	109/113 (96%)	-0.38	2 (1%) 71 70	11, 24, 48, 63	0
3	L	109/113 (96%)	0.15	8 (7%) 18 17	11, 25, 80, 94	0
All	All	722/753 (95%)	-0.17	27 (3%) 45 44	9, 24, 55, 94	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	J	698	SER	9.6
3	L	2102	GLY	8.7
3	L	2101	LYS	8.6
3	J	704	ASN	8.5
3	L	2098	SER	8.2
3	H	303	ASN	7.0
3	H	302	GLY	7.0
3	H	304	ASN	6.5
3	L	2097	ARG	6.4
3	L	2103	ASN	6.2
3	L	2099	ILE	6.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	J	699	ILE	6.1
3	J	697	ARG	6.0
3	J	703	ASN	5.7
3	L	2104	ASN	5.4
3	H	298	SER	4.9
3	H	299	ILE	4.4
3	J	700	LYS	4.0
3	I	504	ASN	3.9
3	H	296	ASP	3.9
3	L	2100	LYS	3.8
3	J	696	ASP	3.4
3	H	313	PRO	2.9
3	H	297	ARG	2.9
3	K	904	ASN	2.7
3	I	513	PRO	2.2
3	K	830	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	5IU	E	1123	20/21	1.00	0.11	-	8,14,21,24	0
2	5IU	F	1135	20/21	0.99	0.11	-	5,11,18,20	0
1	5IU	A	1010	20/21	0.99	0.12	-	8,14,20,28	0
2	5IU	D	1111	20/21	0.99	0.12	-	10,17,21,22	0
1	5IU	B	1022	20/21	0.99	0.10	-	6,13,18,27	0
1	5IU	C	1034	20/21	0.99	0.11	-	4,12,21,27	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	K	K	4004	1/1	0.98	0.11	-0.07	27,27,27,27	0
4	K	I	4003	1/1	0.99	0.10	-0.19	28,28,28,28	0
4	K	L	4002	1/1	0.97	0.10	-0.29	28,28,28,28	0
4	K	J	4005	1/1	0.96	0.10	-0.49	35,35,35,35	0
4	K	G	4001	1/1	0.99	0.09	-0.63	25,25,25,25	0
4	K	H	4006	1/1	0.97	0.08	-2.38	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.