



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:48 AM GMT

PDB ID : 2ITW  
Title : Crystal structure of EGFR kinase domain in complex with AFN941  
Authors : Yun, C.-H.; Boggon, T.J.; Li, Y.; Woo, S.; Greulich, H.; Meyerson, M.; Eck, M.J.  
Deposited on : 2006-05-25  
Resolution : 2.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1945 (2.90-2.86)
Clashscore	102246	2202 (2.90-2.86)
Ramachandran outliers	100387	2149 (2.90-2.86)
Sidechain outliers	100360	2152 (2.90-2.86)
RSRZ outliers	91569	1950 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ITQ	A	2018	X	-	-	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2510 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EPIDERMAL GROWTH FACTOR RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	301	Total	C	N	O	S	0	1	1
			2403	1544	406	437	16			

- Molecule 2 is 1,2,3,4-TETRAHYDROGEN STAUROSPORINE (three-letter code: ITQ) (formula: C<sub>28</sub>H<sub>28</sub>N<sub>4</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			35	28	4	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	72	Total	O	0	0
			72	72		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: EPIDERMAL GROWTH FACTOR RECEPTOR



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	144.45Å 144.45Å 144.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.08 – 2.88 24.07 – 2.88	Depositor EDS
% Data completeness (in resolution range)	99.7 (24.08-2.88) 99.7 (24.07-2.88)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.91 (at 2.89Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.187 , 0.256 0.185 , 0.253	Depositor DCC
$R_{free}$ test set	544 reflections (4.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	56.7	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.1	EDS
Estimated twinning fraction	0.026 for -l,-k,-h	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 11456 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2510	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ITQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.84	3/2457 (0.1%)	0.84	2/3324 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	950	CYS	CB-SG	-5.71	1.72	1.81
1	A	872	GLU	CD-OE2	5.67	1.31	1.25
1	A	797	CYS	CB-SG	-5.37	1.73	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	831	ARG	NE-CZ-NH1	6.70	123.65	120.30
1	A	778	LEU	CA-CB-CG	-6.10	101.26	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2403	0	2447	86	0
2	A	35	0	27	8	0
3	A	72	0	0	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2510	0	2474	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:858:LEU:HD12	3:A:3035:HOH:O	1.43	1.16
1:A:882:ALA:HB1	1:A:884:GLU:OE2	1.47	1.10
1:A:756:ASN:H	1:A:756:ASN:HD22	1.11	0.98
1:A:747:LEU:O	1:A:748:ARG:HB3	1.74	0.88
1:A:747:LEU:HD12	1:A:785:THR:HA	1.55	0.87
1:A:719:GLY:HA3	2:A:2018:ITQ:C26	2.09	0.83
1:A:724:GLY:HA2	1:A:748:ARG:HA	1.65	0.78
1:A:918:ILE:HD12	1:A:926:ILE:HD11	1.68	0.75
1:A:719:GLY:HA3	2:A:2018:ITQ:H26A	1.70	0.73
2:A:2018:ITQ:H16	2:A:2018:ITQ:H26B	1.70	0.73
1:A:919:PRO:HD2	1:A:922:GLU:OE2	1.88	0.73
1:A:748:ARG:O	1:A:749:GLU:HB2	1.87	0.72
1:A:756:ASN:ND2	1:A:756:ASN:H	1.88	0.71
1:A:868:GLU:HB2	1:A:890:ILE:HG23	1.74	0.70
1:A:776[A]:ARG:CG	1:A:776[A]:ARG:HH11	2.04	0.70
1:A:717:VAL:O	1:A:717:VAL:HG13	1.92	0.70
1:A:756:ASN:N	1:A:756:ASN:HD22	1.89	0.69
1:A:697:GLU:HB3	3:A:3002:HOH:O	1.93	0.69
1:A:857:GLY:O	1:A:858:LEU:HG	1.93	0.68
1:A:824:GLY:HA3	1:A:853:ILE:HD12	1.76	0.68
2:A:2018:ITQ:C16	2:A:2018:ITQ:H27B	2.24	0.67
1:A:751:THR:HG23	1:A:756:ASN:HD21	1.59	0.65
1:A:829:GLU:HG3	1:A:893:HIS:CG	2.33	0.63
1:A:916:ASP:HB2	3:A:3052:HOH:O	1.99	0.62
1:A:932:ARG:NH1	1:A:951:TRP:O	2.32	0.62
1:A:831:ARG:HB3	1:A:831:ARG:NH1	2.15	0.62
1:A:776[A]:ARG:HG2	1:A:776[A]:ARG:HH11	1.62	0.62
1:A:776[A]:ARG:HG2	1:A:776[A]:ARG:NH1	2.19	0.58
1:A:983:GLY:O	1:A:984:ASP:C	2.40	0.58
1:A:882:ALA:CB	1:A:884:GLU:OE2	2.39	0.58
1:A:751:THR:HG23	1:A:756:ASN:ND2	2.19	0.57
1:A:734:GLU:HG2	3:A:3005:HOH:O	2.03	0.57
1:A:831:ARG:HH11	1:A:831:ARG:HB3	1.69	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:869:TYR:CE2	1:A:871:ALA:HB2	2.41	0.56
1:A:829:GLU:HG3	1:A:893:HIS:CD2	2.40	0.55
1:A:776[B]:ARG:HH11	1:A:776[B]:ARG:HG2	1.71	0.55
1:A:918:ILE:HD12	1:A:926:ILE:CD1	2.36	0.55
1:A:876:VAL:HG12	1:A:877:PRO:HD2	1.88	0.55
1:A:772:PRO:O	1:A:852:LYS:HE2	2.09	0.53
1:A:965:ILE:O	1:A:969:SER:HB3	2.08	0.53
1:A:717:VAL:O	1:A:717:VAL:CG1	2.57	0.53
1:A:747:LEU:HB2	1:A:785:THR:OG1	2.10	0.52
1:A:705:ARG:NH1	1:A:731:TRP:CH2	2.78	0.52
1:A:1007:MET:HG3	1:A:1008:ASP:H	1.74	0.51
1:A:747:LEU:HG	1:A:786:VAL:N	2.26	0.51
1:A:844:LEU:HG	1:A:854:THR:HG21	1.92	0.51
1:A:705:ARG:HH11	1:A:731:TRP:HH2	1.57	0.50
1:A:868:GLU:CB	1:A:890:ILE:HG23	2.42	0.49
1:A:854:THR:OG1	1:A:855:ASP:N	2.45	0.49
1:A:918:ILE:HG21	1:A:926:ILE:HD12	1.94	0.49
1:A:724:GLY:CA	1:A:748:ARG:HA	2.39	0.49
1:A:749:GLU:O	1:A:750:ALA:C	2.51	0.49
1:A:719:GLY:HA3	2:A:2018:ITQ:O4	2.13	0.49
1:A:776[B]:ARG:NH1	1:A:1014:ASP:OD1	2.46	0.48
1:A:887:LEU:HB2	1:A:888:HIS:ND1	2.27	0.48
1:A:723:PHE:CD1	1:A:724:GLY:N	2.81	0.48
1:A:747:LEU:O	1:A:748:ARG:CB	2.56	0.48
1:A:705:ARG:NH1	1:A:731:TRP:HH2	2.11	0.48
1:A:776[B]:ARG:CG	1:A:776[B]:ARG:HH11	2.27	0.47
1:A:834:VAL:HG12	1:A:836:ARG:HG3	1.96	0.47
1:A:700:ASN:C	1:A:700:ASN:HD22	2.17	0.47
1:A:748:ARG:O	1:A:749:GLU:CB	2.59	0.47
1:A:893:HIS:O	1:A:896:ASP:HB2	2.15	0.46
1:A:720:SER:HB2	1:A:725:THR:HG23	1.97	0.46
1:A:736:GLU:O	1:A:737:LYS:HB2	2.15	0.46
1:A:792:LEU:O	1:A:794:PRO:HD3	2.16	0.45
1:A:909:THR:OG1	1:A:912:SER:HB2	2.17	0.45
1:A:761:ASP:O	1:A:764:TYR:HB3	2.17	0.45
1:A:719:GLY:HA3	2:A:2018:ITQ:H26	1.95	0.45
1:A:971:MET:HG2	1:A:978:TYR:CG	2.52	0.44
1:A:828:LEU:HD22	1:A:833:LEU:HD23	2.00	0.44
1:A:869:TYR:HE2	1:A:871:ALA:HB2	1.83	0.44
1:A:830:ASP:C	1:A:832:ARG:H	2.21	0.44
1:A:747:LEU:CD1	1:A:785:THR:HA	2.37	0.43

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:960:LYS:O	1:A:963:GLU:N	2.45	0.43
1:A:783:THR:OG1	1:A:785:THR:O	2.22	0.43
2:A:2018:ITQ:C16	2:A:2018:ITQ:H26B	2.44	0.43
1:A:984:ASP:C	3:A:3065:HOH:O	2.57	0.43
1:A:717:VAL:O	1:A:718:LEU:C	2.57	0.43
1:A:700:ASN:C	1:A:700:ASN:ND2	2.72	0.42
1:A:833:LEU:HD13	1:A:861:LEU:HD12	2.01	0.42
1:A:716:LYS:HE2	1:A:716:LYS:HB3	1.89	0.42
1:A:887:LEU:C	1:A:888:HIS:ND1	2.73	0.42
2:A:2018:ITQ:H27B	2:A:2018:ITQ:H16	1.98	0.41
1:A:904:VAL:O	1:A:908:MET:HG2	2.19	0.41
1:A:736:GLU:OE1	1:A:1016:TYR:OH	2.33	0.41
1:A:716:LYS:HB3	1:A:717:VAL:H	1.65	0.41
1:A:756:ASN:HA	1:A:759:ILE:HG22	2.02	0.40
1:A:748:ARG:HD2	3:A:3006:HOH:O	2.22	0.40
1:A:756:ASN:ND2	1:A:756:ASN:N	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	298/327 (91%)	255 (86%)	29 (10%)	14 (5%)	<b>3</b> <b>10</b>

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	723	PHE
1	A	749	GLU
1	A	748	ARG
1	A	806	LYS
1	A	718	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	720	SER
1	A	722	ALA
1	A	734	GLU
1	A	747	LEU
1	A	750	ALA
1	A	865	GLU
1	A	753	PRO
1	A	855	ASP
1	A	863	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	263/287 (92%)	234 (89%)	29 (11%)	8 21

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	697	GLU
1	A	700	ASN
1	A	705	ARG
1	A	717	VAL
1	A	748	ARG
1	A	756	ASN
1	A	757	LYS
1	A	768	SER
1	A	776[A]	ARG
1	A	776[B]	ARG
1	A	784	SER
1	A	797	CYS
1	A	806	LYS
1	A	831	ARG
1	A	832	ARG
1	A	841	ARG
1	A	846	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	861	LEU
1	A	881	MET
1	A	884	GLU
1	A	892	THR
1	A	899	SER
1	A	921	SER
1	A	925	SER
1	A	928	GLU
1	A	960	LYS
1	A	984	ASP
1	A	1007	MET
1	A	1008	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	700	ASN
1	A	756	ASN
1	A	808	ASN
1	A	893	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ITQ	A	2018	-	30,42,42	2.61	9 (30%)	27,68,68	2.02	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ITQ	A	2018	-	1/1/5/7	0/4/49/49	0/0/8/8

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2018	ITQ	C11-C18	-2.53	1.39	1.42
2	A	2018	ITQ	C10-C11	-2.41	1.40	1.43
2	A	2018	ITQ	C8-N1	-2.04	1.33	1.37
2	A	2018	ITQ	C1-C20	2.02	1.55	1.49
2	A	2018	ITQ	C24-C23	2.72	1.58	1.53
2	A	2018	ITQ	C9-N1	3.76	1.45	1.33
2	A	2018	ITQ	O6-C22	5.61	1.53	1.42
2	A	2018	ITQ	C22-C23	6.11	1.58	1.52
2	A	2018	ITQ	C10-C9	8.12	1.51	1.41

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2018	ITQ	C10-C9-N1	-4.93	101.62	111.09
2	A	2018	ITQ	C5-C6-C19	-4.09	103.76	110.28
2	A	2018	ITQ	C10-C7-C8	-3.87	104.42	107.98
2	A	2018	ITQ	C16-C17-C12	-2.88	116.58	120.73
2	A	2018	ITQ	C7-C8-N1	2.00	113.53	110.23
2	A	2018	ITQ	C5-C20-N3	2.26	108.17	106.67
2	A	2018	ITQ	O5-C8-C7	3.86	126.71	122.68

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	2018	ITQ	C25

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2018	ITQ	8	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	301/327 (92%)	-0.09	16 (5%) 30 24	30, 49, 89, 104	4 (1%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	750	ALA	4.1
1	A	754	LYS	3.4
1	A	922	GLU	3.3
1	A	696	GLY	3.1
1	A	753	PRO	3.0
1	A	722	ALA	2.9
1	A	875	LYS	2.8
1	A	918	ILE	2.8
1	A	749	GLU	2.7
1	A	865	GLU	2.5
1	A	867	LYS	2.5
1	A	919	PRO	2.3
1	A	864	ALA	2.2
1	A	748	ARG	2.1
1	A	747	LEU	2.1
1	A	874	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ITQ	A	2018	35/35	0.91	0.18	0.06	68,78,82,84	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.