



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:50 AM GMT

PDB ID : 2IXF  
Title : CRYSTAL STRUCTURE OF THE ATPASE DOMAIN OF TAP1 WITH ATP  
(D645Q, Q678H MUTANT)  
Authors : Procko, E.; Ferrin-O'Connell, I.; Ng, S.-L.; Gaudet, R.  
Deposited on : 2006-07-08  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	<div> <div>6%</div> <div>86% 8% 6%</div> </div>
1	B	271	<div> <div>6%</div> <div>87% 7% 6%</div> </div>
1	C	271	<div> <div>6%</div> <div>86% 7% 7%</div> </div>
1	D	271	<div> <div>4%</div> <div>87% 5% 8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	3	-	-	-	X
4	GOL	B	3	-	-	-	X
4	GOL	B	4	-	-	-	X
4	GOL	C	3	-	-	-	X
4	GOL	D	3	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8862 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANTIGEN PEPTIDE TRANSPORTER 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	255	Total	C	N	O	S	0	3	0
			1933	1219	347	359	8			
1	B	255	Total	C	N	O	S	0	4	0
			1942	1227	345	362	8			
1	C	252	Total	C	N	O	S	0	5	0
			1931	1221	342	361	7			
1	D	250	Total	C	N	O	S	0	6	0
			1930	1220	346	357	7			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	464	MET	-	EXPRESSION TAG	UNP P36370
A	726	ALA	-	EXPRESSION TAG	UNP P36370
A	727	ALA	-	EXPRESSION TAG	UNP P36370
A	728	ALA	-	EXPRESSION TAG	UNP P36370
A	729	HIS	-	EXPRESSION TAG	UNP P36370
A	730	HIS	-	EXPRESSION TAG	UNP P36370
A	731	HIS	-	EXPRESSION TAG	UNP P36370
A	732	HIS	-	EXPRESSION TAG	UNP P36370
A	733	HIS	-	EXPRESSION TAG	UNP P36370
A	734	HIS	-	EXPRESSION TAG	UNP P36370
A	645	GLN	ASP	ENGINEERED MUTATION	UNP P36370
A	678	HIS	GLN	ENGINEERED MUTATION	UNP P36370
B	464	MET	-	EXPRESSION TAG	UNP P36370
B	726	ALA	-	EXPRESSION TAG	UNP P36370
B	727	ALA	-	EXPRESSION TAG	UNP P36370
B	728	ALA	-	EXPRESSION TAG	UNP P36370
B	729	HIS	-	EXPRESSION TAG	UNP P36370
B	730	HIS	-	EXPRESSION TAG	UNP P36370
B	731	HIS	-	EXPRESSION TAG	UNP P36370
B	732	HIS	-	EXPRESSION TAG	UNP P36370
B	733	HIS	-	EXPRESSION TAG	UNP P36370

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	734	HIS	-	EXPRESSION TAG	UNP P36370
B	645	GLN	ASP	ENGINEERED MUTATION	UNP P36370
B	678	HIS	GLN	ENGINEERED MUTATION	UNP P36370
C	464	MET	-	EXPRESSION TAG	UNP P36370
C	726	ALA	-	EXPRESSION TAG	UNP P36370
C	727	ALA	-	EXPRESSION TAG	UNP P36370
C	728	ALA	-	EXPRESSION TAG	UNP P36370
C	729	HIS	-	EXPRESSION TAG	UNP P36370
C	730	HIS	-	EXPRESSION TAG	UNP P36370
C	731	HIS	-	EXPRESSION TAG	UNP P36370
C	732	HIS	-	EXPRESSION TAG	UNP P36370
C	733	HIS	-	EXPRESSION TAG	UNP P36370
C	734	HIS	-	EXPRESSION TAG	UNP P36370
C	645	GLN	ASP	ENGINEERED MUTATION	UNP P36370
C	678	HIS	GLN	ENGINEERED MUTATION	UNP P36370
D	464	MET	-	EXPRESSION TAG	UNP P36370
D	726	ALA	-	EXPRESSION TAG	UNP P36370
D	727	ALA	-	EXPRESSION TAG	UNP P36370
D	728	ALA	-	EXPRESSION TAG	UNP P36370
D	729	HIS	-	EXPRESSION TAG	UNP P36370
D	730	HIS	-	EXPRESSION TAG	UNP P36370
D	731	HIS	-	EXPRESSION TAG	UNP P36370
D	732	HIS	-	EXPRESSION TAG	UNP P36370
D	733	HIS	-	EXPRESSION TAG	UNP P36370
D	734	HIS	-	EXPRESSION TAG	UNP P36370
D	645	GLN	ASP	ENGINEERED MUTATION	UNP P36370
D	678	HIS	GLN	ENGINEERED MUTATION	UNP P36370

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	251	Total	O	0	0
			251	251		
5	B	262	Total	O	0	0
			262	262		
5	C	254	Total	O	0	0
			254	254		
5	D	201	Total	O	0	0
			201	201		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: ANTIGEN PEPTIDE TRANSPORTER 1




#### • Molecule 1: ANTIGEN PEPTIDE TRANSPORTER 1



#### • Molecule 1: ANTIGEN PEPTIDE TRANSPORTER 1



#### • Molecule 1: ANTIGEN PEPTIDE TRANSPORTER 1





ALA
ALA
ALA
HIS
HIS
HIS
HIS
HIS
HIS

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.27Å 108.92Å 123.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.49 – 2.00 36.83 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.8 (38.49-2.00) 98.8 (36.83-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.82 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.216 , 0.258 0.218 , 0.261	Depositor DCC
$R_{free}$ test set	3772 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.9	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 37.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 74878 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8862	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 73.62 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.7772e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/1981	0.52	1/2691 (0.0%)
1	B	0.36	0/1991	0.51	0/2706
1	C	0.35	0/1982	0.49	0/2693
1	D	0.33	0/1976	0.50	0/2681
All	All	0.35	0/7930	0.50	1/10771 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	465	SER	N-CA-C	5.88	126.88	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1933	0	1912	12	0
1	B	1942	0	1923	10	0
1	C	1931	0	1910	11	0
1	D	1930	0	1915	8	0
2	A	31	0	12	0	0
2	B	31	0	12	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	31	0	12	0	0
2	D	31	0	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	6	0	8	2	0
4	B	12	0	16	1	0
4	C	6	0	8	1	0
4	D	6	0	8	1	0
5	A	251	0	0	1	0
5	B	262	0	0	1	0
5	C	254	0	0	2	0
5	D	201	0	0	0	0
All	All	8862	0	7748	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:476[B]:MET:HE3	1:B:541:LEU:HD11	1.59	0.83
1:C:698:CYS:SG	5:C:2250:HOH:O	2.45	0.74
1:A:583:THR:HG23	4:A:3:GOL:H31	1.72	0.71
1:B:698:CYS:SG	5:B:2236:HOH:O	2.27	0.70
1:A:698:CYS:SG	5:A:2250:HOH:O	2.54	0.66
1:D:524:VAL:O	1:D:528:LEU:HD13	1.97	0.64
1:C:591:ALA:HA	1:C:594:MET:HE3	1.82	0.60
1:D:468:SER:HB2	1:D:547[B]:VAL:HG12	1.84	0.58
1:A:580:LEU:HD11	1:A:636:ARG:HG3	1.87	0.57
1:C:716:MET:SD	5:C:2052:HOH:O	2.58	0.56
1:A:583:THR:HG23	4:A:3:GOL:C3	2.34	0.56
1:B:559:ALA:HB3	1:B:638:PRO:HG3	1.89	0.53
1:B:563:GLN:HE21	1:B:563:GLN:N	2.06	0.52
1:A:559:ALA:HB3	1:A:638:PRO:HG3	1.94	0.50
1:D:583:THR:HG23	4:D:3:GOL:O2	2.12	0.50
1:B:508:GLY:HA2	1:B:669:ALA:O	2.13	0.48
1:D:471:LEU:HD22	1:D:548:GLN:HB3	1.95	0.48
1:D:567:LEU:HB3	1:D:570:ARG:HD3	1.96	0.48
1:C:552:HIS:O	1:C:556:THR:HG23	2.16	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:638:PRO:O	1:C:671:ARG:HD3	2.15	0.46
1:D:644:ASP:HA	1:D:676:ILE:HB	1.98	0.46
1:B:583:THR:HG23	4:B:3:GOL:O2	2.16	0.45
1:C:626:GLN:CD	1:C:649:ALA:HB3	2.37	0.45
1:A:590:THR:HG23	1:A:599:HIS:CE1	2.51	0.45
1:B:480:VAL:HG11	1:B:528:LEU:HD21	1.99	0.44
1:A:626:GLN:CG	1:A:646:ALA:O	2.66	0.44
1:B:561:VAL:HG22	1:B:633:ALA:HB2	2.00	0.44
1:C:626:GLN:OE1	1:C:649:ALA:HB3	2.18	0.44
1:C:583:THR:HG23	4:C:3:GOL:O2	2.18	0.43
1:A:465:SER:N	1:A:466:PRO:HD3	2.33	0.43
1:C:559:ALA:HB3	1:C:638:PRO:HG3	1.99	0.43
1:B:655:GLN:NE2	1:B:682:LEU:HD11	2.33	0.43
1:A:480:VAL:HG11	1:A:528:LEU:HD21	2.00	0.43
1:D:468:SER:HB2	1:D:547[A]:VAL:HG22	2.00	0.43
1:C:561:VAL:HG22	1:C:633:ALA:HB2	2.01	0.42
1:B:534:PRO:HG3	1:B:539:VAL:HG23	2.02	0.41
1:C:644:ASP:HA	1:C:676:ILE:HB	2.03	0.41
1:A:644:ASP:HA	1:A:676:ILE:HB	2.03	0.41
1:D:654:ASN:OD1	1:D:657:ARG:NH1	2.54	0.40
1:A:580:LEU:CD1	1:A:636:ARG:HG3	2.49	0.40
1:A:580:LEU:HD23	1:A:584:PRO:HG3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/271 (94%)	254 (99%)	2 (1%)	0	100	100
1	B	257/271 (95%)	256 (100%)	1 (0%)	0	100	100
1	C	255/271 (94%)	254 (100%)	1 (0%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	252/271 (93%)	251 (100%)	1 (0%)	0	100	100
All	All	1020/1084 (94%)	1015 (100%)	5 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/219 (92%)	195 (97%)	6 (3%)	48	47
1	B	204/219 (93%)	200 (98%)	4 (2%)	63	65
1	C	202/219 (92%)	198 (98%)	4 (2%)	63	65
1	D	201/219 (92%)	196 (98%)	5 (2%)	55	55
All	All	808/876 (92%)	789 (98%)	19 (2%)	58	58

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	465	SER
1	A	471	LEU
1	A	475	ASN
1	A	531	LEU
1	A	610	ASP
1	A	718	GLU
1	B	563	GLN
1	B	607	GLN
1	B	610	ASP
1	B	619	GLN
1	C	490	PRO
1	C	566	LEU
1	C	570	ARG
1	C	660	ARG
1	D	468	SER
1	D	501	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	570	ARG
1	D	709[A]	ARG
1	D	709[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	492	HIS
1	A	551	HIS
1	A	607	GLN
1	B	563	GLN
1	B	607	GLN
1	B	655	GLN
1	B	679	GLN
1	D	619	GLN
1	D	659	GLN
1	D	700	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ATP	A	1	3	24,33,33	0.91	1 (4%)	31,52,52	1.93	5 (16%)
4	GOL	A	3	-	5,5,5	0.36	0	5,5,5	0.37	0
2	ATP	B	1	3	24,33,33	0.89	1 (4%)	31,52,52	1.90	3 (9%)
4	GOL	B	3	-	5,5,5	0.34	0	5,5,5	0.20	0
4	GOL	B	4	-	5,5,5	0.32	0	5,5,5	0.40	0
2	ATP	C	1	3	24,33,33	0.92	1 (4%)	31,52,52	1.91	2 (6%)
4	GOL	C	3	-	5,5,5	0.36	0	5,5,5	0.34	0
2	ATP	D	1	3	24,33,33	0.98	1 (4%)	31,52,52	1.86	2 (6%)
4	GOL	D	3	-	5,5,5	0.36	0	5,5,5	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	1	3	-	0/18/38/38	0/3/3/3
4	GOL	A	3	-	-	0/4/4/4	0/0/0/0
2	ATP	B	1	3	-	0/18/38/38	0/3/3/3
4	GOL	B	3	-	-	0/4/4/4	0/0/0/0
4	GOL	B	4	-	-	0/4/4/4	0/0/0/0
2	ATP	C	1	3	-	0/18/38/38	0/3/3/3
4	GOL	C	3	-	-	0/4/4/4	0/0/0/0
2	ATP	D	1	3	-	0/18/38/38	0/3/3/3
4	GOL	D	3	-	-	0/4/4/4	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	ATP	C5-C4	2.55	1.46	1.40
2	A	1	ATP	C5-C4	2.68	1.46	1.40
2	C	1	ATP	C5-C4	2.70	1.46	1.40
2	D	1	ATP	C5-C4	2.92	1.47	1.40

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	ATP	N3-C2-N1	-8.70	122.23	128.89

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	ATP	N3-C2-N1	-8.48	122.40	128.89
2	B	1	ATP	N3-C2-N1	-8.35	122.50	128.89
2	D	1	ATP	N3-C2-N1	-8.20	122.61	128.89
2	D	1	ATP	C4-C5-N7	-2.90	106.81	109.48
2	A	1	ATP	C4-C5-N7	-2.83	106.87	109.48
2	B	1	ATP	C4-C5-N7	-2.43	107.24	109.48
2	B	1	ATP	C1'-N9-C4	-2.22	123.59	126.94
2	A	1	ATP	C1'-N9-C4	-2.12	123.74	126.94
2	C	1	ATP	C4-C5-N7	-2.06	107.58	109.48
2	A	1	ATP	O2A-PA-O3A	2.01	114.19	105.09
2	A	1	ATP	C2-N1-C6	2.10	122.52	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	3	GOL	2	0
4	B	3	GOL	1	0
4	C	3	GOL	1	0
4	D	3	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	255/271 (94%)	0.43	17 (6%) 21 22	34, 38, 43, 46	0
1	B	255/271 (94%)	0.43	17 (6%) 21 22	34, 38, 44, 46	0
1	C	252/271 (92%)	0.45	16 (6%) 23 24	34, 38, 43, 48	0
1	D	250/271 (92%)	0.42	11 (4%) 38 39	34, 38, 43, 49	0
All	All	1012/1084 (93%)	0.43	61 (6%) 25 27	34, 38, 43, 49	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	720	LEU	6.5
1	B	619	GLN	5.6
1	A	561	VAL	4.8
1	D	568	PHE	4.6
1	A	643	LEU	4.5
1	B	561	VAL	4.3
1	A	719	ALA	4.2
1	A	465	SER	4.2
1	B	466	PRO	4.1
1	C	561	VAL	4.0
1	C	564	GLU	3.9
1	B	643	LEU	3.8
1	C	676	ILE	3.7
1	C	565	PRO	3.7
1	A	619	GLN	3.6
1	B	474	LEU	3.6
1	C	560	ALA	3.6
1	C	568	PHE	3.6
1	A	676	ILE	3.5
1	D	560	ALA	3.4
1	B	467	LEU	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	468	SER	3.2
1	B	676	ILE	3.2
1	B	465	SER	3.2
1	C	674	LEU	3.1
1	D	561	VAL	3.1
1	C	469	GLY	2.9
1	D	676	ILE	2.9
1	C	642	ILE	2.8
1	B	581	THR	2.8
1	B	551[A]	HIS	2.8
1	B	525	ALA	2.8
1	B	560	ALA	2.8
1	B	646	ALA	2.7
1	C	474	LEU	2.7
1	D	642	ILE	2.7
1	A	642	ILE	2.7
1	C	526	ALA	2.7
1	B	475	ASN	2.6
1	A	560	ALA	2.6
1	D	526	ALA	2.6
1	C	531	LEU	2.5
1	C	646	ALA	2.5
1	A	525	ALA	2.5
1	A	467	LEU	2.4
1	B	674	LEU	2.4
1	A	466	PRO	2.4
1	D	698	CYS	2.4
1	D	704	LEU	2.3
1	A	646	ALA	2.3
1	A	675	LEU	2.2
1	D	674	LEU	2.2
1	C	562	GLY	2.1
1	A	677	THR	2.1
1	B	642	ILE	2.1
1	B	675	LEU	2.1
1	A	641	LEU	2.1
1	D	531	LEU	2.1
1	A	475	ASN	2.0
1	C	559	ALA	2.0
1	A	710	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	C	3	6/6	0.69	0.50	12.38	72,72,73,73	0
4	GOL	D	3	6/6	0.53	0.37	12.15	92,92,92,92	0
4	GOL	B	3	6/6	0.48	0.36	6.03	64,64,65,65	0
4	GOL	B	4	6/6	0.65	0.30	2.83	57,57,58,58	0
4	GOL	A	3	6/6	0.73	0.27	2.64	59,60,60,61	0
2	ATP	A	1	31/31	0.97	0.08	-1.19	19,25,29,29	0
2	ATP	B	1	31/31	0.97	0.08	-1.33	21,26,28,29	0
2	ATP	C	1	31/31	0.97	0.07	-1.37	21,28,30,31	0
2	ATP	D	1	31/31	0.97	0.07	-1.50	25,30,32,33	0
3	MG	A	2	1/1	0.95	0.14	-	27,27,27,27	0
3	MG	C	2	1/1	0.94	0.13	-	34,34,34,34	0
3	MG	B	2	1/1	0.96	0.23	-	30,30,30,30	0
3	MG	D	2	1/1	0.96	0.11	-	27,27,27,27	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.