



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 26, 2016 – 09:31 PM BST

PDB ID : 2J10
Title : P53 TETRAMERIZATION DOMAIN MUTANT T329F Q331K
Authors : Carbajo, R.J.; Mora, P.; Sanchez Del Pino, M.M.; Perez-Paya, E.; Pineda-Lucena, A.
Deposited on : 2006-08-08

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	31	 77% 19% •
1	B	31	 77% 23%
1	C	31	 81% 19%
1	D	31	 77% 19% •

2 Ensemble composition and analysis

This entry contains 30 models. Model 20 is the overall representative, medoid model (most similar to other models). The authors have identified model 5 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:327-A:356, B:326-B:356, C:326-C:356, D:326-D:356 (123)	0.58	20

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 7 clusters and 5 single-model clusters were found.

Cluster number	Models
1	1, 4, 5, 6, 9, 10, 25
2	12, 13, 24, 27, 28
3	7, 15, 17, 20, 29
4	21, 30
5	19, 23
6	11, 26
7	3, 22
Single-model clusters	2; 8; 14; 16; 18

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2132 atoms, of which 1056 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called CELLULAR TUMOR ANTIGEN P53.

Mol	Chain	Residues	Atoms						Trace
1	A	31	Total	C	H	N	O	S	0
			533	172	264	47	49	1	
1	B	31	Total	C	H	N	O	S	0
			533	172	264	47	49	1	
1	C	31	Total	C	H	N	O	S	0
			533	172	264	47	49	1	
1	D	31	Total	C	H	N	O	S	0
			533	172	264	47	49	1	

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	329	PHE	THR	ENGINEERED MUTATION	UNP P04637
A	331	LYS	GLN	ENGINEERED MUTATION	UNP P04637
B	329	PHE	THR	ENGINEERED MUTATION	UNP P04637
B	331	LYS	GLN	ENGINEERED MUTATION	UNP P04637
C	329	PHE	THR	ENGINEERED MUTATION	UNP P04637
C	331	LYS	GLN	ENGINEERED MUTATION	UNP P04637
D	329	PHE	THR	ENGINEERED MUTATION	UNP P04637
D	331	LYS	GLN	ENGINEERED MUTATION	UNP P04637

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: CELLULAR TUMOR ANTIGEN P53




- Molecule 1: CELLULAR TUMOR ANTIGEN P53



- Molecule 1: CELLULAR TUMOR ANTIGEN P53



- Molecule 1: CELLULAR TUMOR ANTIGEN P53




4.2 Residue scores for the representative (medoid) model from the NMR ensemble


The representative model is number 20. Colouring as in section 4.1 above.

- Molecule 1: CELLULAR TUMOR ANTIGEN P53






- Molecule 1: CELLULAR TUMOR ANTIGEN P53



- Molecule 1: CELLULAR TUMOR ANTIGEN P53



- Molecule 1: CELLULAR TUMOR ANTIGEN P53



5 Refinement protocol and experimental data overview

The models were refined using the following method: *CNS*.

Of the 30 calculated structures, 30 were deposited, based on the following criterion: *TOTAL ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
SPARKY	structure solution	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	260	256	256	9±3
1	B	269	264	261	9±2
1	C	269	264	261	10±3
1	D	269	264	261	10±3
All	All	32010	31440	31170	709

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

5 of 306 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:D:330:LEU:HD23	1:D:332:ILE:HD11	0.94	1.35	29	6
1:B:344:LEU:HD22	1:C:344:LEU:HD11	0.91	1.41	7	2
1:B:344:LEU:HD22	1:C:344:LEU:HD22	0.89	1.44	28	3
1:A:344:LEU:HD22	1:D:344:LEU:HD21	0.85	1.44	16	1
1:A:344:LEU:HD11	1:D:344:LEU:HD21	0.85	1.48	5	5

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	29/31 (94%)	27±1 (94±4%)	1±1 (5±4%)	0±0 (1±2%)	19	64
1	B	29/31 (94%)	27±1 (94±3%)	1±1 (4±3%)	1±0 (2±2%)	13	53
1	C	29/31 (94%)	27±1 (94±5%)	1±1 (4±4%)	0±0 (2±2%)	17	61
1	D	29/31 (94%)	27±1 (94±4%)	1±1 (5±3%)	0±0 (2±2%)	17	61
All	All	3480/3720 (94%)	3265 (94%)	157 (5%)	58 (2%)	16	59

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	327	TYR	18
1	D	327	TYR	14
1	C	327	TYR	13
1	A	327	TYR	12
1	C	355	ALA	1

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	25/26 (96%)	21±2 (82±7%)	4±2 (18±7%)	6	41
1	B	26/26 (100%)	22±2 (84±7%)	4±2 (16±7%)	7	44
1	C	26/26 (100%)	21±2 (81±7%)	5±2 (19±7%)	5	38
1	D	26/26 (100%)	21±1 (82±6%)	5±1 (18±6%)	6	41
All	All	3090/3120 (99%)	2547 (82%)	543 (18%)	6	41

5 of 85 unique residues with a non-rotameric sidechain are listed below. They are sorted by the

frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	D	344	LEU	16
1	B	342	ARG	16
1	A	344	LEU	15
1	A	335	ARG	14
1	A	342	ARG	14

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided