



wwPDB EM Map/Model Validation Report ⓘ

Apr 10, 2016 – 01:44 PM BST

PDB ID : 3J2Y
EMDB ID: : EMD-5578
Title : Electron Cryo-microscopy of Chikungunya VLP in complex with neutralizing antibody Fab 9.8B
Authors : Sun, S.; Xiang, Y.; Rossmann, M.G.
Deposited on : 2013-01-28
Resolution : 14.90 Å(reported)
Based on PDB ID : 4GQ9

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.
For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/EMValidationReportHelp>

MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : trunk27241


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 14.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|-----------------------------|-----------------------------|
| Clashscore | 114402 | 924 |
| Ramachandran outliers | 111179 | 726 |
| Sidechain outliers | 111093 | 686 |

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 212 | 83% 16% . |
| 1 | C | 212 | 83% 15% . |
| 1 | E | 212 | 85% 13% . |
| 1 | G | 212 | 85% 14% . |
| 2 | B | 218 | 78% 19% . |
| 2 | D | 218 | 79% 18% . |
| 2 | F | 218 | 79% 17% . |
| 2 | H | 218 | 77% 20% . |

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 13120 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 9.8B light chain.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 1 | A | 212 | Total | C | N | O | S | 0 | 0 |
| | | | 1633 | 1015 | 273 | 336 | 9 | | |
| 1 | C | 212 | Total | C | N | O | S | 0 | 0 |
| | | | 1633 | 1015 | 273 | 336 | 9 | | |
| 1 | E | 212 | Total | C | N | O | S | 0 | 0 |
| | | | 1633 | 1015 | 273 | 336 | 9 | | |
| 1 | G | 212 | Total | C | N | O | S | 0 | 0 |
| | | | 1633 | 1015 | 273 | 336 | 9 | | |


- Molecule 2 is a protein called 9.8B heavy chain.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 2 | B | 218 | Total | C | N | O | S | 0 | 0 |
| | | | 1647 | 1039 | 275 | 326 | 7 | | |
| 2 | D | 218 | Total | C | N | O | S | 0 | 0 |
| | | | 1647 | 1039 | 275 | 326 | 7 | | |
| 2 | F | 218 | Total | C | N | O | S | 0 | 0 |
| | | | 1647 | 1039 | 275 | 326 | 7 | | |
| 2 | H | 218 | Total | C | N | O | S | 0 | 0 |
| | | | 1647 | 1039 | 275 | 326 | 7 | | |


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 9.8B light chain




- Molecule 1: 9.8B light chain




- Molecule 1: 9.8B light chain




- Molecule 1: 9.8B light chain




- Molecule 2: 9.8B heavy chain






- Molecule 2: 9.8B heavy chain

Chain D: 79% 18% •




- Molecule 2: 9.8B heavy chain

Chain F: 79% 17% •



- Molecule 2: 9.8B heavy chain

Chain H: 77% 20% •



4 Experimental information

| Property | Value | Source |
|--------------------------------------|-------------------------|-----------|
| Reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, I | Depositor |
| Number of images | 1820 | Depositor |
| Resolution determination method | FSC at 0.5 cut-off | Depositor |
| CTF correction method | Each Micrograph | Depositor |
| Microscope | FEI/PHILIPS CM300FEG/HE | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | 20 | Depositor |
| Minimum defocus (nm) | Not provided | Depositor |
| Maximum defocus (nm) | Not provided | Depositor |
| Magnification | 38000 | Depositor |
| Image detector | Not provided | Depositor |

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------|-------------|---------|
| | | RMSZ | # Z >2 | RMSZ | # Z >2 |
| 1 | A | 0.34 | 0/1669 | 0.48 | 0/2268 |
| 1 | C | 0.34 | 0/1669 | 0.48 | 0/2268 |
| 1 | E | 0.33 | 0/1669 | 0.48 | 0/2268 |
| 1 | G | 0.33 | 0/1669 | 0.48 | 0/2268 |
| 2 | B | 0.33 | 0/1689 | 0.50 | 0/2308 |
| 2 | D | 0.33 | 0/1689 | 0.50 | 0/2308 |
| 2 | F | 0.33 | 0/1689 | 0.50 | 0/2308 |
| 2 | H | 0.33 | 0/1689 | 0.50 | 0/2308 |
| All | All | 0.33 | 0/13432 | 0.49 | 0/18304 |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 1633 | 0 | 1567 | 20 | 0 |
| 1 | C | 1633 | 0 | 1564 | 18 | 0 |
| 1 | E | 1633 | 0 | 1564 | 16 | 0 |
| 1 | G | 1633 | 0 | 1564 | 17 | 0 |
| 2 | B | 1647 | 0 | 1620 | 32 | 0 |
| 2 | D | 1647 | 0 | 1617 | 30 | 0 |
| 2 | F | 1647 | 0 | 1617 | 32 | 0 |
| 2 | H | 1647 | 0 | 1617 | 34 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| All | All | 13120 | 0 | 12730 | 199 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 199 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|------------------|--------------------------|-------------------|
| 1:G:3033:TYR:OH | 1:G:3090:ARG:NH1 | 1.58 | 1.36 |
| 1:E:2033:TYR:OH | 1:E:2090:ARG:NH1 | 1.58 | 1.36 |
| 1:C:1033:TYR:OH | 1:C:1090:ARG:NH1 | 1.58 | 1.34 |
| 1:A:33:TYR:OH | 1:A:90:ARG:NH1 | 1.58 | 1.31 |
| 2:D:1076:SER:HA | 2:D:1077:SER:HB3 | 1.16 | 1.14 |

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|----------|-------------|----|
| 1 | A | 210/212 (99%) | 195 (93%) | 12 (6%) | 3 (1%) | 14 | 58 |
| 1 | C | 210/212 (99%) | 195 (93%) | 12 (6%) | 3 (1%) | 14 | 58 |
| 1 | E | 210/212 (99%) | 195 (93%) | 12 (6%) | 3 (1%) | 14 | 58 |
| 1 | G | 210/212 (99%) | 195 (93%) | 12 (6%) | 3 (1%) | 14 | 58 |
| 2 | B | 216/218 (99%) | 188 (87%) | 22 (10%) | 6 (3%) | 6 | 44 |
| 2 | D | 216/218 (99%) | 188 (87%) | 22 (10%) | 6 (3%) | 6 | 44 |
| 2 | F | 216/218 (99%) | 188 (87%) | 22 (10%) | 6 (3%) | 6 | 44 |
| 2 | H | 216/218 (99%) | 188 (87%) | 22 (10%) | 6 (3%) | 6 | 44 |
| All | All | 1704/1720 (99%) | 1532 (90%) | 136 (8%) | 36 (2%) | 13 | 50 |

5 of 36 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|------|------|
| 2 | B | 77 | SER |
| 2 | D | 1077 | SER |
| 2 | F | 2077 | SER |
| 2 | H | 3077 | SER |
| 1 | A | 93 | TYR |

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|------------------|------------|----------|-------------|----|
| 1 | A | 187/187 (100%) | 178 (95%) | 9 (5%) | 31 | 67 |
| 1 | C | 187/187 (100%) | 178 (95%) | 9 (5%) | 31 | 67 |
| 1 | E | 187/187 (100%) | 178 (95%) | 9 (5%) | 31 | 67 |
| 1 | G | 187/187 (100%) | 178 (95%) | 9 (5%) | 31 | 67 |
| 2 | B | 189/189 (100%) | 182 (96%) | 7 (4%) | 41 | 73 |
| 2 | D | 189/189 (100%) | 182 (96%) | 7 (4%) | 41 | 73 |
| 2 | F | 189/189 (100%) | 182 (96%) | 7 (4%) | 41 | 73 |
| 2 | H | 189/189 (100%) | 182 (96%) | 7 (4%) | 41 | 73 |
| All | All | 1504/1504 (100%) | 1440 (96%) | 64 (4%) | 40 | 70 |

5 of 64 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|------|------|
| 1 | G | 3077 | MET |
| 1 | G | 3109 | ASP |
| 2 | H | 3061 | ASN |
| 1 | G | 3089 | GLN |
| 1 | G | 3104 | GLU |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|------|------|
| 2 | B | 50 | ASN |
| 2 | D | 1039 | GLN |
| 2 | H | 3050 | ASN |
| 2 | B | 61 | ASN |
| 2 | D | 1035 | HIS |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.