



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:40 PM GMT

PDB ID : 4J4B  
Title : PylD in complex with L-lysine-Ne-D-ornithine and NADH  
Authors : Quitterer, F.; Beck, P.; Bacher, A.; Groll, M.  
Deposited on : 2013-02-06  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	<div> <div>21%</div> <div>88%</div> <div>11%</div> <div>.</div> </div>
1	B	259	<div> <div>4%</div> <div>84%</div> <div>14%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NA	B	303	-	-	-	X
4	MG	B	304	-	-	-	X
5	0TF	B	301	-	-	-	X

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 4239 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PylD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	259	Total	C	N	O	S	0	0	0
			1952	1244	314	383	11			
1	B	259	Total	C	N	O	S	0	0	0
			1952	1244	314	383	11			

- Molecule 2 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (three-letter code: NAI) (formula:  $C_{21}H_{29}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Na 1 1	0	0
3	A	1	Total Na 1 1	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Mg 1 1	0	0
4	A	1	Total Mg 1 1	0	0

- Molecule 5 is N 6 -D-ORNITHYL-L-LYSINE (three-letter code: OTF) (formula: C<sub>11</sub>H<sub>24</sub>N<sub>4</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 18 11 4 3	0	0
5	B	1	Total C N O 18 11 4 3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	46	Total O 46 46	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	161	Total	O	0	0
			161	161		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: PylD



#### • Molecule 1: PylD



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.38Å 142.44Å 167.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.90 10.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (10.00-1.90) 99.9 (10.00-1.90)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.47 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.205 , 0.240 0.205 , 0.240	Depositor DCC
$R_{free}$ test set	2320 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.8	Xtriage
Anisotropy	0.407	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.50 , 70.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 45696 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4239	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 0TF, MG, NAI, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.30	0/1988	0.52	0/2700
1	B	0.36	0/1988	0.58	0/2700
All	All	0.33	0/3976	0.55	0/5400

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1952	0	1946	14	0
1	B	1952	0	1946	25	0
2	A	44	0	27	1	0
2	B	44	0	27	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	18	0	23	2	0
5	B	18	0	23	3	0
6	A	46	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	161	0	0	2	1
All	All	4239	0	3992	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:204:THR:HG22	1:B:206:CYS:H	1.38	0.88
1:B:143:ASP:OD1	1:B:166:ARG:HD2	1.89	0.73
1:B:36:LYS:HG3	6:B:405:HOH:O	1.92	0.69
1:B:204:THR:HG21	1:B:209:THR:HG21	1.73	0.68
1:A:215:LEU:HD11	1:A:230:ILE:HD11	1.76	0.67
1:B:120:ASN:HD22	1:B:122:GLN:H	1.43	0.66
1:A:204:THR:HG22	1:A:206:CYS:H	1.59	0.65
1:B:41:THR:HG21	1:B:256:TYR:O	1.95	0.65
1:A:41:THR:HG21	1:A:256:TYR:O	1.97	0.65
1:B:202:GLU:OE1	1:B:204:THR:HB	1.97	0.64
5:B:301:0TF:H7	2:B:302:NAI:H42N	1.79	0.64
1:A:103:ALA:HB2	1:A:108:PHE:CD1	2.36	0.61
1:A:120:ASN:HD22	1:A:122:GLN:H	1.49	0.60
1:B:20:ASP:O	1:B:24:GLN:HG3	2.01	0.60
1:B:215:LEU:HD11	1:B:230:ILE:HD11	1.85	0.58
1:B:204:THR:CG2	1:B:206:CYS:H	2.15	0.56
1:A:112:ASN:HD21	1:A:114:LYS:HB3	1.71	0.56
2:A:901:NAI:H42N	5:A:904:0TF:H7	1.88	0.55
1:B:143:ASP:OD1	1:B:166:ARG:CD	2.54	0.54
1:B:112:ASN:HD22	1:B:114:LYS:H	1.56	0.53
1:B:144:VAL:HG11	1:B:160:LEU:HD13	1.90	0.53
1:B:112:ASN:HD22	1:B:112:ASN:C	2.13	0.51
1:B:112:ASN:ND2	1:B:114:LYS:H	2.08	0.50
1:B:120:ASN:ND2	1:B:122:GLN:H	2.07	0.50
1:B:103:ALA:HB2	1:B:108:PHE:CD1	2.48	0.49
1:B:112:ASN:HD21	1:B:114:LYS:HB2	1.78	0.48
1:A:154:PHE:HB3	1:A:155:PRO:HD3	1.96	0.48
1:A:47:LYS:HA	1:A:77:ASP:O	2.14	0.47
1:A:112:ASN:HD22	1:A:114:LYS:H	1.62	0.47
5:B:301:0TF:N17	5:B:301:0TF:N13	2.62	0.47
1:B:210:ILE:HG12	1:B:229:ALA:HB1	1.99	0.45
1:B:47:LYS:HA	1:B:77:ASP:O	2.17	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:245:GLU:HG2	1:A:246:PRO:HD2	1.99	0.44
1:A:120:ASN:ND2	1:A:122:GLN:H	2.16	0.43
5:B:301:OTF:C16	2:B:302:NAI:H42N	2.48	0.42
1:B:204:THR:HG22	1:B:206:CYS:N	2.20	0.42
1:B:178:GLU:HB3	6:B:509:HOH:O	2.20	0.41
5:A:904:OTF:N17	5:A:904:OTF:N13	2.68	0.41
1:B:143:ASP:OD1	1:B:166:ARG:CG	2.68	0.41
1:A:204:THR:HG21	1:A:209:THR:HG21	2.02	0.41
1:A:112:ASN:HD22	1:A:112:ASN:C	2.24	0.41
1:B:181:THR:HG22	1:B:186:ILE:HG13	2.03	0.41
1:A:181:THR:HG21	1:A:188:PRO:HG3	2.02	0.40
1:B:133:ALA:HB1	1:B:199:ILE:CG2	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:517:HOH:O	6:B:517:HOH:O[3_454]	1.97	0.23

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/259 (99%)	248 (96%)	8 (3%)	1 (0%)	39	27
1	B	257/259 (99%)	250 (97%)	7 (3%)	0	100	100
All	All	514/518 (99%)	498 (97%)	15 (3%)	1 (0%)	52	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	227	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/214 (100%)	201 (94%)	12 (6%)	26	14
1	B	213/214 (100%)	197 (92%)	16 (8%)	17	7
All	All	426/428 (100%)	398 (93%)	28 (7%)	21	10

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	LEU
1	A	41	THR
1	A	48	VAL
1	A	57	ASN
1	A	112	ASN
1	A	120	ASN
1	A	137	LEU
1	A	184	LEU
1	A	195	LYS
1	A	202	GLU
1	A	228	CYS
1	A	235	ARG
1	B	11	ASN
1	B	25	GLU
1	B	29	LEU
1	B	41	THR
1	B	57	ASN
1	B	112	ASN
1	B	120	ASN
1	B	137	LEU
1	B	140	ASP
1	B	145	LEU
1	B	166	ARG
1	B	184	LEU
1	B	195	LYS
1	B	230	ILE
1	B	235	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	242	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	62	ASN
1	A	112	ASN
1	A	120	ASN
1	A	162	GLN
1	B	11	ASN
1	B	15	GLN
1	B	62	ASN
1	B	112	ASN
1	B	120	ASN
1	B	162	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAI	A	901	4	38,48,48	1.06	3 (7%)	48,73,73	1.57	4 (8%)
5	0TF	A	904	-	14,17,17	0.36	0	13,20,20	0.48	0
5	0TF	B	301	-	14,17,17	0.35	0	13,20,20	0.54	0
2	NAI	B	302	4	38,48,48	1.09	3 (7%)	48,73,73	1.63	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAI	A	901	4	-	0/25/72/72	0/5/5/5
5	0TF	A	904	-	-	0/16/20/20	0/0/0/0
5	0TF	B	301	-	-	0/16/20/20	0/0/0/0
2	NAI	B	302	4	-	0/25/72/72	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	NAI	C2N-C3N	2.02	1.39	1.34
2	B	302	NAI	C2N-C3N	2.29	1.40	1.34
2	A	901	NAI	C5A-C4A	3.15	1.47	1.40
2	B	302	NAI	C5A-C4A	3.22	1.47	1.40
2	A	901	NAI	C6N-C5N	3.37	1.39	1.33
2	B	302	NAI	C6N-C5N	3.62	1.40	1.33

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	302	NAI	N3A-C2A-N1A	-7.83	122.90	128.89
2	A	901	NAI	N3A-C2A-N1A	-7.28	123.32	128.89
2	A	901	NAI	PN-O3-PA	-4.30	120.66	132.73
2	B	302	NAI	PN-O3-PA	-3.65	122.48	132.73
2	A	901	NAI	C4A-C5A-N7A	-3.12	106.61	109.48
2	B	302	NAI	C4A-C5A-N7A	-3.08	106.65	109.48
2	A	901	NAI	C1D-N1N-C2N	-2.54	116.49	120.91
2	B	302	NAI	C1D-N1N-C2N	-2.31	116.88	120.91
2	B	302	NAI	C1B-N9A-C4A	-2.08	123.81	126.94
2	B	302	NAI	C2D-C1D-N1N	2.07	118.93	113.34

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	NAI	1	0
5	A	904	0TF	2	0
5	B	301	0TF	3	0
2	B	302	NAI	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	259/259 (100%)	1.06	54 (20%) 1 1	29, 47, 73, 86	0
1	B	259/259 (100%)	-0.04	10 (3%) 43 47	9, 19, 44, 63	0
All	All	518/518 (100%)	0.51	64 (12%) 5 5	9, 35, 67, 86	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	192	ALA	8.4
1	A	226	ILE	5.5
1	A	14	MET	5.4
1	B	1	MET	5.2
1	A	193	ASN	5.0
1	A	179	ARG	4.9
1	A	140	ASP	4.5
1	A	24	GLN	4.2
1	A	1	MET	4.2
1	A	227	PRO	4.2
1	A	175	THR	4.1
1	A	166	ARG	4.1
1	B	192	ALA	3.8
1	A	40	GLY	3.7
1	A	213	ALA	3.6
1	A	164	ASP	3.4
1	A	18	LYS	3.4
1	A	23	VAL	3.3
1	A	37	ALA	3.2
1	B	179	ARG	3.2
1	A	178	GLU	3.2
1	A	176	LEU	3.2
1	A	45	SER	3.1
1	A	233	GLU	3.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	217	GLU	3.0
1	A	236	ASP	3.0
1	B	175	THR	3.0
1	A	235	ARG	3.0
1	B	140	ASP	2.9
1	A	218	ASN	2.9
1	A	10	ILE	2.9
1	B	178	GLU	2.8
1	A	114	LYS	2.7
1	A	232	GLU	2.6
1	A	138	LYS	2.6
1	A	142	LYS	2.6
1	A	57	ASN	2.6
1	B	2	ALA	2.6
1	A	177	LEU	2.6
1	A	139	ALA	2.5
1	B	174	GLU	2.5
1	A	183	ASN	2.5
1	A	182	SER	2.5
1	A	162	GLN	2.4
1	A	228	CYS	2.3
1	A	28	GLY	2.3
1	A	214	VAL	2.3
1	A	17	GLN	2.3
1	A	94	GLN	2.3
1	A	238	TYR	2.2
1	A	194	PRO	2.2
1	A	7	ASP	2.2
1	A	11	ASN	2.2
1	B	193	ASN	2.2
1	A	2	ALA	2.2
1	A	174	GLU	2.1
1	A	250	GLY	2.1
1	B	217	GLU	2.1
1	A	188	PRO	2.1
1	A	219	CYS	2.0
1	A	237	LYS	2.0
1	A	106	ARG	2.0
1	A	184	LEU	2.0
1	A	190	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	MG	B	304	1/1	0.92	0.16	5.26	20,20,20,20	0
5	0TF	B	301	18/18	0.89	0.21	2.92	27,31,43,45	0
3	NA	B	303	1/1	0.86	0.19	2.06	25,25,25,25	0
3	NA	A	902	1/1	0.65	0.33	1.95	41,41,41,41	0
5	0TF	A	904	18/18	0.89	0.17	1.49	36,39,48,49	0
4	MG	A	903	1/1	0.97	0.10	-0.27	38,38,38,38	0
2	NAI	B	302	44/44	0.97	0.08	-0.52	14,20,33,36	0
2	NAI	A	901	44/44	0.95	0.10	-0.64	32,38,42,43	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.