



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 03:26 PM BST

PDB ID : 1J9V
Title : Solution structure of a lactam analogue (DabD) of HIV gp41 600-612 loop.
Authors : Phan Chan Du, A.; Limal, D.; Semetey, V.; Dali, H.; Jolivet, M.; Desgranges, C.; Cung, M.T.; Briand, J.P.; Petit, M.C.; Muller, S.
Deposited on : 2001-05-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 47%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	14	

2 Ensemble composition and analysis ⓘ

This entry contains 49 models.

Cyrange was unable to find well-defined residues.

Error message: Only domains with < 8 residues could be identified.

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 199 atoms, of which 101 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41.

Mol	Chain	Residues	Atoms					Trace
1	A	14	Total	C	H	N	O	0
			199	63	101	16	19	

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ACE	-	INSERTION	UNP P12488
A	5	DAB	CYS	ENGINEERED	UNP P12488
A	11	ASP	CYS	ENGINEERED	UNP P12488

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A: 




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41


Chain A: 



4.2.2 Score per residue for model 2


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A: 




4.2.3 Score per residue for model 3

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41




4.2.4 Score per residue for model 4

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41




4.2.5 Score per residue for model 5

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41




4.2.6 Score per residue for model 6

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41




4.2.7 Score per residue for model 7

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41




4.2.8 Score per residue for model 8

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41




4.2.9 Score per residue for model 9

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41




4.2.10 Score per residue for model 10

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41




4.2.11 Score per residue for model 11

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41



4.2.12 Score per residue for model 12


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41



4.2.13 Score per residue for model 13


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41


Chain A:  86% 14%



4.2.14 Score per residue for model 14


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  86% 14%



4.2.15 Score per residue for model 15


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  86% 14%



4.2.16 Score per residue for model 16


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41


Chain A:  79% 21%



4.2.17 Score per residue for model 17


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  86% 14%



4.2.18 Score per residue for model 18


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.19 Score per residue for model 19


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.20 Score per residue for model 20


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.21 Score per residue for model 21


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.22 Score per residue for model 22


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.23 Score per residue for model 23


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.24 Score per residue for model 24


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.25 Score per residue for model 25


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.26 Score per residue for model 26

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.27 Score per residue for model 27

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  71% 21% 7%



4.2.28 Score per residue for model 28


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  71% 21% 7%



4.2.29 Score per residue for model 29


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.30 Score per residue for model 30


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.31 Score per residue for model 31


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.32 Score per residue for model 32

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.33 Score per residue for model 33

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  71% 29%



4.2.34 Score per residue for model 34


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  71% 29%



4.2.35 Score per residue for model 35

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.36 Score per residue for model 36


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  71% 29%



4.2.37 Score per residue for model 37


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.38 Score per residue for model 38


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.39 Score per residue for model 39


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.40 Score per residue for model 40


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41


Chain A:  79% 21%



4.2.41 Score per residue for model 41

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  86% 14%



4.2.42 Score per residue for model 42


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  71% 29%



4.2.43 Score per residue for model 43


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.44 Score per residue for model 44


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.45 Score per residue for model 45


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.46 Score per residue for model 46

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.47 Score per residue for model 47


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  71% 29%



4.2.48 Score per residue for model 48


- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



4.2.49 Score per residue for model 49

- Molecule 1: DabD (Ace)IWG(DAB)SGKLIDTTA ANALOGUE OF HIV GP41

Chain A:  79% 21%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *Torsion angle dynamics, molecular dynamics, energy minimization..*

Of the 50 calculated structures, 49 were deposited, based on the following criterion: *target function.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	refinement	1.5
DISCOVER	refinement	3

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5109
Number of chemical shift lists	1
Total number of shifts	91
Number of shifts mapped to atoms	91
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	47%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DAB, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.45±0.01	1±0/89 (1.1±0.0%)	1.32±0.05	1±0/117 (0.9±0.0%)
All	All	1.45	49/4361 (1.1%)	1.32	49/5733 (0.9%)

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	14	ALA	C-OXT	8.17	1.38	1.23	4	49

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	3	TRP	CD1-NE1-CE2	-5.70	103.87	109.00	28	49

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	98	101	98	0±0
All	All	4802	4949	4814	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:10:ILE:HD12	1:A:10:ILE:O	0.41	2.16	36	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	11/14 (79%)	6±1 (51±13%)	5±1 (48±12%)	0±0 (1±2%)	31	76
All	All	539/686 (79%)	276 (51%)	259 (48%)	4 (1%)	31	76

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	8	LYS	3
1	A	4	GLY	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	9/9 (100%)	8±1 (87±8%)	1±1 (13±8%)	9	50
All	All	441/441 (100%)	383 (87%)	58 (13%)	9	50

All 3 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	11	ASP	37
1	A	8	LYS	11
1	A	9	LEU	10

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	DAB	A	5	1	4,6,7	1.19±0.10	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	DAB	A	5	1	3,6,8	1.07±0.03	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DAB	A	5	1	-	0±0,3,5,7	0±0,0,0,0

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 47% for the well-defined parts and 47% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 5109

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	91
Number of shifts mapped to atoms	91
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 47%, i.e. 62 atoms were assigned a chemical shift out of a possible 132. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	24/60 (40%)	24/24 (100%)	0/24 (0%)	0/12 (0%)
Sidechain	32/60 (53%)	32/34 (94%)	0/25 (0%)	0/1 (0%)
Aromatic	6/12 (50%)	6/6 (100%)	0/5 (0%)	0/1 (0%)
Overall	62/132 (47%)	62/64 (97%)	0/54 (0%)	0/14 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 47%, i.e. 62 atoms were assigned a chemical shift out of a possible 132. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	24/60 (40%)	24/24 (100%)	0/24 (0%)	0/12 (0%)
Sidechain	32/60 (53%)	32/34 (94%)	0/25 (0%)	0/1 (0%)
Aromatic	6/12 (50%)	6/6 (100%)	0/5 (0%)	0/1 (0%)
Overall	62/132 (47%)	62/64 (97%)	0/54 (0%)	0/14 (0%)

7.1.4 Statistically unusual chemical shifts ⓘ

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

