



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:44 PM GMT

PDB ID : 4JDC
Title : Crystal structure of the Delta-pyrroline-5-carboxylate dehydrogenase from Mycobacterium tuberculosis
Authors : Lagautriere, T.; Bashiri, G.; Baker, E.N.
Deposited on : 2013-02-24
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	563	 79% 15% • 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4528 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-pyrroline-5-carboxylate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	536	4112	2604	729	768	11	0	1	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP I6X0K4
A	-18	GLY	-	EXPRESSION TAG	UNP I6X0K4
A	-17	SER	-	EXPRESSION TAG	UNP I6X0K4
A	-16	SER	-	EXPRESSION TAG	UNP I6X0K4
A	-15	HIS	-	EXPRESSION TAG	UNP I6X0K4
A	-14	HIS	-	EXPRESSION TAG	UNP I6X0K4
A	-13	HIS	-	EXPRESSION TAG	UNP I6X0K4
A	-12	HIS	-	EXPRESSION TAG	UNP I6X0K4
A	-11	HIS	-	EXPRESSION TAG	UNP I6X0K4
A	-10	HIS	-	EXPRESSION TAG	UNP I6X0K4
A	-9	SER	-	EXPRESSION TAG	UNP I6X0K4
A	-8	SER	-	EXPRESSION TAG	UNP I6X0K4
A	-7	GLY	-	EXPRESSION TAG	UNP I6X0K4
A	-6	LEU	-	EXPRESSION TAG	UNP I6X0K4
A	-5	VAL	-	EXPRESSION TAG	UNP I6X0K4
A	-4	PRO	-	EXPRESSION TAG	UNP I6X0K4
A	-3	ARG	-	EXPRESSION TAG	UNP I6X0K4
A	-2	GLY	-	EXPRESSION TAG	UNP I6X0K4
A	-1	SER	-	EXPRESSION TAG	UNP I6X0K4
A	0	HIS	-	EXPRESSION TAG	UNP I6X0K4
A	505	ASP	GLY	ENGINEERED MUTATION	UNP I6X0K4

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	27	10	5	10	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	389	Total	O	0	0
			389	389		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1-pyrroline-5-carboxylate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 6 2 2	Depositor
Cell constants a, b, c, α , β , γ	163.13Å 163.13Å 96.18Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.80 – 1.60 29.84 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.80-1.60) 99.8 (29.84-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.170 , 0.202 0.179 , 0.205	Depositor DCC
R_{free} test set	4908 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	21.7	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 98322 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4528	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CME, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.42	18/4201 (0.4%)	1.32	33/5728 (0.6%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	439	GLU	CD-OE1	10.00	1.36	1.25
1	A	519	TRP	CD2-CE2	6.98	1.49	1.41
1	A	95	TRP	CD2-CE2	6.97	1.49	1.41
1	A	363	SER	CA-CB	6.65	1.62	1.52
1	A	274	TRP	CD2-CE2	6.15	1.48	1.41
1	A	507	ASN	CG-ND2	6.06	1.48	1.32
1	A	340	TRP	CD2-CE2	5.93	1.48	1.41
1	A	199	PRO	N-CD	5.87	1.56	1.47
1	A	537	TYR	CG-CD2	5.84	1.46	1.39
1	A	513	PRO	N-CD	5.69	1.55	1.47
1	A	13	GLU	CD-OE1	5.66	1.31	1.25
1	A	429	PRO	N-CD	5.60	1.55	1.47
1	A	521	SER	CB-OG	5.51	1.49	1.42
1	A	462	ALA	N-CA	5.21	1.56	1.46
1	A	164	PRO	N-CD	5.20	1.55	1.47
1	A	399	TYR	CG-CD2	5.18	1.45	1.39
1	A	488	PRO	N-CD	5.10	1.54	1.47
1	A	441	TYR	CG-CD2	5.04	1.45	1.39

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	307	ARG	NE-CZ-NH1	-19.25	110.67	120.30
1	A	307	ARG	NE-CZ-NH2	15.56	128.08	120.30
1	A	361	ASP	CB-CG-OD1	10.61	127.85	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	25	ARG	NE-CZ-NH2	-7.90	116.35	120.30
1	A	69	ARG	NE-CZ-NH1	-7.29	116.65	120.30
1	A	109	ARG	NE-CZ-NH2	-7.18	116.71	120.30
1	A	244	ASP	CB-CG-OD1	7.05	124.65	118.30
1	A	518	ARG	NE-CZ-NH2	-6.68	116.96	120.30
1	A	25	ARG	NE-CZ-NH1	6.62	123.61	120.30
1	A	361	ASP	CB-CG-OD2	-6.60	112.36	118.30
1	A	500	GLY	N-CA-C	-6.46	96.94	113.10
1	A	177	ARG	NE-CZ-NH1	-6.45	117.08	120.30
1	A	41	ASP	CB-CG-OD2	-6.17	112.75	118.30
1	A	309	ASP	CB-CG-OD2	5.98	123.69	118.30
1	A	41	ASP	CB-CG-OD1	5.98	123.69	118.30
1	A	321	ASP	CB-CG-OD1	5.95	123.66	118.30
1	A	49	ARG	NE-CZ-NH2	5.87	123.23	120.30
1	A	70	LEU	CB-CG-CD2	5.81	120.88	111.00
1	A	332	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	A	371	ASP	CB-CG-OD1	5.70	123.43	118.30
1	A	416	ASP	C-N-CD	5.68	140.34	128.40
1	A	69	ARG	NE-CZ-NH2	5.50	123.05	120.30
1	A	383	GLU	OE1-CD-OE2	-5.49	116.72	123.30
1	A	529	PHE	CB-CG-CD1	-5.43	117.00	120.80
1	A	441	TYR	CZ-CE2-CD2	-5.43	114.92	119.80
1	A	92	LYS	CD-CE-NZ	-5.31	99.49	111.70
1	A	507	ASN	N-CA-CB	5.27	120.08	110.60
1	A	476	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	A	534	ASP	CB-CG-OD1	5.20	122.98	118.30
1	A	463	ASP	CB-CG-OD2	-5.14	113.67	118.30
1	A	54	ASP	CB-CG-OD1	5.13	122.92	118.30
1	A	487	LYS	C-N-CD	5.11	139.12	128.40
1	A	284	TYR	CD1-CE1-CZ	-5.03	115.28	119.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4112	0	4023	46	0
2	A	27	0	12	1	0
3	A	389	0	0	12	0
All	All	4528	0	4035	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:542:VAL:HA	1:A:543:ASP:C	2.00	0.82
1:A:474:ARG:HH11	1:A:474:ARG:HG2	1.50	0.76
1:A:299:PHE:HB3	3:A:1084:HOH:O	1.90	0.71
1:A:298:ASP:C	3:A:1084:HOH:O	2.29	0.70
1:A:298:ASP:OD2	1:A:328:SER:HB3	1.94	0.66
1:A:164:PRO:HD3	1:A:174:ILE:HG13	1.77	0.66
1:A:542:VAL:CA	1:A:543:ASP:C	2.66	0.62
1:A:418:THR:HG22	1:A:418:THR:O	1.98	0.62
1:A:480:GLY:O	1:A:481:ASN:ND2	2.28	0.62
1:A:373:ARG:NH1	3:A:809:HOH:O	2.34	0.60
1:A:489:THR:CG2	3:A:1038:HOH:O	2.51	0.57
1:A:355:ARG:HD2	3:A:893:HOH:O	2.03	0.57
1:A:474:ARG:HG2	1:A:474:ARG:NH1	2.14	0.57
1:A:136:GLN:OE1	1:A:326:LYS:HE3	2.05	0.57
1:A:240:LEU:C	1:A:240:LEU:HD23	2.26	0.56
1:A:299:PHE:N	3:A:1084:HOH:O	2.39	0.54
1:A:489:THR:HG23	3:A:1038:HOH:O	2.08	0.53
1:A:481:ASN:ND2	1:A:499[B]:GLY:HA3	2.24	0.53
1:A:261:ILE:HB	1:A:290:LEU:HD23	1.91	0.52
1:A:328:SER:O	1:A:489:THR:HG23	2.10	0.51
1:A:326:LYS:O	1:A:328:SER:N	2.44	0.51
1:A:253:LEU:HD23	1:A:276:TRP:CG	2.47	0.49
1:A:384:ARG:NH2	3:A:946:HOH:O	2.46	0.48
1:A:298:ASP:CG	1:A:457:THR:OG1	2.52	0.47
1:A:498:PHE:O	1:A:499[B]:GLY:O	2.31	0.47
1:A:346:GLU:HG3	3:A:932:HOH:O	2.14	0.47
1:A:180:ASP:O	1:A:259:ALA:HB2	2.15	0.47
1:A:489:THR:HG22	3:A:1038:HOH:O	2.14	0.46
1:A:25:ARG:O	1:A:29:ARG:HG2	2.16	0.46
1:A:352:ALA:HB2	1:A:395:VAL:HG21	1.98	0.46
1:A:227:LEU:C	1:A:227:LEU:HD23	2.37	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:514:LEU:N	1:A:514:LEU:HD22	2.31	0.45
1:A:249:SER:OG	2:A:601:NAD:N1A	2.46	0.45
1:A:150:ARG:HD2	3:A:728:HOH:O	2.16	0.44
1:A:507:ASN:HA	3:A:888:HOH:O	2.18	0.43
1:A:240:LEU:HD23	1:A:241:VAL:N	2.34	0.43
1:A:187:THR:HG22	1:A:198:LEU:HD12	2.00	0.43
1:A:280:ASN:O	1:A:283:ARG:HG2	2.18	0.42
1:A:263:PHE:O	1:A:292:GLY:HA3	2.19	0.42
1:A:456:LEU:HG	1:A:457:THR:HG23	2.02	0.41
1:A:223:LEU:HD23	1:A:223:LEU:HA	1.95	0.41
1:A:225:MET:HG3	1:A:240:LEU:HD12	2.01	0.41
1:A:298:ASP:OD2	1:A:457:THR:OG1	2.38	0.40
1:A:326:LYS:O	1:A:327:CME:C	2.69	0.40
1:A:444:ILE:HA	1:A:444:ILE:HD13	1.98	0.40
1:A:326:LYS:C	1:A:328:SER:N	2.72	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	532/563 (94%)	517 (97%)	13 (2%)	2 (0%)	39 17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	499[A]	GLY
1	A	499[B]	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	415/439 (94%)	408 (98%)	7 (2%)	68 44

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	131	SER
1	A	212	LYS
1	A	328	SER
1	A	453	ARG
1	A	474	ARG
1	A	489	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	481	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CME	A	327	1	8,9,10	0.78	0	6,9,11	3.51	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	327	1	-	0/5/8/10	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	327	CME	O-C-CA	-2.55	118.84	125.49
1	A	327	CME	CB-SG-SD	7.87	119.29	103.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	327	CME	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAD	A	601	-	23,29,48	2.17	5 (21%)	27,45,73	2.18	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	601	-	-	0/12/32/62	0/3/3/5

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	NAD	C5A-C4A	2.09	1.45	1.40
2	A	601	NAD	PN-O5D	2.17	1.63	1.55
2	A	601	NAD	PN-O3	3.11	1.68	1.61
2	A	601	NAD	PN-O2N	3.36	1.61	1.50
2	A	601	NAD	O4B-C1B	7.42	1.50	1.41

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	NAD	N3A-C2A-N1A	-7.31	123.30	128.89
2	A	601	NAD	C1B-N9A-C4A	-2.70	122.87	126.94
2	A	601	NAD	O5B-C5B-C4B	-2.54	99.76	109.12
2	A	601	NAD	O2A-PA-O3	2.11	114.66	105.09
2	A	601	NAD	O2N-PN-O3	2.14	114.88	106.09
2	A	601	NAD	O2A-PA-O1A	2.48	125.94	112.53
2	A	601	NAD	C2B-C1B-N9A	5.12	122.11	114.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	NAD	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	535/563 (95%)	-0.33	7 (1%) 79 79	14, 22, 41, 90	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	426	TYR	7.4
1	A	454	TYR	6.7
1	A	543	ASP	3.4
1	A	453	ARG	2.8
1	A	452	SER	2.8
1	A	490	GLY	2.7
1	A	542	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CME	A	327	10/11	0.77	0.14	-	45,60,78,85	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAD	A	601	27/44	0.90	0.10	0.99	20,29,37,45	27

6.5 Other polymers [i](#)

There are no such residues in this entry.