



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 08:15 PM GMT

PDB ID : 1JIE  
Title : Crystal structure of bleomycin-binding protein from bleomycin-producing *Streptomyces verticillus* complexed with metal-free bleomycin  
Authors : Sugiyama, M.; Kumagai, T.; Hayashida, M.; Maruyama, M.; Matoba, Y.  
Deposited on : 2001-07-02  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	122	
1	B	122	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2229 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called bleomycin-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	122	Total	C	N	O	S	0	3	0
			950	595	169	183	3			
1	B	122	Total	C	N	O	S	0	2	0
			948	593	169	183	3			

- Molecule 2 is BLEOMYCIN A2 (three-letter code: BLM) (formula:  $C_{55}H_{85}N_{17}O_{21}S_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	S	0	0
			96	55	17	21	3		
2	A	1	Total	C	N	O	S	0	0
			96	55	17	21	3		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	73	Total 73	O 73	0	0
3	B	66	Total 66	O 66	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: bleomycin-binding protein

Chain A:  84% 13% . .



- Molecule 1: bleomycin-binding protein

Chain B:  81% 16% .



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	35.52Å 83.42Å 43.24Å 90.00° 104.88° 90.00°	Depositor
Resolution (Å)	10.00 – 1.80	Depositor
% Data completeness (in resolution range)	75.9 (10.00-1.80)	Depositor
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.156 , 0.218	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2229	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BLM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	0/988	1.42	8/1346 (0.6%)
1	B	0.53	0/980	1.31	1/1335 (0.1%)
All	All	0.54	0/1968	1.37	9/2681 (0.3%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	47	ARG	NE-CZ-NH1	8.19	124.39	120.30
1	A	33	PHE	CB-CG-CD1	7.27	125.89	120.80
1	A	42	ARG	NE-CZ-NH1	6.89	123.74	120.30
1	A	109	ARG	NE-CZ-NH2	-6.45	117.07	120.30
1	B	270	ASP	CB-CG-OD1	6.41	124.07	118.30
1	A	110	ASP	CB-CG-OD1	5.55	123.30	118.30
1	A	42	ARG	CD-NE-CZ	5.44	131.21	123.60
1	A	80	ARG	NE-CZ-NH1	-5.12	117.74	120.30
1	A	42	ARG	NE-CZ-NH2	-5.01	117.80	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	950	0	898	17	0
1	B	948	0	894	14	0
2	A	96	0	85	6	0
2	B	96	0	85	6	0
3	A	73	0	0	4	0
3	B	66	0	0	4	0
All	All	2229	0	1962	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:402:BLM:H541	2:A:402:BLM:H44	1.67	0.76
2:B:401:BLM:H611	3:B:609:HOH:O	1.86	0.75
1:A:47:ARG:HD2	2:B:401:BLM:H543	1.72	0.70
1:B:284:THR:HG22	3:B:614:HOH:O	1.97	0.65
1:A:80:ARG:N	1:A:80:ARG:HE	1.94	0.64
1:A:47:ARG:HH11	2:B:401:BLM:H541	1.62	0.63
1:B:285:ASP:OD2	1:B:288:ASP:HB2	1.97	0.63
2:A:402:BLM:O12	2:A:402:BLM:HAA	2.01	0.59
1:A:47:ARG:HH11	2:B:401:BLM:C54	2.15	0.59
1:A:89:THR:HG21	3:A:559:HOH:O	2.05	0.57
1:B:216:VAL:HB	1:B:217:PRO:HD3	1.87	0.57
2:B:401:BLM:HAA	2:B:401:BLM:O12	2.06	0.55
1:A:89:THR:HG23	3:A:620:HOH:O	2.09	0.53
1:A:1:MET:O	1:A:1:MET:HG3	2.08	0.53
1:B:292:PRO:HG3	1:B:311:PRO:HG3	1.93	0.50
2:A:402:BLM:H542	1:B:233:PHE:CE2	2.47	0.49
1:B:322:GLU:OXT	1:B:322:GLU:HG3	2.12	0.48
1:B:254:GLU:HG2	3:B:554:HOH:O	2.13	0.47
1:B:288:ASP:CG	1:B:290:SER:HG	2.18	0.47
1:A:32:ASP:CG	1:A:42:ARG:HH21	2.18	0.47
1:A:89:THR:HG22	1:A:89:THR:O	2.16	0.45
1:B:214:VAL:HG22	3:B:518:HOH:O	2.16	0.45
1:A:80:ARG:CA	1:A:80:ARG:HE	2.28	0.45
1:A:6:GLY:HA2	3:A:618:HOH:O	2.17	0.45
1:A:65:TRP:O	1:A:66[B]:ILE:HD13	2.17	0.44
1:B:232:ASP:OD2	1:B:242[B]:ARG:NH2	2.50	0.44
1:B:236[B]:ARG:HB2	1:B:236[B]:ARG:HE	1.28	0.44
1:A:115:CYS:HB2	2:A:402:BLM:HAC	2.00	0.43

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:ARG:HB2	1:A:47:ARG:HG2	2.00	0.43
1:A:71:PRO:HD2	3:A:544:HOH:O	2.17	0.43
1:B:268:VAL:O	1:B:321:GLY:N	2.48	0.43
1:B:232:ASP:CG	1:B:242[B]:ARG:HE	2.23	0.42
1:A:77:GLU:O	1:A:80:ARG:HG2	2.19	0.42
1:A:47:ARG:HD2	2:B:401:BLM:C54	2.46	0.41
1:B:282:VAL:HB	1:B:294:MET:CE	2.50	0.41
2:A:402:BLM:HNP	2:A:402:BLM:H522	1.43	0.41
2:A:402:BLM:C54	2:A:402:BLM:H44	2.45	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	123/122 (101%)	122 (99%)	1 (1%)	0	100	100
1	B	122/122 (100%)	120 (98%)	2 (2%)	0	100	100
All	All	245/244 (100%)	242 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	100/97 (103%)	91 (91%)	9 (9%)	12	3
1	B	99/97 (102%)	93 (94%)	6 (6%)	23	8
All	All	199/194 (103%)	184 (92%)	15 (8%)	18	5

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LEU
1	A	32	ASP
1	A	36[A]	ARG
1	A	36[B]	ARG
1	A	42	ARG
1	A	76	GLU
1	A	80	ARG
1	A	105	GLU
1	A	122	GLU
1	B	201	MET
1	B	203	LYS
1	B	217	PRO
1	B	254	GLU
1	B	289	THR
1	B	322	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	249	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	BLM	A	402	-	89,101,101	1.87	13 (14%)	102,143,143	2.05	29 (28%)
2	BLM	B	401	-	89,101,101	1.85	10 (11%)	102,143,143	2.17	34 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BLM	A	402	-	-	0/85/142/142	0/6/6/6
2	BLM	B	401	-	-	0/85/142/142	0/6/6/6

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	BLM	C10-C9	-5.78	1.31	1.40
2	B	401	BLM	C7-NE	-5.68	1.24	1.34
2	A	402	BLM	C10-C9	-5.06	1.32	1.40
2	A	402	BLM	C7-NE	-4.75	1.25	1.34
2	A	402	BLM	C10-C12	-4.62	1.43	1.50
2	A	402	BLM	C45-C46	-4.27	1.41	1.49
2	B	401	BLM	C10-C12	-4.19	1.43	1.50
2	B	401	BLM	C45-C46	-4.09	1.41	1.49
2	B	401	BLM	C42-C43	-3.61	1.47	1.49
2	A	402	BLM	C42-C43	-3.41	1.47	1.49
2	A	402	BLM	C46-NO	-3.01	1.27	1.31
2	B	401	BLM	C46-NO	-2.17	1.28	1.31
2	A	402	BLM	C46-S46	2.00	1.76	1.73
2	A	402	BLM	C47-S46	2.02	1.73	1.70
2	A	402	BLM	C70-NQ	2.03	1.38	1.33
2	A	402	BLM	C12-NH	2.03	1.38	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	BLM	C7-NG	2.11	1.38	1.34
2	A	402	BLM	C1-NA	2.96	1.38	1.32
2	B	401	BLM	C1-NA	3.21	1.39	1.32
2	B	401	BLM	C8-NE	6.84	1.45	1.35
2	A	402	BLM	C8-NE	7.94	1.47	1.35
2	A	402	BLM	C10-NG	8.51	1.48	1.34
2	B	401	BLM	C10-NG	9.12	1.49	1.34

All (63) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402	BLM	C9-C10-NG	-5.81	117.97	123.25
2	B	401	BLM	CA-C9-C8	-5.09	114.83	120.05
2	B	401	BLM	O12-C12-C10	-3.76	115.87	121.00
2	B	401	BLM	O1-C1-NA	-3.64	117.68	123.08
2	B	401	BLM	C63-OH1-C14	-3.55	109.38	115.80
2	B	401	BLM	O62-C60-C59	-3.54	103.03	109.68
2	B	401	BLM	O70-C70-NQ	-3.43	118.77	125.34
2	B	401	BLM	C12-C10-NG	-3.32	109.05	115.31
2	A	402	BLM	O4-C4-C5	-3.26	113.85	120.93
2	A	402	BLM	OH3-C38-C37	-3.24	102.09	109.08
2	B	401	BLM	C9-C10-NG	-3.23	120.31	123.25
2	A	402	BLM	C48-C49-NP	-3.17	111.15	115.42
2	B	401	BLM	C42-C41-NM	-3.17	104.25	111.97
2	B	401	BLM	O49-C49-C48	-2.80	115.28	121.23
2	A	402	BLM	C12-C10-NG	-2.68	110.26	115.31
2	A	402	BLM	C27-C28-NI	-2.48	104.08	108.56
2	A	402	BLM	C45-C44-S43	-2.45	108.78	111.79
2	B	401	BLM	C3-C2-NB	-2.43	105.72	109.56
2	B	401	BLM	O36-C36-NL	-2.40	118.22	122.93
2	B	401	BLM	OH1-C63-O62	-2.29	104.88	110.68
2	A	402	BLM	O70-C70-NQ	-2.28	120.98	125.34
2	A	402	BLM	C33-C34-C36	-2.17	106.61	109.55
2	A	402	BLM	C3-C2-NB	-2.14	106.18	109.56
2	B	401	BLM	C27-C28-NI	-2.14	104.70	108.56
2	A	402	BLM	CA-C9-C10	-2.14	118.41	122.99
2	A	402	BLM	O59-C59-C58	-2.12	105.56	110.34
2	B	401	BLM	O58-C58-C57	-2.11	104.89	109.87
2	B	401	BLM	C50-NP-C49	-2.09	117.47	122.15
2	A	402	BLM	OH1-C63-O62	-2.08	105.41	110.68
2	B	401	BLM	C51-C52-S53	-2.04	107.73	111.63
2	B	401	BLM	O56-C64-O64	-2.02	105.56	110.68

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	BLM	C5-C4-ND	2.05	120.59	116.08
2	B	401	BLM	C55-S53-C52	2.09	105.61	101.73
2	A	402	BLM	C50-NP-C49	2.15	126.96	122.15
2	A	402	BLM	O64-C65-C67	2.17	113.76	109.68
2	B	401	BLM	C63-O62-C60	2.22	118.05	113.75
2	B	401	BLM	O36-C36-C34	2.32	125.91	120.56
2	A	402	BLM	C64-C69-C68	2.32	114.17	110.00
2	A	402	BLM	O62-C60-C59	2.33	114.06	109.68
2	B	401	BLM	C58-C59-C60	2.39	114.36	110.20
2	A	402	BLM	C64-O64-C65	2.58	118.75	113.75
2	B	401	BLM	C64-C69-C68	2.61	114.70	110.00
2	A	402	BLM	C55-S53-C52	2.68	106.70	101.73
2	B	401	BLM	C28-NI-C29	2.83	110.17	105.71
2	A	402	BLM	O56-C57-C58	2.90	114.65	107.17
2	A	402	BLM	C37-NL-C36	2.97	129.67	121.85
2	B	401	BLM	O58-C58-C59	2.99	117.06	110.34
2	A	402	BLM	O1-C1-C2	3.01	124.43	120.33
2	B	401	BLM	C64-O64-C65	3.03	119.62	113.75
2	A	402	BLM	C5-C4-ND	3.08	122.88	116.08
2	B	401	BLM	O68-C68-C69	3.17	114.80	107.76
2	B	401	BLM	O62-C63-C57	3.62	116.78	109.47
2	A	402	BLM	C54-S53-C52	3.73	108.65	101.73
2	B	401	BLM	O1-C1-C2	3.74	125.43	120.33
2	A	402	BLM	C58-C59-C60	3.97	117.11	110.20
2	A	402	BLM	O68-C68-C69	4.05	116.75	107.76
2	A	402	BLM	O49-C49-NP	4.18	131.37	122.58
2	B	401	BLM	O49-C49-NP	4.27	131.56	122.58
2	B	401	BLM	C68-O68-C70	4.48	123.49	117.03
2	B	401	BLM	C10-C9-C8	6.00	120.83	115.69
2	A	402	BLM	CB-C31-NK	6.39	116.96	109.64
2	B	401	BLM	NF-C8-NE	7.62	128.00	116.95
2	A	402	BLM	C10-C9-C8	7.63	122.22	115.69

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	402	BLM	6	0
2	B	401	BLM	6	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.