



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:58 AM GMT

PDB ID : 3KDT
Title : Crystal structure of peroxisome proliferator-activated receptor alpha (PPAR α) complex with N-3-((2-(4-Chlorophenyl)-5-methyl-1,3-oxazol-4-yl)methoxy)benzyl)-N-(methoxycarbonyl)glycine
Authors : Muckelbauer, J.K.
Deposited on : 2009-10-23
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	<div> <div>5%</div> <div>77%</div> <div>17%</div> <div>• •</div> </div>
1	B	277	<div> <div>5%</div> <div>78%</div> <div>18%</div> <div>• •</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4245 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peroxisome proliferator-activated receptor alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	0	1
			2068	1324	345	381	18			
1	B	272	Total	C	N	O	S	0	0	1
			2113	1352	353	390	18			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	192	GLY	-	EXPRESSION TAG	UNP Q07869
A	193	SER	-	EXPRESSION TAG	UNP Q07869
A	194	HIS	-	EXPRESSION TAG	UNP Q07869
A	195	MET	-	EXPRESSION TAG	UNP Q07869
B	192	GLY	-	EXPRESSION TAG	UNP Q07869
B	193	SER	-	EXPRESSION TAG	UNP Q07869
B	194	HIS	-	EXPRESSION TAG	UNP Q07869
B	195	MET	-	EXPRESSION TAG	UNP Q07869

- Molecule 2 is N-(3-{[2-(4-CHLOROPHENYL)-5-METHYL-1,3-OXAZOL-4-YL]METHOXY}BENZYL)-N-(METHOXYCARBONYL)GLYCINE (three-letter code: 7HA) (formula: C₂₂H₂₁ClN₂O₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			31	22	1	2	6		
2	B	1	Total	C	Cl	N	O	0	0
			31	22	1	2	6		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor alpha



- Molecule 1: Peroxisome proliferator-activated receptor alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	64.75Å 64.75Å 123.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.70 45.79 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.0 (50.00-2.70) 99.0 (45.79-2.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.253 , 0.316 0.251 , 0.317	Depositor DCC
R_{free} test set	688 reflections (5.22%)	DCC
Wilson B-factor (Å ²)	49.3	Xtriage
Anisotropy	0.585	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 23.9	EDS
Estimated twinning fraction	0.469 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	1 of 13909 reflections (0.007%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4245	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 7HA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/2104	0.59	0/2839
1	B	0.48	0/2149	0.58	0/2899
All	All	0.48	0/4253	0.58	0/5738

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2068	0	2074	27	0
1	B	2113	0	2101	30	0
2	A	31	0	20	1	0
2	B	31	0	20	2	0
3	A	2	0	0	0	0
All	All	4245	0	4215	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:418:ASP:OD1	1:B:418:ASP:N	2.24	0.71
1:B:276:CYS:SG	2:B:501:7HA:H1B	2.33	0.69
1:A:388:ARG:O	1:A:391:LEU:HG	1.93	0.68
1:A:341:ARG:NH2	1:A:356:GLU:OE2	2.25	0.65
1:A:235:ASN:CG	1:A:236:ASN:H	1.99	0.65
1:B:207:ALA:HA	1:B:377:LEU:HD21	1.79	0.64
1:B:385:CYS:SG	1:B:387:ASP:HB2	2.38	0.64
1:A:234:SER:O	1:A:235:ASN:HB3	1.99	0.63
1:A:418:ASP:OD1	1:A:418:ASP:N	2.25	0.62
1:A:391:LEU:HD13	1:A:397:ILE:HD13	1.82	0.61
1:A:351:PHE:CE1	1:A:451:GLU:HG3	2.36	0.60
1:A:276:CYS:SG	2:A:501:7HA:H1B	2.42	0.60
1:A:234:SER:O	1:A:235:ASN:CB	2.50	0.59
1:A:464:TYR:HA	1:A:467:MET:HG2	1.86	0.58
1:B:388:ARG:O	1:B:391:LEU:HG	2.03	0.58
1:A:280:SER:O	1:A:284:VAL:HG23	2.04	0.57
1:B:267:GLU:HG2	1:B:268:ALA:H	1.69	0.57
1:B:341:ARG:NH2	1:B:356:GLU:OE2	2.35	0.56
1:B:465:ARG:O	1:B:466:ASP:HB2	2.07	0.55
1:B:200:THR:HG22	1:B:203:LEU:HD12	1.90	0.53
1:B:276:CYS:SG	2:B:501:7HA:C1	2.96	0.52
1:A:351:PHE:HE1	1:A:451:GLU:HG3	1.75	0.52
1:B:246:THR:O	1:B:250:ALA:HB2	2.09	0.50
1:A:230:SER:HB2	1:A:232:LYS:H	1.77	0.50
1:A:324:VAL:HB	1:A:331:LEU:HB2	1.93	0.49
1:A:235:ASN:CG	1:A:236:ASN:N	2.66	0.48
1:A:210:ILE:HG22	1:A:376:SER:HB2	1.96	0.48
1:A:354:ILE:HD11	1:A:447:ILE:HD12	1.96	0.47
1:B:409:ARG:HG3	1:B:423:PHE:CD2	2.49	0.47
1:A:206:LEU:HD12	1:A:295:PRO:HG2	1.96	0.47
1:B:223:VAL:O	1:B:227:VAL:HG23	2.16	0.46
1:B:351:PHE:HE1	1:B:451:GLU:HG3	1.81	0.46
1:B:198:SER:HA	1:B:201:ALA:HB3	1.97	0.46
1:B:356:GLU:HB3	1:B:357:PRO:HD3	1.97	0.46
1:A:356:GLU:HB3	1:A:357:PRO:HD3	1.99	0.45
1:A:421:PHE:C	1:A:424:PRO:HD2	2.38	0.44
1:B:248:CYS:HA	1:B:251:GLU:OE1	2.17	0.44
1:B:267:GLU:O	1:B:271:ARG:NH1	2.51	0.44
1:B:235:ASN:O	1:B:237:PRO:HD3	2.17	0.44
1:B:250:ALA:HB1	1:B:254:LEU:HD12	1.98	0.44
1:B:281:VAL:HG21	1:B:457:HIS:CD2	2.53	0.44
1:A:209:ARG:HD2	1:A:295:PRO:HG3	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:363:MET:CE	1:B:363:MET:HA	2.49	0.42
1:B:251:GLU:HA	1:B:255:VAL:HG22	2.01	0.42
1:B:405:VAL:HG13	1:B:423:PHE:HZ	1.84	0.42
1:A:291:ALA:HA	1:A:294:ILE:HD12	2.00	0.42
1:A:426:LEU:HA	1:A:426:LEU:HD23	1.76	0.42
1:A:419:ASP:OD2	1:A:425:LYS:NZ	2.39	0.42
1:B:236:ASN:HA	1:B:237:PRO:HD3	1.85	0.42
1:B:267:GLU:O	1:B:271:ARG:HG3	2.19	0.41
1:A:243:ASP:HA	1:A:343:PHE:CE1	2.55	0.41
1:B:269:GLU:HB3	1:B:351:PHE:CD2	2.55	0.41
1:B:412:LEU:HD22	1:B:422:LEU:HD23	2.03	0.41
1:B:340:THR:OG1	1:B:343:PHE:CB	2.69	0.41
1:A:302:LEU:O	1:A:306:VAL:HG23	2.21	0.41
1:A:361:PHE:CZ	1:A:433:LEU:HG	2.56	0.41
1:B:324:VAL:HB	1:B:331:LEU:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/277 (94%)	239 (92%)	18 (7%)	4 (2%)	13	32
1	B	266/277 (96%)	242 (91%)	19 (7%)	5 (2%)	10	25
All	All	527/554 (95%)	481 (91%)	37 (7%)	9 (2%)	11	29

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	235	ASN
1	B	466	ASP
1	B	238	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	263	ILE
1	B	267	GLU
1	B	465	ARG
1	A	244	MET
1	A	252	LYS
1	A	238	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/242 (93%)	215 (96%)	10 (4%)	35	65
1	B	226/242 (93%)	217 (96%)	9 (4%)	38	69
All	All	451/484 (93%)	432 (96%)	19 (4%)	36	68

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	243	ASP
1	A	246	THR
1	A	247	LEU
1	A	275	CYS
1	A	346	SER
1	A	418	ASP
1	A	439	GLU
1	A	448	LYS
1	A	466	ASP
1	A	467	MET
1	B	219	ASN
1	B	243	ASP
1	B	245	GLU
1	B	269	GLU
1	B	275	CYS
1	B	306	VAL
1	B	418	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	444	VAL
1	B	448	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	7HA	A	501	-	26,33,33	0.77	2 (7%)	33,45,45	2.17	9 (27%)
2	7HA	B	501	-	26,33,33	0.78	2 (7%)	33,45,45	2.10	8 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7HA	A	501	-	-	0/17/23/23	0/2/3/3
2	7HA	B	501	-	-	0/17/23/23	0/2/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	7HA	O15-C16	-2.13	1.32	1.37
2	A	501	7HA	O15-C16	-2.02	1.32	1.37
2	A	501	7HA	C1-C2	2.01	1.51	1.48
2	B	501	7HA	C1-C2	2.02	1.51	1.48

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	7HA	C20-C22-N23	-4.06	106.19	113.10
2	A	501	7HA	C12-C7-C5	-3.49	114.80	120.51
2	B	501	7HA	C20-C22-N23	-2.95	108.08	113.10
2	B	501	7HA	O30-C28-O29	-2.81	120.15	124.67
2	B	501	7HA	C12-C7-C5	-2.76	116.00	120.51
2	A	501	7HA	O30-C28-O29	-2.63	120.43	124.67
2	B	501	7HA	O29-C28-N23	-2.49	119.58	124.32
2	A	501	7HA	O29-C28-N23	-2.35	119.86	124.32
2	A	501	7HA	C9-C10-CL13	2.06	122.73	119.35
2	B	501	7HA	C14-O15-C16	2.16	123.50	117.70
2	B	501	7HA	C8-C7-C5	2.21	124.12	120.51
2	A	501	7HA	C8-C7-C5	2.55	124.68	120.51
2	A	501	7HA	C14-O15-C16	2.57	124.58	117.70
2	B	501	7HA	C31-O30-C28	3.17	119.35	115.40
2	A	501	7HA	C31-O30-C28	3.66	119.96	115.40
2	A	501	7HA	O30-C28-N23	8.11	119.70	111.53
2	B	501	7HA	O30-C28-N23	8.67	120.27	111.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	7HA	1	0
2	B	501	7HA	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/277 (95%)	0.41	14 (5%) 30 28	32, 54, 88, 94	0
1	B	272/277 (98%)	0.38	14 (5%) 32 30	29, 55, 97, 112	0
All	All	537/554 (96%)	0.39	28 (5%) 31 30	29, 54, 94, 112	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	263	ILE	16.5
1	A	264	GLN	7.1
1	A	254	LEU	5.3
1	A	454	ALA	5.1
1	B	194	HIS	5.1
1	B	193	SER	5.0
1	B	198	SER	4.3
1	A	265	ASN	3.4
1	B	254	LEU	3.4
1	A	227	VAL	3.2
1	B	443	LEU	3.2
1	B	262	GLY	3.0
1	B	196	GLU	2.7
1	B	268	ALA	2.7
1	B	197	ASP	2.6
1	B	255	VAL	2.6
1	A	452	SER	2.6
1	A	453	ASP	2.5
1	A	255	VAL	2.3
1	A	311	TYR	2.3
1	A	269	GLU	2.2
1	A	347	LEU	2.1
1	B	449	LYS	2.1
1	B	338	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	274	HIS	2.1
1	A	231	GLY	2.1
1	A	250	ALA	2.1
1	B	228	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	7HA	B	501	31/31	0.87	0.30	1.26	52,55,68,69	0
2	7HA	A	501	31/31	0.88	0.20	-0.19	48,53,65,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.