



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 08:29 PM GMT

PDB ID : 1KHF  
Title : PEPCK complex with PEP  
Authors : Dunten, P.; Belunis, C.; Crowther, R.; Hollfelder, K.; Kammlott, U.; Levin, W.; Michel, H.; Ramsey, G.B.; Swain, A.; Weber, D.; Wertheimer, S.J.  
Deposited on : 2001-11-29  
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	7858 (2.04-2.00)
Clashscore	102246	9060 (2.04-2.00)
Ramachandran outliers	100387	8952 (2.04-2.00)
Sidechain outliers	100360	8951 (2.04-2.00)
RSRZ outliers	91569	7873 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	625	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4905 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoenolpyruvate Carboxykinase, cytosolic (GTP).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	603	Total	C	N	O	S	0	0	0
			4721	3012	807	867	35			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	CLONING ARTIFACT	UNP P35558
A	-1	GLU	-	CLONING ARTIFACT	UNP P35558
A	0	LEU	-	CLONING ARTIFACT	UNP P35558
A	267	VAL	ILE	VARIANT	UNP P35558
A	586	ASP	GLU	VARIANT	UNP P35558
A	597	VAL	GLU	VARIANT	UNP P35558


- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mn	0	0
			1	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is PHOSPHOENOLPYRUVATE (three-letter code: PEP) (formula: C<sub>3</sub>H<sub>5</sub>O<sub>6</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	164	Total 164	O 164	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphoenolpyruvate Carboxykinase, cytosolic (GTP)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.25Å 60.68Å 62.03Å 88.69° 70.00° 72.54°	Depositor
Resolution (Å)	20.00 – 2.02 29.83 – 2.02	Depositor EDS
% Data completeness (in resolution range)	89.2 (20.00-2.02) 86.6 (29.83-2.02)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.27 (at 2.01Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.183 , 0.244 0.183 , 0.244	Depositor DCC
$R_{free}$ test set	1724 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.7	Xtriage
Anisotropy	0.306	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 42.7	EDS
Estimated twinning fraction	0.012 for -h,-l,-k	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 34475 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4905	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MN, PEP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.28	8/4842 (0.2%)	1.16	20/6557 (0.3%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	140	TYR	CD1-CE1	7.26	1.50	1.39
1	A	516	TRP	CE3-CZ3	6.57	1.49	1.38
1	A	95	VAL	CB-CG1	6.03	1.65	1.52
1	A	463	GLU	CD-OE1	5.41	1.31	1.25
1	A	89	GLU	CD-OE1	5.40	1.31	1.25
1	A	536	VAL	CB-CG1	5.37	1.64	1.52
1	A	196	VAL	CB-CG1	5.24	1.63	1.52
1	A	316	LYS	CD-CE	5.14	1.64	1.51

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	591	ASP	CB-CG-OD2	9.72	127.05	118.30
1	A	562	ASP	CB-CG-OD2	8.99	126.39	118.30
1	A	253	ARG	NE-CZ-NH2	-7.84	116.38	120.30
1	A	545	ASP	CB-CG-OD2	6.82	124.44	118.30
1	A	365	ASP	CB-CG-OD2	6.60	124.24	118.30
1	A	84	ASP	CB-CG-OD2	6.32	123.99	118.30
1	A	87	ARG	NE-CZ-NH1	-6.29	117.15	120.30
1	A	318	ASP	CB-CG-OD1	6.29	123.96	118.30
1	A	520	ASP	CB-CG-OD1	6.15	123.83	118.30
1	A	253	ARG	NE-CZ-NH1	6.05	123.33	120.30
1	A	395	ASP	CB-CG-OD2	6.02	123.72	118.30
1	A	115	ARG	NE-CZ-NH1	-5.83	117.38	120.30
1	A	186	ASP	CB-CG-OD2	5.81	123.53	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	21	ASP	CB-CG-OD2	5.72	123.45	118.30
1	A	73	ASP	CB-CG-OD2	5.61	123.35	118.30
1	A	41	ASP	CB-CG-OD2	5.42	123.17	118.30
1	A	101	ASP	CB-CG-OD2	5.40	123.16	118.30
1	A	310	ASP	CB-CG-OD2	5.28	123.05	118.30
1	A	562	ASP	CB-CG-OD1	-5.24	113.59	118.30
1	A	603	ASP	CB-CG-OD2	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4721	0	4663	76	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	10	0	2	0	0
5	A	8	0	12	2	0
6	A	164	0	0	1	0
All	All	4905	0	4677	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:ARG:HH11	1:A:100:ARG:HG3	0.98	1.14
1:A:100:ARG:NH1	1:A:100:ARG:HG3	1.77	0.93
1:A:473:LYS:C	1:A:474:ILE:HD13	1.92	0.89
1:A:10:ASN:CG	1:A:11:LEU:H	1.79	0.86
1:A:10:ASN:ND2	1:A:11:LEU:H	1.77	0.82
1:A:594:LYS:HE3	1:A:598:ASP:OD2	1.86	0.75
1:A:387:LYS:HG3	1:A:389:LYS:HE2	1.70	0.72

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:473:LYS:N	1:A:473:LYS:HD3	2.06	0.70
1:A:619:ILE:O	1:A:622:MET:HG2	1.93	0.69
1:A:255:ALA:HB1	1:A:260:TRP:O	1.93	0.69
1:A:49:SER:HB2	1:A:51:GLU:OE1	1.93	0.68
1:A:10:ASN:CG	1:A:11:LEU:N	2.48	0.67
1:A:473:LYS:O	1:A:474:ILE:HD13	1.95	0.66
1:A:100:ARG:HH11	1:A:100:ARG:CG	1.90	0.63
1:A:558:ILE:HB	1:A:559:PRO:CD	2.30	0.61
1:A:377:LEU:HD11	1:A:383:ILE:HD11	1.82	0.61
1:A:474:ILE:N	1:A:474:ILE:HD13	2.15	0.59
1:A:586:ASP:O	1:A:590:GLU:HG3	2.07	0.55
1:A:287:ALA:O	1:A:436:ARG:HD2	2.07	0.55
1:A:255:ALA:HB1	1:A:260:TRP:C	2.28	0.53
1:A:558:ILE:HB	1:A:559:PRO:HD2	1.91	0.53
1:A:10:ASN:ND2	1:A:11:LEU:N	2.53	0.53
1:A:38:CYS:HB3	1:A:134:MET:HG3	1.92	0.52
1:A:200:LEU:HA	1:A:201:PRO:C	2.30	0.52
1:A:142:ILE:HG23	1:A:176:MET:CG	2.39	0.52
1:A:594:LYS:CE	1:A:598:ASP:OD2	2.59	0.50
1:A:293:LEU:HD12	1:A:296:MET:HE3	1.92	0.50
1:A:288:CYS:SG	1:A:436:ARG:HG3	2.53	0.49
1:A:498:LEU:HD13	1:A:607:GLU:HB2	1.94	0.49
1:A:221:HIS:NE2	1:A:249:ARG:HD2	2.28	0.49
1:A:100:ARG:NH1	1:A:100:ARG:CG	2.60	0.49
1:A:253:ARG:HG2	1:A:488:TYR:HB3	1.95	0.49
1:A:443:LEU:HD22	1:A:577:LEU:HD21	1.95	0.49
1:A:583:GLU:O	1:A:587:LYS:HG3	2.12	0.49
1:A:246:PHE:C	1:A:246:PHE:HD1	2.17	0.48
1:A:63:GLU:O	1:A:209:ASN:OD1	2.31	0.48
1:A:151:SER:HG	1:A:153:LEU:H	1.59	0.48
1:A:268:LEU:HD12	1:A:268:LEU:C	2.33	0.48
1:A:102:THR:HG22	1:A:103:VAL:HG13	1.96	0.47
1:A:293:LEU:CD1	1:A:296:MET:HE3	2.45	0.47
1:A:178:THR:N	1:A:179:PRO:CD	2.77	0.47
1:A:246:PHE:C	1:A:246:PHE:CD1	2.87	0.46
1:A:19:SER:O	1:A:20:LEU:C	2.54	0.46
1:A:434:GLY:O	1:A:460:MET:HE3	2.15	0.46
1:A:69:LEU:HD11	1:A:77:LEU:HB2	1.99	0.44
1:A:221:HIS:C	1:A:222:LEU:HD12	2.38	0.44
1:A:368:VAL:CG2	5:A:705:EDO:H21	2.48	0.44
1:A:89:GLU:HG3	6:A:792:HOH:O	2.18	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:ASN:OD1	1:A:272:ASN:C	2.57	0.43
1:A:156:ILE:HD13	1:A:184:LEU:HB2	2.00	0.43
1:A:450:TRP:HA	5:A:704:EDO:H22	2.01	0.43
1:A:248:LEU:HD23	1:A:248:LEU:HA	1.89	0.43
1:A:143:PRO:HA	1:A:158:ILE:HD13	2.00	0.43
1:A:142:ILE:HD11	1:A:167:VAL:HG22	2.01	0.42
1:A:242:GLY:HA2	1:A:246:PHE:HB3	2.01	0.42
1:A:151:SER:OG	1:A:152:PRO:N	2.52	0.42
1:A:315:MET:HA	1:A:324:ARG:O	2.19	0.42
1:A:200:LEU:HB3	1:A:201:PRO:HA	2.00	0.42
1:A:527:TRP:HA	1:A:528:PRO:HD3	1.94	0.42
1:A:87:ARG:HG2	1:A:87:ARG:NH1	2.35	0.42
1:A:582:LYS:HB2	1:A:622:MET:HG3	2.00	0.42
1:A:261:LEU:HD11	1:A:500:MET:SD	2.60	0.42
1:A:391:TRP:CE2	1:A:399:CYS:HB3	2.54	0.41
1:A:503:HIS:HA	1:A:504:PRO:HD3	1.91	0.41
1:A:217:THR:HA	1:A:231:PHE:O	2.20	0.41
1:A:336:ALA:HB3	1:A:337:PRO:HD3	2.02	0.41
1:A:413:CYS:HB3	1:A:416:ILE:HB	2.03	0.41
1:A:604:LEU:HA	1:A:605:PRO:HD3	1.96	0.41
1:A:278:LYS:HG3	1:A:541:PHE:CE1	2.56	0.41
1:A:473:LYS:N	1:A:473:LYS:CD	2.76	0.41
1:A:249:ARG:O	1:A:487:GLY:HA3	2.21	0.41
1:A:73:ASP:C	1:A:73:ASP:OD1	2.58	0.40
1:A:293:LEU:HD12	1:A:296:MET:CE	2.51	0.40
1:A:441:VAL:HA	1:A:442:PRO:HD3	1.88	0.40
1:A:221:HIS:CE1	1:A:249:ARG:HD2	2.56	0.40
1:A:73:ASP:HB3	1:A:353:LYS:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	597/625 (96%)	580 (97%)	16 (3%)	1 (0%)	52 47

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	236	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	508/523 (97%)	489 (96%)	19 (4%)	41 37

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	LEU
1	A	100	ARG
1	A	123	GLU
1	A	226	ARG
1	A	246	PHE
1	A	253	ARG
1	A	290	LYS
1	A	353	LYS
1	A	387	LYS
1	A	392	SER
1	A	393	SER
1	A	395	ASP
1	A	473	LYS
1	A	493	TYR
1	A	521	LYS
1	A	560	LYS
1	A	575	MET
1	A	581	SER
1	A	616	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	292	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	PEP	A	703	3	5,9,9	2.81	2 (40%)	8,13,13	2.98	3 (37%)
5	EDO	A	704	-	3,3,3	0.50	0	2,2,2	0.83	0
5	EDO	A	705	-	3,3,3	0.43	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEP	A	703	3	-	0/5/9/9	0/0/0/0
5	EDO	A	704	-	-	0/1/1/1	0/0/0/0
5	EDO	A	705	-	-	0/1/1/1	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	703	PEP	P-O2P	3.82	1.68	1.54
4	A	703	PEP	P-O3P	4.77	1.71	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	703	PEP	O3P-P-O2P	-2.25	98.82	107.38
4	A	703	PEP	O3P-P-O2	2.26	112.64	105.25
4	A	703	PEP	C1-C2-C3	7.68	134.56	120.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	704	EDO	1	0
5	A	705	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	603/625 (96%)	-0.21	10 (1%) 73 74	4, 12, 30, 47	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	393	SER	4.0
1	A	152	PRO	3.8
1	A	394	GLU	3.6
1	A	379	SER	3.2
1	A	10	ASN	3.2
1	A	396	GLY	2.6
1	A	473	LYS	2.5
1	A	378	ALA	2.2
1	A	432	ILE	2.1
1	A	21	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	EDO	A	704	4/4	0.97	0.16	1.52	15,18,23,25	0
5	EDO	A	705	4/4	0.97	0.08	-0.61	17,19,19,21	0
4	PEP	A	703	10/10	0.97	0.09	-0.78	12,15,18,19	0
3	NA	A	702	1/1	0.98	0.10	-	20,20,20,20	0
2	MN	A	701	1/1	1.00	0.06	-	10,10,10,10	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.