



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 03:20 PM BST

PDB ID : 1KSM
Title : AVERAGE NMR SOLUTION STRUCTURE OF CA LN CALBINDIN D9K
Authors : Bertini, I.; Donaire, A.; Luchinat, C.; Piccioli, M.; Poggi, L.; Parigi, G.; Jimenez, B.
Deposited on : 2002-01-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	79	

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1203 atoms, of which 602 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called VITAMIN D-DEPENDENT CALCIUM-BINDING PROTEIN.

Mol	Chain	Residues	Atoms						Trace
1	A	75	Total	C	H	N	O	S	0
			1202	384	602	91	124	1	

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	43	MET	PRO	ENGINEERED	UNP P02633


- Molecule 2 is LANTHANUM (III) ION (three-letter code: LA) (formula: La).

Mol	Chain	Residues	Atoms	
2	A	1	Total	La
			1	1

4 Residue-property plots

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: VITAMIN D-DEPENDENT CALCIUM-BINDING PROTEIN



5 Refinement protocol and experimental data overview

The models were refined using the following method: *SIMULATED ANNEALING TORSION ANGLE DYNAMICS*.

Of the 30 calculated structures, 1 were deposited, based on the following criterion: *minimized average structure*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	1.5
PSEUDYANA	structure solution	1.5
PSEUDYANA	refinement	3.1

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section:
LA

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	600	602	604	27
All	All	601	602	604	27

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:49:LEU:HD23	1:A:53:LEU:HD11	0.91	1.42
1:A:61:VAL:O	1:A:61:VAL:HG12	0.70	1.86
1:A:23:LEU:HD12	1:A:28:LEU:HD13	0.70	1.62
1:A:49:LEU:HD23	1:A:53:LEU:CD1	0.55	2.26
1:A:31:LEU:O	1:A:35:GLU:N	0.52	2.42
1:A:10:PHE:CE2	1:A:23:LEU:HD21	0.51	2.41
1:A:28:LEU:CD2	1:A:61:VAL:HG11	0.51	2.35
1:A:30:LEU:O	1:A:33:GLN:CB	0.50	2.60
1:A:61:VAL:CG1	1:A:61:VAL:O	0.49	2.58
1:A:36:PHE:CE2	1:A:73:ILE:HG21	0.49	2.42
1:A:10:PHE:CZ	1:A:23:LEU:HD21	0.48	2.43
1:A:35:GLU:HG3	1:A:36:PHE:CD2	0.48	2.44
1:A:25:LYS:HB2	1:A:50:PHE:CD1	0.47	2.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:6:LEU:O	1:A:10:PHE:CB	0.47	2.62
1:A:13:TYR:CE2	1:A:31:LEU:HA	0.45	2.47
1:A:32:LEU:O	1:A:36:PHE:O	0.44	2.35
1:A:32:LEU:O	1:A:36:PHE:C	0.44	2.55
1:A:10:PHE:CD1	1:A:11:GLU:N	0.43	2.87
1:A:17:GLU:O	1:A:27:GLU:CD	0.43	2.57
1:A:41:LYS:O	1:A:42:GLY:O	0.43	2.37
1:A:59:GLY:C	1:A:60:GLU:OE1	0.42	2.58
1:A:25:LYS:CG	1:A:46:LEU:HD12	0.42	2.44
1:A:13:TYR:O	1:A:27:GLU:OE2	0.42	2.38
1:A:31:LEU:C	1:A:33:GLN:N	0.41	2.72
1:A:22:GLN:CD	1:A:60:GLU:OE2	0.41	2.59
1:A:17:GLU:O	1:A:27:GLU:OE2	0.40	2.39
1:A:13:TYR:CG	1:A:31:LEU:HD12	0.40	2.51

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	73/79 (92%)	61 (84%)	11 (15%)	1 (1%)	19	64
All	All	73/79 (92%)	61 (84%)	11 (15%)	1 (1%)	19	64

All 1 Ramachandran outliers are listed below.

Mol	Chain	Res	Type
1	A	42	GLY

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	68/71 (96%)	45 (66%)	23 (34%)	1	11
All	All	68/71 (96%)	45 (66%)	23 (34%)	1	11

All 23 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	74	SER
1	A	4	GLU
1	A	26	GLU
1	A	64	GLU
1	A	58	ASP
1	A	23	LEU
1	A	75	GLN
1	A	62	SER
1	A	38	SER
1	A	11	GLU
1	A	56	ASN
1	A	7	LYS
1	A	1	LYS
1	A	49	LEU
1	A	2	SER
1	A	5	GLU
1	A	46	LEU
1	A	39	LEU
1	A	34	THR
1	A	51	GLU
1	A	31	LEU
1	A	47	ASP
1	A	35	GLU

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided