



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:05 AM GMT

PDB ID : 3KSQ  
Title : Discovery of C-Imidazole Azaheptapyridine FPT Inhibitors  
Authors : Zhu, H.Y.; Cooper, A.B.; Njoroge, G.; Kirschmeier, P.; Strickland, C.; Hruza, A.; Girijavallabhan, V.M.  
Deposited on : 2009-11-23  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	<div> <div>2%</div> <div>77%</div> <div>6%</div> <div>16%</div> </div>
2	B	437	<div> <div>4%</div> <div>85%</div> <div>7%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	B	1001	-	-	-	X
5	FPP	B	438	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6211 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Farnesyltransferase, CAAX box, alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	0	0	0
			2680	1706	470	499	5			

- Molecule 2 is a protein called Protein farnesyltransferase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	403	Total	C	N	O	S	0	0	0
			3169	2026	545	575	23			


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		

- Molecule 5 is FARNESYL DIPHOSPHATE (three-letter code: FPP) (formula: C<sub>15</sub>H<sub>28</sub>O<sub>7</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	O	P	0	0
			24	15	7	2		

- Molecule 6 is TERT-BUTYL 4-[(11S)-8-CHLORO-6-[(R)-HYDROXY(1-METHYL-1H-IMIDAZOL-5-YL)METHYL]-11H-BENZO[5,6]CYCLOHEPTA[1,2-B]PYRIDIN-11-YL]PIPERAZINE-1-CARBOXYLATE (three-letter code: Z96) (formula:  $C_{28}H_{32}ClN_5O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	Cl	N	O	0
			37	28	1	5	3	


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	142	Total 142	O 142	0	0
7	B	157	Total 157	O 157	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Farnesyltransferase, CAAX box, alpha



- Molecule 2: Protein farnesyltransferase subunit beta



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	171.12Å 171.12Å 69.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.64 – 2.10 29.64 – 2.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (29.64-2.10) 99.5 (29.64-2.10)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.46 (at 2.10Å)	Xtriage
Refinement program	BUSTER 2.9.2	Depositor
R, $R_{free}$	0.193 , 0.219 0.188 , 0.209	Depositor DCC
$R_{free}$ test set	3413 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	35.6	Xtriage
Anisotropy	0.209	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 47.4	EDS
Estimated twinning fraction	0.023 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 67705 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6211	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: Z96, MG, ZN, FPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/2746	0.60	0/3728
2	B	0.52	0/3255	0.64	0/4421
All	All	0.51	0/6001	0.62	0/8149

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2680	0	2600	10	0
2	B	3169	0	3098	14	0
3	B	1	0	0	0	0
4	B	1	0	0	0	0
5	B	24	0	25	1	0
6	B	37	0	32	1	0
7	A	142	0	0	0	0
7	B	157	0	0	0	0
All	All	6211	0	5755	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:414:HIS:O	2:B:417:GLN:HG3	1.90	0.70
2:B:21:GLU:HB2	2:B:22:PRO:HD2	1.74	0.69
2:B:21:GLU:HB2	2:B:22:PRO:CD	2.23	0.68
1:A:342:GLU:HG2	1:A:346:LYS:HD2	1.86	0.57
1:A:287:ARG:HB3	1:A:291:ARG:HD3	1.88	0.55
2:B:301:SER:O	2:B:305:ALA:HB3	2.06	0.55
1:A:326:GLN:HA	1:A:330:LYS:HE3	1.87	0.55
1:A:77:PRO:HB2	1:A:101:ASP:HB3	1.89	0.55
2:B:192:LEU:HD23	2:B:199:VAL:HG23	1.89	0.55
2:B:253:PHE:HA	2:B:307:LEU:HD21	1.92	0.50
1:A:300:LEU:HD12	1:A:303:GLN:HG3	1.96	0.47
2:B:239:ILE:HB	2:B:252:THR:HA	1.97	0.47
1:A:67:ARG:HH22	1:A:94:GLU:CD	2.17	0.46
2:B:268:LEU:HD13	2:B:273:LEU:HD12	1.98	0.46
2:B:201:VAL:HG23	2:B:241:GLY:O	2.17	0.45
2:B:75:ARG:NH2	2:B:393:VAL:O	2.51	0.44
1:A:287:ARG:HB3	1:A:291:ARG:CD	2.48	0.44
1:A:75:ILE:HD11	1:A:109:ARG:HD2	2.00	0.43
1:A:164:LYS:O	6:B:439:Z96:CL	2.74	0.43
2:B:348:GLY:O	2:B:358:ARG:HD2	2.19	0.42
2:B:202:ARG:HD2	5:B:438:FPP:H152	2.01	0.42
2:B:350:LEU:HB2	2:B:363:THR:HA	2.02	0.41
2:B:23:LEU:HD12	2:B:26:LEU:HD22	2.03	0.41
1:A:55:PHE:CE2	1:A:121:ARG:HD3	2.56	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/377 (83%)	297 (95%)	14 (4%)	2 (1%)	30	24
2	B	401/437 (92%)	396 (99%)	5 (1%)	0	100	100
All	All	714/814 (88%)	693 (97%)	19 (3%)	2 (0%)	46	45

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	286	ASP
1	A	304	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/338 (87%)	281 (96%)	12 (4%)	37	36
2	B	340/371 (92%)	331 (97%)	9 (3%)	54	58
All	All	633/709 (89%)	612 (97%)	21 (3%)	45	47

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75	ILE
1	A	76	ASP
1	A	85	SER
1	A	94	GLU
1	A	285	GLN
1	A	286	ASP
1	A	297	ASN
1	A	300	LEU
1	A	323	LEU
1	A	330	LYS
1	A	363	LEU
1	A	368	SER
2	B	21	GLU
2	B	83	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	84	ARG
2	B	108	LEU
2	B	198	GLU
2	B	266	ARG
2	B	351	LEU
2	B	352	ASP
2	B	417	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	285	GLN
2	B	275	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	FPP	B	438	-	21,23,23	1.55	4 (19%)	27,31,31	1.66	7 (25%)
6	Z96	B	439	4	36,41,41	2.64	14 (38%)	46,61,61	1.71	10 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	FPP	B	438	-	-	0/25/25/25	0/0/0/0
6	Z96	B	439	4	-	0/13/47/47	0/5/5/5

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	438	FPP	O1-C1	-3.17	1.39	1.43
5	B	438	FPP	PA-O2A	2.23	1.64	1.54
6	B	439	Z96	C5-C10	2.36	1.44	1.37
6	B	439	Z96	C5-C4	2.50	1.44	1.38
5	B	438	FPP	PA-O1A	2.65	1.60	1.51
6	B	439	Z96	C2-C6	2.67	1.54	1.51
6	B	439	Z96	C1-C13	2.69	1.43	1.38
6	B	439	Z96	C28-N3	2.74	1.39	1.34
6	B	439	Z96	C15-C14	2.89	1.43	1.38
6	B	439	Z96	C3-C12	2.96	1.51	1.46
5	B	438	FPP	PB-O1B	3.47	1.62	1.51
6	B	439	Z96	C8-C9	3.93	1.53	1.49
6	B	439	Z96	C13-C14	4.09	1.46	1.38
6	B	439	Z96	C1-C7	4.25	1.45	1.39
6	B	439	Z96	C11-C9	4.28	1.56	1.51
6	B	439	Z96	C2-N4	4.52	1.39	1.34
6	B	439	Z96	C3-C2	5.12	1.49	1.41
6	B	439	Z96	C8-C7	7.41	1.49	1.40

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	438	FPP	PA-O3A-PB	-3.69	120.30	132.67
6	B	439	Z96	N3-C28-N2	-3.33	107.78	112.28
5	B	438	FPP	O3B-PB-O1B	-3.11	100.58	110.58
6	B	439	Z96	O1-C25-N1	-3.05	118.51	124.32
6	B	439	Z96	C15-C14-CL	-2.81	115.66	119.14

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	438	FPP	O1-PA-O1A	-2.03	101.76	109.62
6	B	439	Z96	C11-C9-C12	2.07	123.96	119.58
6	B	439	Z96	C3-C2-C6	2.17	122.71	120.02
6	B	439	Z96	C4-C3-C12	2.18	121.22	117.62
5	B	438	FPP	C6-C5-C3	2.25	120.03	112.71
5	B	438	FPP	O2A-PA-O3A	2.32	115.62	105.09
6	B	439	Z96	C21-N3-C28	2.89	110.27	105.71
6	B	439	Z96	C23-N2-C27	3.14	128.22	124.35
5	B	438	FPP	O3A-PA-O1	3.19	111.39	102.94
5	B	438	FPP	C10-C8-C9	3.41	120.61	115.41
6	B	439	Z96	O2-C25-N1	3.69	116.66	110.96
6	B	439	Z96	C29-O2-C25	4.89	126.62	121.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	438	FPP	1	0
6	B	439	Z96	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	315/377 (83%)	-0.06	9 (2%) 55 63	23, 42, 73, 100	0
2	B	403/437 (92%)	0.09	17 (4%) 40 49	24, 37, 61, 96	0
All	All	718/814 (88%)	0.02	26 (3%) 46 55	23, 40, 68, 100	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	55	PHE	7.3
2	B	64	PHE	6.7
2	B	423	PHE	5.3
1	A	304	PRO	3.9
1	A	84	PRO	3.4
2	B	319	GLY	3.1
2	B	355	GLY	3.0
1	A	367	HIS	2.9
1	A	305	SER	2.9
2	B	422	GLY	2.7
1	A	54	GLY	2.7
1	A	368	SER	2.5
2	B	22	PRO	2.5
2	B	21	GLU	2.4
1	A	306	HIS	2.3
2	B	380	ALA	2.3
2	B	301	SER	2.3
2	B	417	GLN	2.3
2	B	288	PHE	2.3
2	B	390	PRO	2.2
2	B	70	ARG	2.1
2	B	28	PRO	2.1
1	A	286	ASP	2.1
2	B	389	VAL	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	277	VAL	2.0
2	B	419	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	MG	B	1001	1/1	0.99	0.29	7.06	27,27,27,27	0
5	FPP	B	438	24/24	0.84	0.21	2.62	33,58,91,92	0
6	Z96	B	439	37/37	0.91	0.14	0.54	45,53,60,64	0
4	ZN	B	2001	1/1	1.00	0.03	-2.53	38,38,38,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.