



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:15 PM GMT

PDB ID : 4L1F  
Title : Electron transferring flavoprotein of Acidaminococcus fermentans: Towards a mechanism of flavin-based electron bifurcation  
Authors : Mowafy, A.M.; Chowdhury, N.P.; Demmer, J.; Upadhyay, V.; Kolzer, S.; Jayamani, E.; Kahnt, J.; Demmer, U.; Ermler, U.; Buckel, W.  
Deposited on : 2013-06-03  
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	383	<div> <div style="width: 90%;"></div> <div style="width: 8%;"></div> <div style="width: 2%;"></div> <div style="width: 1%;"></div> <div style="width: 1%;"></div> </div> <div>90% 8% ..</div>
1	B	383	<div> <div style="width: 90%;"></div> <div style="width: 8%;"></div> <div style="width: 2%;"></div> <div style="width: 1%;"></div> <div style="width: 1%;"></div> </div> <div>90% 8% ..</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	COS	A	402	-	-	-	X
3	COS	B	402	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6653 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-CoA dehydrogenase domain protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	380	Total	C	N	O	S	0	1	0
			2910	1843	477	566	24			
1	B	380	Total	C	N	O	S	0	1	0
			2910	1843	477	566	24			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is COENZYME A PERSULFIDE (three-letter code: COS) (formula:  $C_{21}H_{36}N_7O_{16}P_3S_2$ ).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			49	21	7	16	3	2		
3	B	1	Total	C	N	O	P	S	0	0
			49	21	7	16	3	2		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		
4	A	2	Total	Na	0	0
			2	2		

- Molecule 5 is 1,3-PROPANDIOL (three-letter code: PDO) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			5	3	2		

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula:  $O_4P$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	313	Total 313	O 313	0	0
7	B	303	Total 303	O 303	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acyl-CoA dehydrogenase domain protein



- Molecule 1: Acyl-CoA dehydrogenase domain protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	109.10 Å   141.40 Å   64.10 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	47.13 – 1.79 47.13 – 1.79	Depositor EDS
% Data completeness (in resolution range)	99.3 (47.13-1.79) 99.3 (47.13-1.79)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.04 (at 1.79 Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.153   ,   0.183 0.165   ,   0.193	Depositor DCC
$R_{free}$ test set	4666 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.7	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 93074 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6653	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PDO, NA, COS, PO4, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.11	4/2959 (0.1%)	1.09	12/3980 (0.3%)
1	B	1.08	3/2959 (0.1%)	1.06	8/3980 (0.2%)
All	All	1.10	7/5918 (0.1%)	1.07	20/7960 (0.3%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	166	TYR	CG-CD1	6.75	1.48	1.39
1	A	274	SER	CB-OG	-6.60	1.33	1.42
1	B	274	SER	CB-OG	-6.20	1.34	1.42
1	B	75	GLU	CD-OE1	-6.16	1.18	1.25
1	B	105	GLU	CD-OE1	5.57	1.31	1.25
1	A	104	GLU	CD-OE2	5.11	1.31	1.25
1	A	88	SER	CA-CB	5.02	1.60	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	273	ARG	NE-CZ-NH2	-10.30	115.15	120.30
1	B	325	ASP	CB-CG-OD1	7.29	124.86	118.30
1	B	351	ARG	NE-CZ-NH1	7.15	123.88	120.30
1	A	174	LYS	CD-CE-NZ	6.83	127.41	111.70
1	B	171	MET	CG-SD-CE	-6.80	89.32	100.20
1	B	362	TYR	CB-CG-CD2	-6.62	117.03	121.00
1	A	14	ASP	CB-CG-OD1	6.60	124.24	118.30
1	A	351	ARG	NE-CZ-NH1	6.24	123.42	120.30
1	A	31	ARG	NE-CZ-NH1	6.19	123.39	120.30
1	A	362	TYR	CB-CG-CD1	6.10	124.66	121.00
1	B	370	ARG	NE-CZ-NH1	6.00	123.30	120.30
1	A	362	TYR	CB-CG-CD2	-5.78	117.53	121.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	109	LYS	CD-CE-NZ	-5.67	98.67	111.70
1	B	362	TYR	CB-CG-CD1	5.40	124.24	121.00
1	A	354	ARG	NE-CZ-NH2	5.38	122.99	120.30
1	A	274	SER	N-CA-CB	5.36	118.54	110.50
1	A	273	ARG	NE-CZ-NH1	5.22	122.91	120.30
1	B	362	TYR	CA-CB-CG	5.12	123.14	113.40
1	B	273	ARG	NE-CZ-NH2	-5.10	117.75	120.30
1	A	33	LYS	CD-CE-NZ	5.06	123.33	111.70

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2910	0	2879	16	0
1	B	2910	0	2879	19	0
2	A	53	0	31	3	0
2	B	53	0	31	5	0
3	A	49	0	32	1	0
3	B	49	0	32	0	0
4	A	2	0	0	0	0
4	B	1	0	0	0	0
5	A	5	0	8	1	0
6	B	5	0	0	0	0
7	A	313	0	0	3	0
7	B	303	0	0	5	0
All	All	6653	0	5892	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:10:GLN:NE2	7:B:621:HOH:O	2.14	0.72
1:B:156:PHE:O	2:B:401:FAD:C4X	2.38	0.70
1:A:358:ILE:HD11	2:A:401:FAD:HM83	1.80	0.63
1:A:358:ILE:CD1	2:A:401:FAD:HM83	2.30	0.62
1:A:197:LYS:HE2	7:A:799:HOH:O	2.00	0.60
1:A:200:ASP:OD1	7:A:561:HOH:O	2.16	0.60
1:B:358:ILE:HD11	2:B:401:FAD:HM83	1.86	0.56
1:B:306:LYS:NZ	7:B:767:HOH:O	2.39	0.56
1:B:13:LYS:HE2	7:B:737:HOH:O	2.09	0.52
1:A:156:PHE:O	2:A:401:FAD:C4X	2.58	0.52
1:B:379:ARG:NE	7:B:729:HOH:O	2.43	0.51
1:A:13:LYS:HE2	7:A:730:HOH:O	2.11	0.50
1:A:5:LEU:O	5:A:404:PDO:H21	2.11	0.50
1:A:247:ALA:HB1	1:A:299:VAL:HG22	1.94	0.49
1:B:156:PHE:HB3	2:B:401:FAD:C9A	2.43	0.49
1:A:343:TYR:CZ	1:B:358:ILE:HB	2.47	0.49
1:B:351:ARG:NH2	7:B:760:HOH:O	2.48	0.47
1:B:358:ILE:CD1	2:B:401:FAD:HM83	2.45	0.46
1:A:70:ILE:HG23	1:A:248:ALA:HB2	1.97	0.46
1:B:156:PHE:HB3	2:B:401:FAD:C9	2.46	0.46
1:B:74:GLU:OE2	1:B:255:GLU:OE1	2.34	0.46
1:A:363:GLU:N	3:A:402:COS:S'P	2.76	0.45
1:B:93:LEU:HB3	1:B:123:PHE:HB2	1.98	0.44
1:B:147:LYS:HD2	1:B:218:GLU:HB3	1.98	0.44
1:B:124:GLY:HA2	1:B:157:ILE:HD12	1.99	0.44
1:A:93:LEU:HB3	1:A:123:PHE:HB2	1.99	0.44
1:A:132:THR:HG23	1:A:368:VAL:CG2	2.48	0.43
1:B:59:TYR:O	1:B:112:ALA:HA	2.19	0.42
1:A:82:GLY:HA3	1:A:206:THR:OG1	2.19	0.42
1:B:132:THR:HG23	1:B:368:VAL:CG2	2.50	0.42
1:B:247:ALA:HB1	1:B:299:VAL:HG22	2.02	0.42
1:A:379:ARG:NH1	1:A:379:ARG:HB3	2.34	0.42
1:A:14:ASP:O	1:A:18:GLU:HG3	2.20	0.41
1:B:158:THR:O	1:B:159:ASN:HB2	2.20	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/383 (99%)	371 (98%)	8 (2%)	0	100	100
1	B	379/383 (99%)	368 (97%)	11 (3%)	0	100	100
All	All	758/766 (99%)	739 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/300 (99%)	294 (99%)	4 (1%)	76	68
1	B	298/300 (99%)	293 (98%)	5 (2%)	68	57
All	All	596/600 (99%)	587 (98%)	9 (2%)	72	62

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	221	LYS
1	A	273	ARG
1	A	362	TYR
1	A	379	ARG
1	B	35	HIS
1	B	40	LYS
1	B	144	LYS
1	B	362	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	379	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	179	HIS
1	A	205	HIS
1	B	179	HIS
1	B	205	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	FAD	A	401	-	48,58,58	1.55	6 (12%)	54,89,89	2.98	22 (40%)
3	COS	A	402	-	40,51,51	0.95	1 (2%)	52,76,76	2.39	16 (30%)
5	PDO	A	404	-	4,4,4	0.50	0	3,3,3	0.26	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	B	401	-	48,58,58	1.49	6 (12%)	54,89,89	2.61	15 (27%)
3	COS	B	402	-	40,51,51	1.30	5 (12%)	52,76,76	2.16	15 (28%)
6	PO4	B	403	-	4,4,4	0.45	0	6,6,6	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	401	-	-	0/30/50/50	0/6/6/6
3	COS	A	402	-	-	0/44/65/65	0/3/3/3
5	PDO	A	404	-	-	0/2/2/2	0/0/0/0
2	FAD	B	401	-	-	0/30/50/50	0/6/6/6
3	COS	B	402	-	-	0/44/65/65	0/3/3/3
6	PO4	B	403	-	-	0/0/0/0	0/0/0/0

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	FAD	C6-C5X	-5.85	1.32	1.41
2	B	401	FAD	C6-C5X	-2.98	1.37	1.41
3	B	402	COS	C2A-N3A	2.01	1.35	1.32
3	B	402	COS	C2A-N1A	2.06	1.37	1.33
2	B	401	FAD	C4X-N5	2.31	1.37	1.33
3	B	402	COS	C8A-N7A	2.48	1.39	1.34
2	A	401	FAD	C4-C4X	2.79	1.46	1.41
2	B	401	FAD	C9A-C5X	2.81	1.48	1.42
2	A	401	FAD	C4X-N5	2.86	1.37	1.33
3	A	402	COS	C5A-C4A	3.00	1.47	1.40
2	A	401	FAD	C4X-C10	3.04	1.46	1.41
2	A	401	FAD	O4B-C1B	3.16	1.45	1.41
2	B	401	FAD	O4B-C1B	3.40	1.45	1.41
3	B	402	COS	C5A-C4A	3.54	1.48	1.40
2	B	401	FAD	C4-C4X	3.60	1.48	1.41
2	A	401	FAD	C5'-C4'	3.64	1.57	1.51
3	B	402	COS	O4B-C1B	3.68	1.45	1.41
2	B	401	FAD	C4X-C10	4.17	1.48	1.41

All (68) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	COS	N3A-C2A-N1A	-8.84	122.13	128.89
2	A	401	FAD	C4X-C10-N10	-8.40	115.57	120.52
3	B	402	COS	N3A-C2A-N1A	-7.79	122.93	128.89
2	B	401	FAD	N3A-C2A-N1A	-7.75	122.96	128.89
2	A	401	FAD	C4-C4X-C10	-6.38	115.86	119.94
2	B	401	FAD	C4-C4X-C10	-6.00	116.10	119.94
3	B	402	COS	CEP-CBP-CCP	-5.90	100.85	108.50
2	A	401	FAD	N3A-C2A-N1A	-5.79	124.46	128.89
3	A	402	COS	CEP-CBP-CCP	-5.68	101.14	108.50
2	B	401	FAD	C4X-C10-N10	-5.15	117.49	120.52
3	A	402	COS	C1B-N9A-C4A	-4.44	120.25	126.94
2	A	401	FAD	C4X-C4-N3	-4.36	117.63	123.59
2	B	401	FAD	C4X-C4-N3	-4.26	117.76	123.59
2	A	401	FAD	O4'-C4'-C5'	-4.24	100.96	110.19
3	A	402	COS	CDP-CBP-CCP	-4.20	103.05	108.50
3	A	402	COS	O3B-P3B-O7A	-4.11	96.84	107.11
3	B	402	COS	C1B-N9A-C4A	-4.06	120.82	126.94
2	A	401	FAD	P-O3P-PA	-4.05	121.35	132.73
3	B	402	COS	C2B-C1B-N9A	-3.79	108.51	114.29
3	A	402	COS	O5P-C5P-C6P	-3.54	115.88	121.98
2	A	401	FAD	C9A-C5X-N5	-3.46	117.24	122.36
2	B	401	FAD	C2B-C1B-N9A	-3.37	109.15	114.29
3	B	402	COS	P2A-O3A-P1A	-3.18	123.79	132.73
3	A	402	COS	C2B-C1B-N9A	-3.10	109.55	114.29
2	A	401	FAD	O3P-PA-O5B	-3.01	94.95	102.94
3	A	402	COS	P2A-O3A-P1A	-2.99	124.33	132.73
3	A	402	COS	C6P-C7P-N8P	-2.86	105.61	111.88
2	A	401	FAD	C9-C9A-C5X	-2.83	114.58	119.62
3	B	402	COS	O3B-P3B-O7A	-2.83	100.05	107.11
2	A	401	FAD	O3'-C3'-C2'	-2.63	102.13	108.75
3	B	402	COS	C3P-N4P-C5P	-2.59	117.70	122.79
2	B	401	FAD	C9A-C5X-N5	-2.58	118.55	122.36
2	A	401	FAD	O3B-C3B-C2B	-2.50	103.71	111.83
3	B	402	COS	C6P-C7P-N8P	-2.47	106.46	111.88
2	B	401	FAD	P-O3P-PA	-2.40	125.98	132.73
2	A	401	FAD	C4A-C5A-N7A	-2.40	107.27	109.48
2	A	401	FAD	C1B-N9A-C4A	-2.40	123.32	126.94
2	B	401	FAD	O3'-C3'-C2'	-2.29	102.97	108.75
2	B	401	FAD	C1B-N9A-C4A	-2.20	123.62	126.94
3	A	402	COS	C4A-C5A-N7A	-2.20	107.45	109.48
3	A	402	COS	O3B-C3B-C4B	-2.19	101.39	109.99
2	B	401	FAD	C4X-N5-C5X	2.03	119.09	116.76
2	A	401	FAD	C4X-N5-C5X	2.13	119.21	116.76

Continued on next page...



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	COS	CDP-CBP-CAP	2.13	113.23	109.34
2	A	401	FAD	O2A-PA-O1A	2.13	124.07	112.53
3	B	402	COS	N6A-C6A-N1A	2.20	123.92	119.20
2	A	401	FAD	O4B-C4B-C3B	2.25	109.69	105.15
3	B	402	COS	O9A-P3B-O8A	2.26	115.97	107.38
3	B	402	COS	P3B-O3B-C3B	2.27	127.00	121.56
3	B	402	COS	C2A-N1A-C6A	2.28	122.85	118.77
2	A	401	FAD	O2P-P-O1P	2.29	124.94	112.53
2	A	401	FAD	C1'-N10-C9A	2.30	121.44	118.86
3	A	402	COS	O2A-P1A-O1A	2.31	125.06	112.53
2	B	401	FAD	O2'-C2'-C1'	2.41	115.88	109.94
2	B	401	FAD	N6A-C6A-N1A	2.65	124.89	119.20
2	A	401	FAD	O2'-C2'-C1'	2.84	116.93	109.94
3	A	402	COS	C2A-N1A-C6A	2.85	123.86	118.77
3	A	402	COS	CEP-CBP-CDP	3.13	115.56	109.28
3	B	402	COS	O4B-C1B-N9A	3.29	114.98	108.10
3	A	402	COS	CEP-CBP-CAP	3.40	115.56	109.34
3	A	402	COS	O9A-P3B-O7A	3.47	121.74	110.58
3	B	402	COS	CEP-CBP-CAP	3.54	115.81	109.34
2	B	401	FAD	C2A-N1A-C6A	3.55	125.11	118.77
2	A	401	FAD	C6-C5X-C9A	3.96	124.20	118.98
2	B	401	FAD	C5X-C9A-N10	6.50	122.56	117.62
2	A	401	FAD	C5X-C9A-N10	7.84	123.57	117.62
2	B	401	FAD	C4-N3-C2	9.43	123.40	115.25
2	A	401	FAD	C4-N3-C2	9.53	123.49	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	FAD	3	0
3	A	402	COS	1	0
5	A	404	PDO	1	0
2	B	401	FAD	5	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	380/383 (99%)	-0.21	0 100 100	7, 15, 29, 50	0
1	B	380/383 (99%)	-0.15	2 (0%) 91 90	8, 16, 32, 52	0
All	All	760/766 (99%)	-0.18	2 (0%) 94 92	7, 15, 30, 52	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	379	ARG	3.2
1	B	146	ASP	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	COS	A	402	49/49	0.90	0.20	4.84	21,30,47,59	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	COS	B	402	49/49	0.91	0.20	4.54	20,31,50,52	0
4	NA	A	403	1/1	0.96	0.09	1.96	31,31,31,31	0
4	NA	A	405	1/1	0.98	0.10	1.69	31,31,31,31	0
6	PO4	B	403	5/5	0.92	0.13	1.22	52,60,65,70	0
4	NA	B	404	1/1	0.92	0.09	0.54	42,42,42,42	0
2	FAD	B	401	53/53	0.97	0.07	-0.45	9,12,18,21	0
2	FAD	A	401	53/53	0.97	0.07	-0.45	9,13,18,20	0
5	PDO	A	404	5/5	0.69	0.35	-	44,47,48,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.