



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 12:53 AM BST

PDB ID : 2L2K  
Title : Solution NMR structure of the R/G STEM LOOP RNA-ADAR2 DSRBM2 Complex  
Authors : Allain, F.H.-T.; Oberstrass, F.C.; Steff, R.  
Deposited on : 2010-08-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*



The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958
RNA backbone	3027	600

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	B	71	 89% 7%
2	A	42	 98%

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	B:48-B:113 (66)	0.19	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 6 single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 5, 6, 12, 15, 19
2	10, 17
3	7, 14
4	13, 16
Single-model clusters	2; 8; 9; 11; 18; 20

### 3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2438 atoms, of which 1006 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Adenosine deaminase.

Mol	Chain	Residues	Atoms						Trace
1	B	71	Total	C	H	N	O	S	0
			1093	344	551	96	100	2	

- Molecule 2 is a RNA chain called RNA (42-MER).


Mol	Chain	Residues	Atoms						Trace
2	A	42	Total	C	H	N	O	P	0
			1345	398	455	157	294	41	

## 4 Residue-property plots [i](#)


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Adenosine deaminase



- Molecule 2: RNA (42-MER)




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: Adenosine deaminase




- Molecule 2: RNA (42-MER)



### 4.2.2 Score per residue for model 2

- Molecule 1: Adenosine deaminase

Chain B:  85% 8% 7%




- Molecule 2: RNA (42-MER)

Chain A:  95% 5%



### 4.2.3 Score per residue for model 3

- Molecule 1: Adenosine deaminase

Chain B:  83% 10% 7%




- Molecule 2: RNA (42-MER)

Chain A:  93% 5%



### 4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: Adenosine deaminase

Chain B:  86% 7% 7%




- Molecule 2: RNA (42-MER)

Chain A:  98%



### 4.2.5 Score per residue for model 5

- Molecule 1: Adenosine deaminase

Chain B:  83% 10% 7%




- Molecule 2: RNA (42-MER)

Chain A:  98% .



### 4.2.6 Score per residue for model 6

- Molecule 1: Adenosine deaminase

Chain B:  86% 7% 7%




- Molecule 2: RNA (42-MER)

Chain A:  100%

There are no outlier residues in this chain.

### 4.2.7 Score per residue for model 7

- Molecule 1: Adenosine deaminase

Chain B:  87% 6% 7%




- Molecule 2: RNA (42-MER)

Chain A:  98% .



### 4.2.8 Score per residue for model 8

- Molecule 1: Adenosine deaminase

Chain B:  86% • • 7%




- Molecule 2: RNA (42-MER)

Chain A:  93% 7%



### 4.2.9 Score per residue for model 9

- Molecule 1: Adenosine deaminase

Chain B:  82% 10% • 7%




- Molecule 2: RNA (42-MER)

Chain A:  95% 5%



### 4.2.10 Score per residue for model 10

- Molecule 1: Adenosine deaminase

Chain B:  86% 7% 7%




- Molecule 2: RNA (42-MER)

Chain A:  98% •



### 4.2.11 Score per residue for model 11

- Molecule 1: Adenosine deaminase

Chain B:  86% 6% 7%




- Molecule 2: RNA (42-MER)


Chain A:  93% 7%




### 4.2.12 Score per residue for model 12

- Molecule 1: Adenosine deaminase

Chain B:  87% 6% 7%




- Molecule 2: RNA (42-MER)


Chain A:  90% 10%



### 4.2.13 Score per residue for model 13

- Molecule 1: Adenosine deaminase

Chain B:  89% 6% 7%




- Molecule 2: RNA (42-MER)

Chain A:  98%



#### 4.2.14 Score per residue for model 14


- Molecule 1: Adenosine deaminase

Chain B:  83% 10% 7%



- Molecule 2: RNA (42-MER)


Chain A:  93% • 5%



#### 4.2.15 Score per residue for model 15


- Molecule 1: Adenosine deaminase

Chain B:  89% • 7%




- Molecule 2: RNA (42-MER)

Chain A:  93% 7%



#### 4.2.16 Score per residue for model 16


- Molecule 1: Adenosine deaminase

Chain B:  87% 6% 7%




- Molecule 2: RNA (42-MER)

Chain A:  95% 5%



### 4.2.17 Score per residue for model 17

- Molecule 1: Adenosine deaminase

Chain B:  85% 8% 7%




- Molecule 2: RNA (42-MER)


Chain A:  90% 10%



### 4.2.18 Score per residue for model 18

- Molecule 1: Adenosine deaminase

Chain B:  85% 8% 7%




- Molecule 2: RNA (42-MER)

Chain A:  98%



### 4.2.19 Score per residue for model 19

- Molecule 1: Adenosine deaminase

Chain B:  86% 7% 7%




- Molecule 2: RNA (42-MER)

Chain A:  93% 7%



#### 4.2.20 Score per residue for model 20


- Molecule 1: Adenosine deaminase

Chain B:  87% 6% 7%



- Molecule 2: RNA (42-MER)

Chain A:  93% • 5%



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *SIMULATED ANNEALING*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *STRUCTURES WITH THE LOWEST ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
AMBER	refinement	
TOPSPIN	structure solution	2.0
CYANA	structure solution	
PROCHECKNMR	structure solution	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	B	0.49±0.00	0±0/517 (0.0±0.0%)	0.71±0.02	0±0/691 (0.0±0.0%)
2	A	0.95±0.00	0±0/993 (0.0±0.0%)	1.47±0.01	1±1/1546 (0.1±0.1%)
All	All	0.82	0/30200 (0.0%)	1.29	20/44740 (0.0%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	A	22	C	O4'-C1'-N1	5.75	112.80	108.20	19	7
2	A	22	C	P-O3'-C3'	5.50	126.31	119.70	20	1
2	A	23	G	O4'-C1'-N9	5.20	112.36	108.20	20	1
2	A	19	C	O4'-C1'-N1	5.17	112.33	108.20	7	7
2	A	27	U	O4'-C1'-N1	5.08	112.27	108.20	2	4

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	508	515	515	0±0
2	A	890	455	455	0±0
All	All	27960	19400	19400	11

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:55:LEU:HB3	1:B:109:PHE:CE2	0.66	2.25	9	3
1:B:81:VAL:HG12	1:B:82:ASP:H	0.45	1.72	8	1
1:B:63:PHE:C	1:B:64:LEU:HD22	0.44	2.33	18	2
2:A:20:U:H2'	2:A:22:C:C5	0.43	2.48	14	2
1:B:112:HIS:CG	1:B:113:LEU:H	0.41	2.34	14	2
2:A:27:U:C5	2:A:28:C:C5	0.40	3.08	8	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	65/71 (92%)	58±1 (90±2%)	6±2 (10±2%)	1±1 (1±1%)	29	74
All	All	1300/1420 (92%)	1165 (90%)	124 (10%)	11 (1%)	29	74

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	65	SER	9
1	B	81	VAL	1
1	B	82	ASP	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	54/58 (93%)	50±1 (93±1%)	4±1 (7±1%)	22	68

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1080/1160 (93%)	1001 (93%)	79 (7%)	22 68

All 11 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	59	LEU	20
1	B	76	VAL	20
1	B	66	GLU	18
1	B	79	VAL	9
1	B	55	LEU	4
1	B	110	ASN	3
1	B	56	ARG	1
1	B	50	MET	1
1	B	95	LEU	1
1	B	60	LYS	1
1	B	93	LYS	1

### 6.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
2	A	41/42 (98%)	1±1 (2±2%)	0±0 (1±1%)	0.73±0.02
All	All	820/840 (98%)	15 (2%)	6 (1%)	0.73

The overall RNA backbone suiteness is 0.73.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
2	A	21	U	9
2	A	23	G	4
2	A	22	C	1
2	A	24	G	1

All unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
2	A	20	U	3
2	A	21	U	2
2	A	22	C	1

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided