



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:12 AM GMT

PDB ID : 3L4B
Title : Crystal Structure of an Octomeric Two-Subunit TrkA K⁺ Channel Ring Gating Assembly, TM1088A:TM1088B, from *Thermotoga maritima*
Authors : Deller, M.C.; Johnson, H.A.; Miller, M.; Spraggon, G.; Wilson, I.A.; Lesley, S.A.; Joint Center for Structural Genomics (JCSG)
Deposited on : 2009-12-18
Resolution : 3.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1000 (3.56-3.36)
Clashscore	102246	1090 (3.56-3.36)
Ramachandran outliers	100387	1057 (3.56-3.36)
Sidechain outliers	100360	1058 (3.56-3.36)
RSRZ outliers	91569	1005 (3.56-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	143	<div> <div>81%</div> <div>10%</div> <div>8%</div> </div>
1	B	143	<div> <div>2%</div> <div>78%</div> <div>11%</div> <div>10%</div> </div>
1	E	143	<div> <div>2%</div> <div>78%</div> <div>7%</div> <div>13%</div> </div>
1	F	143	<div> <div>2%</div> <div>78%</div> <div>12%</div> <div>10%</div> </div>
2	C	218	<div> <div>3%</div> <div>84%</div> <div>14%</div> <div>2%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	218	<div><div></div><div>2%</div><div>82%</div><div>14%</div><div></div><div></div></div>
2	G	218	<div><div></div><div>4%</div><div>86%</div><div>9%</div><div></div><div></div></div>
2	H	218	<div><div></div><div>3%</div><div>86%</div><div>7%</div><div></div><div>6%</div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9045 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TrkA K+ Channel protien TM1088A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	131	Total	C	N	O	S	0	0	0
			908	572	152	177	7			
1	B	129	Total	C	N	O	S	0	0	0
			907	568	155	176	8			
1	E	125	Total	C	N	O	S	0	0	0
			818	508	139	165	6			
1	F	128	Total	C	N	O	S	0	0	0
			900	567	146	179	8			

- Molecule 2 is a protein called TrkA K+ Channel protien TM1088B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	216	Total	C	N	O	S	0	0	0
			1409	878	247	279	5			
2	D	216	Total	C	N	O	S	0	0	0
			1422	884	248	284	6			
2	G	211	Total	C	N	O	S	0	0	0
			1324	814	236	268	6			
2	H	205	Total	C	N	O	S	0	0	0
			1264	767	236	255	6			

- Molecule 3 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	E	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	F	1	Total	C	N	O	P	0	0
			23	10	5	7	1		


- Molecule 4 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	X	0	0
			1	1		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TrkA K+ Channel protien TM1088A

Chain A: 




- Molecule 1: TrkA K+ Channel protien TM1088A

Chain B: 




- Molecule 1: TrkA K+ Channel protien TM1088A

Chain E: 





- Molecule 1: TrkA K+ Channel protien TM1088A

Chain F: 




- Molecule 2: TrkA K+ Channel protien TM1088B


Chain C: 




- Molecule 2: TrkA K⁺ Channel protien TM1088B



- Molecule 2: TrkA K⁺ Channel protien TM1088B



- Molecule 2: TrkA K⁺ Channel protien TM1088B



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	96.78Å 96.78Å 305.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.39 – 3.45 49.33 – 3.44	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.39-3.45) 99.9 (49.33-3.44)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 3.48Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.259 , 0.308 0.255 , 0.303	Depositor DCC
R_{free} test set	1164 reflections (5.41%)	DCC
Wilson B-factor (Å ²)	102.1	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 105.6	EDS
Estimated twinning fraction	0.098 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 22804 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	9045	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/925	0.51	0/1259
1	B	0.36	0/923	0.49	0/1252
1	E	0.34	0/829	0.46	0/1131
1	F	0.36	0/916	0.47	0/1247
2	C	0.35	0/1428	0.50	0/1960
2	D	0.35	0/1441	0.53	1/1980 (0.1%)
2	G	0.33	0/1340	0.50	0/1838
2	H	0.31	0/1279	0.48	0/1754
All	All	0.35	0/9081	0.50	1/12421 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	117	LEU	CA-CB-CG	5.72	128.45	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	908	0	733	8	0
1	B	907	0	742	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	818	0	618	10	0
1	F	900	0	746	10	0
2	C	1409	0	1144	14	0
2	D	1422	0	1146	18	0
2	G	1324	0	983	10	0
2	H	1264	0	900	6	0
3	A	23	0	12	3	0
3	B	23	0	12	1	0
3	E	23	0	12	4	0
3	F	23	0	12	2	0
4	B	1	0	0	0	0
All	All	9045	0	7060	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:126:ALA:O	1:E:129:MET:HG3	1.90	0.71
1:A:130:ILE:HA	1:A:133:VAL:HG12	1.78	0.65
2:C:28:ILE:HG21	2:C:39:PHE:HD2	1.63	0.64
1:E:61:GLU:HG2	2:H:14:TYR:OH	1.97	0.64
2:D:28:ILE:HG21	2:D:39:PHE:HD2	1.63	0.62
1:E:124:CYS:O	1:E:127:VAL:HG12	1.99	0.61
2:D:86:VAL:HG13	2:D:92:VAL:HB	1.82	0.60
1:F:76:VAL:HG23	1:F:99:VAL:HG21	1.84	0.59
2:G:4:ILE:HD13	2:G:67:ASP:HB3	1.84	0.58
2:C:98:LEU:HD12	2:C:116:VAL:HG21	1.88	0.56
1:E:124:CYS:O	1:E:128:LEU:HD12	2.06	0.55
1:E:36:VAL:HG21	1:E:68:CYS:SG	2.46	0.55
2:C:128:GLU:OE1	2:D:22:ARG:NH1	2.39	0.55
2:D:73:THR:HG23	2:D:79:ASN:OD1	2.06	0.55
2:C:188:ARG:HG3	2:C:189:GLY:H	1.73	0.54
2:G:73:THR:HG23	2:G:79:ASN:OD1	2.07	0.54
1:E:16:GLY:HA3	3:E:144:AMP:P	2.47	0.54
1:B:61:GLU:HG2	2:C:14:TYR:OH	2.08	0.54
2:G:70:VAL:HG22	2:G:96:VAL:HB	1.89	0.53
1:A:43:PHE:HE2	1:A:56:VAL:HG22	1.73	0.53
1:A:16:GLY:HA3	3:A:144:AMP:P	2.49	0.53
2:G:4:ILE:N	2:G:4:ILE:HD12	2.24	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:ARG:N	3:A:144:AMP:O1P	2.40	0.52
1:F:93:ALA:HB1	1:F:99:VAL:HG11	1.92	0.51
1:F:125:PRO:O	1:F:129:MET:HG2	2.10	0.51
2:C:28:ILE:HG21	2:C:39:PHE:CD2	2.46	0.50
1:B:88:PHE:HB2	2:D:77:GLU:HA	1.94	0.50
1:B:38:LYS:HB2	3:B:144:AMP:C4	2.46	0.50
2:C:74:PRO:HB2	2:D:10:THR:HG21	1.95	0.49
2:D:2:LYS:HG3	2:D:67:ASP:OD1	2.13	0.49
2:D:50:GLY:HA3	2:D:57:ILE:HD12	1.95	0.48
2:C:16:ALA:O	2:C:20:LEU:HD12	2.14	0.48
1:A:38:LYS:HB2	3:A:144:AMP:C4	2.49	0.48
1:F:43:PHE:HE2	1:F:56:VAL:HG22	1.78	0.47
1:B:9:TYR:HD2	1:B:73:ALA:HA	1.80	0.47
2:D:128:GLU:HA	2:D:131:ILE:CG1	2.45	0.47
2:C:73:THR:HG23	2:C:79:ASN:OD1	2.15	0.47
2:D:63:VAL:HG22	2:D:90:PHE:CD2	2.50	0.46
2:G:10:THR:HG21	2:H:74:PRO:HB2	1.97	0.46
2:D:118:ASN:OD1	2:D:120:THR:HG23	2.15	0.46
2:D:58:LEU:O	2:D:63:VAL:HG13	2.15	0.46
1:F:17:ARG:HG3	3:F:144:AMP:O1P	2.15	0.46
1:F:36:VAL:HG21	1:F:68:CYS:SG	2.56	0.45
1:E:16:GLY:HA3	3:E:144:AMP:O2P	2.16	0.45
2:D:28:ILE:HG21	2:D:39:PHE:CD2	2.49	0.44
2:G:34:GLU:HA	2:G:34:GLU:OE1	2.18	0.44
2:H:118:ASN:OD1	2:H:120:THR:HG23	2.18	0.43
2:C:118:ASN:OD1	2:C:120:THR:HG23	2.18	0.43
2:C:98:LEU:HD12	2:C:116:VAL:CG2	2.48	0.43
1:B:85:THR:HA	2:D:77:GLU:HB2	2.00	0.43
1:B:87:PHE:HZ	2:D:84:GLN:HG3	1.84	0.43
1:A:87:PHE:HB2	1:A:115:PHE:CZ	2.54	0.43
1:B:87:PHE:HB2	1:B:115:PHE:CZ	2.54	0.43
1:E:17:ARG:N	3:E:144:AMP:O1P	2.46	0.42
2:D:75:ARG:NH1	2:D:77:GLU:OE2	2.51	0.42
1:F:37:ASP:OD1	3:F:144:AMP:O2'	2.36	0.42
1:B:93:ALA:HB3	1:B:102:VAL:HG11	2.00	0.42
2:C:188:ARG:HG3	2:C:189:GLY:N	2.34	0.42
1:F:93:ALA:HB1	1:F:99:VAL:CG1	2.49	0.42
1:B:93:ALA:CB	1:B:102:VAL:HG11	2.50	0.42
2:G:74:PRO:HB2	2:H:10:THR:HG21	2.02	0.42
1:E:87:PHE:HB2	1:E:115:PHE:CZ	2.55	0.42
2:D:101:ASP:HA	2:D:102:PRO:HD2	1.93	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:34:VAL:HG22	1:B:53:PHE:HD2	1.85	0.41
2:G:118:ASN:OD1	2:G:120:THR:HG23	2.20	0.41
1:F:86:ASN:HA	1:F:89:ILE:HD12	2.03	0.41
1:A:35:VAL:HB	1:A:54:THR:HG22	2.03	0.41
2:C:70:VAL:HG22	2:C:96:VAL:HB	2.01	0.41
2:G:22:ARG:NH1	2:G:22:ARG:HG2	2.36	0.41
2:G:4:ILE:N	2:G:4:ILE:CD1	2.84	0.41
1:E:16:GLY:HA3	3:E:144:AMP:O1P	2.21	0.41
1:A:36:VAL:HG21	1:A:68:CYS:SG	2.60	0.41
2:D:86:VAL:HG13	2:D:92:VAL:CB	2.50	0.41
2:C:72:LEU:CD2	2:C:98:LEU:HD13	2.51	0.40
2:H:101:ASP:HA	2:H:102:PRO:HD2	1.95	0.40
2:H:73:THR:HG23	2:H:79:ASN:OD1	2.21	0.40
1:B:35:VAL:HB	1:B:54:THR:HG22	2.04	0.40
1:F:40:GLU:HG3	1:F:43:PHE:HD2	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/143 (90%)	121 (94%)	8 (6%)	0	100	100
1	B	127/143 (89%)	120 (94%)	7 (6%)	0	100	100
1	E	121/143 (85%)	115 (95%)	6 (5%)	0	100	100
1	F	126/143 (88%)	119 (94%)	7 (6%)	0	100	100
2	C	214/218 (98%)	194 (91%)	15 (7%)	5 (2%)	8	46
2	D	214/218 (98%)	195 (91%)	14 (6%)	5 (2%)	8	46
2	G	205/218 (94%)	187 (91%)	13 (6%)	5 (2%)	7	45
2	H	199/218 (91%)	180 (90%)	14 (7%)	5 (2%)	7	44

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1335/1444 (92%)	1231 (92%)	84 (6%)	20 (2%)	13	54

All (20) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	139	ILE
2	D	139	ILE
2	G	139	ILE
2	H	139	ILE
2	C	142	LEU
2	G	142	LEU
2	H	142	LEU
2	C	8	GLY
2	D	8	GLY
2	D	136	PHE
2	D	142	LEU
2	G	8	GLY
2	G	136	PHE
2	H	8	GLY
2	H	136	PHE
2	C	136	PHE
2	G	7	GLY
2	D	7	GLY
2	H	7	GLY
2	C	7	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	72/124 (58%)	70 (97%)	2 (3%)	51	82
1	B	72/124 (58%)	68 (94%)	4 (6%)	26	66
1	E	57/124 (46%)	53 (93%)	4 (7%)	19	58
1	F	77/124 (62%)	74 (96%)	3 (4%)	39	75

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	104/192 (54%)	91 (88%)	13 (12%)	6	28
2	D	104/192 (54%)	90 (86%)	14 (14%)	5	25
2	G	86/192 (45%)	77 (90%)	9 (10%)	8	37
2	H	75/192 (39%)	69 (92%)	6 (8%)	15	51
All	All	647/1264 (51%)	592 (92%)	55 (8%)	13	49

All (55) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	94	ARG
1	A	96	MET
1	B	53	PHE
1	B	56	VAL
1	B	94	ARG
1	B	96	MET
2	C	10	THR
2	C	11	THR
2	C	19	MET
2	C	27	VAL
2	C	33	ARG
2	C	38	GLU
2	C	73	THR
2	C	98	LEU
2	C	106	GLU
2	C	114	THR
2	C	120	THR
2	C	206	GLU
2	C	210	THR
2	D	2	LYS
2	D	10	THR
2	D	11	THR
2	D	19	MET
2	D	22	ARG
2	D	63	VAL
2	D	73	THR
2	D	86	VAL
2	D	98	LEU
2	D	114	THR
2	D	120	THR
2	D	143	GLU
2	D	166	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	210	THR
1	E	96	MET
1	E	127	VAL
1	E	128	LEU
1	E	129	MET
1	F	53	PHE
1	F	96	MET
1	F	128	LEU
2	G	19	MET
2	G	20	LEU
2	G	22	ARG
2	G	31	LYS
2	G	73	THR
2	G	106	GLU
2	G	114	THR
2	G	120	THR
2	G	210	THR
2	H	19	MET
2	H	22	ARG
2	H	73	THR
2	H	106	GLU
2	H	114	THR
2	H	120	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 1 is unknown - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	AMP	A	144	-	20,25,25	0.74	1 (5%)	22,38,38	2.71	5 (22%)
3	AMP	B	144	-	20,25,25	0.80	1 (5%)	22,38,38	2.44	2 (9%)
3	AMP	E	144	-	20,25,25	0.88	1 (5%)	22,38,38	2.60	6 (27%)
3	AMP	F	144	-	20,25,25	0.75	0	22,38,38	2.37	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AMP	A	144	-	-	0/6/26/26	0/3/3/3
3	AMP	B	144	-	-	0/6/26/26	0/3/3/3
3	AMP	E	144	-	-	0/6/26/26	0/3/3/3
3	AMP	F	144	-	-	0/6/26/26	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	144	AMP	C5-N7	-2.16	1.32	1.39
3	A	144	AMP	C5-N7	-2.02	1.32	1.39
3	E	144	AMP	O4'-C1'	2.61	1.44	1.41

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	144	AMP	N3-C2-N1	-10.53	120.83	128.89
3	A	144	AMP	N3-C2-N1	-10.02	121.22	128.89
3	E	144	AMP	N3-C2-N1	-9.70	121.46	128.89
3	F	144	AMP	N3-C2-N1	-9.70	121.47	128.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	144	AMP	C2'-C1'-N9	-4.84	106.90	114.29
3	E	144	AMP	C2'-C1'-N9	-3.17	109.45	114.29
3	A	144	AMP	O3P-P-O1P	-2.67	102.00	110.58
3	E	144	AMP	O3P-P-O1P	-2.50	102.54	110.58
3	A	144	AMP	C4-C5-N7	-2.22	107.44	109.48
3	E	144	AMP	O5'-P-O1P	-2.16	101.64	107.14
3	F	144	AMP	C1'-N9-C4	-2.13	123.73	126.94
3	E	144	AMP	C4'-O4'-C1'	2.21	112.15	109.72
3	B	144	AMP	O2P-P-O5'	2.52	113.81	106.56
3	A	144	AMP	O2P-P-O5'	2.74	114.46	106.56
3	E	144	AMP	O2P-P-O5'	3.62	117.00	106.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	144	AMP	3	0
3	B	144	AMP	1	0
3	E	144	AMP	4	0
3	F	144	AMP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	131/143 (91%)	-0.17	0 100 100	35, 49, 55, 58	0
1	B	129/143 (90%)	-0.23	2 (1%) 74 68	42, 49, 55, 58	0
1	E	125/143 (87%)	0.08	3 (2%) 62 55	28, 49, 53, 58	0
1	F	128/143 (89%)	-0.10	3 (2%) 64 57	42, 49, 54, 58	0
2	C	216/218 (99%)	-0.16	6 (2%) 56 50	39, 49, 53, 57	0
2	D	216/218 (99%)	-0.13	5 (2%) 64 57	40, 49, 54, 60	0
2	G	211/218 (96%)	0.00	9 (4%) 39 33	39, 49, 54, 58	0
2	H	205/218 (94%)	-0.17	6 (2%) 55 48	41, 49, 54, 58	0
All	All	1361/1444 (94%)	-0.11	34 (2%) 61 54	28, 49, 54, 60	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	133	VAL	5.1
2	D	200	TYR	3.6
2	G	137	SER	3.3
1	B	133	VAL	3.2
2	D	148	PHE	3.2
1	F	134	LYS	3.0
2	G	111	MET	2.9
2	H	172	ASP	2.9
2	C	200	TYR	2.9
2	C	151	VAL	2.8
2	C	147	GLU	2.8
2	G	150	SER	2.7
1	E	26	ALA	2.7
2	D	215	LEU	2.7
2	G	108	PHE	2.6
2	G	148	PHE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	27	VAL	2.5
2	D	198	LYS	2.5
2	C	201	VAL	2.5
2	H	3	VAL	2.5
1	F	43	PHE	2.4
2	H	5	ILE	2.4
2	G	84	GLN	2.3
2	G	202	ILE	2.3
1	B	135	GLU	2.3
2	H	177	ALA	2.2
2	D	39	PHE	2.2
2	H	4	ILE	2.1
1	E	44	HIS	2.1
2	G	71	ILE	2.1
2	C	150	SER	2.1
2	G	161	GLY	2.1
2	C	187	PRO	2.0
1	E	29	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	AMP	A	144	23/23	0.93	0.22	0.24	78,80,90,95	0
3	AMP	B	144	23/23	0.95	0.18	-0.23	61,65,72,74	0
3	AMP	E	144	23/23	0.89	0.15	-0.70	84,96,100,102	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	AMP	F	144	23/23	0.95	0.15	-1.07	63,67,76,80	0
4	UNX	B	145	1/1	0.91	0.38	-	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.