



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:16 PM GMT

PDB ID : 4L4H  
Title : Crystal structure of mouse Ryanodine Receptor isoform 2 (RyR2) 1-547  
Authors : Kimlicka, L.; Tung, C.C.; Van Petegem, F.  
Deposited on : 2013-06-07  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	547	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PGE	A	602	-	-	X	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	A	603	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3665 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ryanodine receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	466	Total	C	N	O	S	0	5	0
			3564	2250	631	665	18			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	82	Total	O	0	1
			83	83		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 4 <sub>2</sub> 2 <sub>1</sub> 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.04 Å   78.04 Å   248.38 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	48.71 – 2.00 48.59 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.71-2.00) 99.7 (48.59-2.00)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.16	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.56 (at 2.00 Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.220   ,   0.251 0.219   ,   0.250	Depositor DCC
$R_{free}$ test set	2645 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	33.7	Xtriage
Anisotropy	0.263	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 54.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 52836 reflections (0.004%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3665	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, PGE, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.99	4/3648 (0.1%)	0.91	6/4945 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	131	CYS	CB-SG	-7.69	1.69	1.82
1	A	534	TYR	CD1-CE1	5.54	1.47	1.39
1	A	132	CYS	CB-SG	-5.08	1.73	1.81
1	A	532	SER	CA-CB	5.06	1.60	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	542	ARG	NE-CZ-NH2	-5.79	117.40	120.30
1	A	258	ARG	NE-CZ-NH1	-5.73	117.44	120.30
1	A	417	ARG	NE-CZ-NH1	-5.69	117.45	120.30
1	A	235	ARG	NE-CZ-NH2	-5.61	117.49	120.30
1	A	183	LEU	CB-CG-CD2	-5.16	102.23	111.00
1	A	542	ARG	NE-CZ-NH1	5.06	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3564	0	3380	71	0
2	A	1	0	0	0	0
3	A	10	0	14	8	0
4	A	7	0	10	3	0
5	A	83	0	0	2	0
All	All	3665	0	3404	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:602:PGE:H5	5:A:763:HOH:O	1.51	1.06
1:A:162:ILE:CD1	1:A:183:LEU:CD2	2.34	1.04
1:A:162:ILE:HD12	1:A:183:LEU:HD21	1.45	0.95
1:A:162:ILE:CD1	1:A:183:LEU:HD21	1.97	0.94
1:A:258:ARG:HH12	1:A:317:MET:HE2	1.38	0.86
1:A:224:ALA:H	1:A:288:HIS:HE1	1.25	0.84
1:A:162:ILE:HD12	1:A:183:LEU:CD2	2.05	0.84
1:A:162:ILE:HD13	1:A:183:LEU:CD2	2.06	0.82
1:A:125:TYR:CZ	1:A:417:ARG:HD2	2.15	0.81
1:A:224:ALA:H	1:A:288:HIS:CE1	2.01	0.78
1:A:162:ILE:HD13	1:A:183:LEU:HD23	1.66	0.76
1:A:125:TYR:CE1	1:A:417:ARG:HD2	2.22	0.74
1:A:233:VAL:H	3:A:602:PGE:H22	1.53	0.74
1:A:42:PHE:HB3	4:A:603:PEG:H11	1.70	0.74
1:A:162:ILE:CD1	1:A:183:LEU:HD23	2.18	0.74
1:A:308:LEU:HD13	1:A:393:MET:HG3	1.75	0.69
1:A:394:HIS:CD2	1:A:396:GLU:H	2.10	0.68
1:A:463:PHE:O	1:A:542:ARG:NH2	2.25	0.68
1:A:276:ARG:HH12	3:A:602:PGE:H42	1.58	0.67
1:A:431:ARG:NH2	5:A:702:HOH:O	2.31	0.64
1:A:355:LYS:HE3	1:A:358:ASP:OD2	1.98	0.64
1:A:119:ILE:CD1	1:A:162:ILE:HD11	2.28	0.63
1:A:232:ASP:HA	3:A:602:PGE:C3	2.29	0.62
1:A:258:ARG:NH1	1:A:317:MET:HE2	2.13	0.62
1:A:276:ARG:HH12	3:A:602:PGE:H52	1.66	0.61
1:A:119:ILE:HD13	1:A:162:ILE:HD11	1.83	0.61
1:A:69:LEU:HD23	1:A:121:LEU:CD2	2.32	0.60
1:A:232:ASP:HA	3:A:602:PGE:H3	1.83	0.60
1:A:276:ARG:NH1	3:A:602:PGE:H42	2.17	0.59

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:ARG:O	1:A:299:HIS:HE1	1.85	0.58
1:A:193:HIS:HD2	1:A:204:ASP:OD1	1.86	0.57
1:A:287:SER:HA	1:A:349:MET:SD	2.45	0.56
1:A:394:HIS:HD2	1:A:397:GLY:H	1.51	0.56
1:A:258:ARG:HH12	1:A:317:MET:CE	2.14	0.54
1:A:247:VAL:HG11	1:A:316:LEU:HD11	1.90	0.54
1:A:25:THR:CG2	1:A:32:GLN:HG3	2.38	0.54
1:A:224:ALA:N	1:A:288:HIS:HE1	2.01	0.53
1:A:258:ARG:NH1	1:A:317:MET:CE	2.71	0.53
1:A:112:THR:HG21	1:A:174:LYS:HE3	1.92	0.52
1:A:181:LEU:HD11	1:A:214:VAL:HG21	1.92	0.52
1:A:42:PHE:CB	4:A:603:PEG:H11	2.38	0.52
1:A:42:PHE:HB3	4:A:603:PEG:C1	2.41	0.50
1:A:394:HIS:HD2	1:A:396:GLU:H	1.53	0.50
1:A:394:HIS:CD2	1:A:397:GLY:H	2.30	0.49
1:A:365:HIS:HB2	1:A:393:MET:CE	2.42	0.49
1:A:299:HIS:HD2	1:A:302:THR:OG1	1.95	0.49
1:A:228:LEU:HD13	1:A:289:ILE:HB	1.94	0.49
1:A:52:THR:O	1:A:52:THR:CG2	2.61	0.48
1:A:417:ARG:HG3	1:A:417:ARG:HH11	1.78	0.47
1:A:232:ASP:HA	3:A:602:PGE:H32	1.96	0.47
1:A:69:LEU:HD23	1:A:121:LEU:HD23	1.95	0.47
1:A:481:ALA:O	1:A:485:ARG:HG3	2.15	0.46
1:A:515:ALA:HB2	1:A:523:GLY:HA3	1.98	0.45
1:A:377:VAL:O	1:A:377:VAL:HG23	2.15	0.45
1:A:365:HIS:HB2	1:A:393:MET:HE1	1.98	0.45
1:A:315:LEU:HD12	1:A:317:MET:CE	2.46	0.45
1:A:500:GLU:O	1:A:504:ARG:HG2	2.17	0.45
1:A:125:TYR:OH	1:A:417:ARG:HD2	2.16	0.44
1:A:365:HIS:HD2	1:A:368:THR:OG1	2.01	0.43
1:A:169:ARG:NH2	1:A:176:ARG:HD3	2.34	0.43
1:A:281:ARG:HB2	1:A:281:ARG:HE	1.47	0.42
1:A:315:LEU:HD12	1:A:317:MET:HE3	2.01	0.42
1:A:181:LEU:CD1	1:A:214:VAL:HG21	2.50	0.41
1:A:228:LEU:HD13	1:A:289:ILE:CB	2.50	0.41
1:A:193:HIS:CD2	1:A:204:ASP:OD1	2.70	0.41
1:A:52:THR:HG22	1:A:52:THR:O	2.21	0.41
1:A:276:ARG:HD2	1:A:278:GLU:OE2	2.21	0.41
1:A:125:TYR:CE1	1:A:417:ARG:CD	2.98	0.41
1:A:284:TRP:CE2	1:A:348:GLY:HA2	2.56	0.40
1:A:69:LEU:HD23	1:A:121:LEU:HD21	2.03	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:HIS:CE1	1:A:125:TYR:HB3	2.56	0.40
1:A:463:PHE:HB3	1:A:539:ALA:HB1	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	457/547 (84%)	448 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/470 (78%)	346 (94%)	21 (6%)	25	19

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LEU
1	A	32	GLN
1	A	62	LEU
1	A	66	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	74	SER
1	A	117	HIS
1	A	136	SER
1	A	213	SER
1	A	219	SER
1	A	228	LEU
1	A	229	ILE
1	A	234	LEU
1	A	251	GLU
1	A	281	ARG
1	A	315	LEU
1	A	388	GLN
1	A	421	SER
1	A	475	LYS
1	A	477	ASN
1	A	479	LEU
1	A	497	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	HIS
1	A	150	GLN
1	A	193	HIS
1	A	288	HIS
1	A	299	HIS
1	A	365	HIS
1	A	394	HIS
1	A	395	HIS
1	A	427	ASN
1	A	496	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PGE	A	602	-	9,9,9	0.54	0	8,8,8	0.97	0
4	PEG	A	603	-	6,6,6	0.71	0	5,5,5	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	A	602	-	-	0/7/7/7	0/0/0/0
4	PEG	A	603	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	PGE	8	0
4	A	603	PEG	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	466/547 (85%)	0.26	21 (4%) 37 38	18, 43, 73, 97	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	75	VAL	5.3
1	A	199	SER	4.3
1	A	29	HIS	4.3
1	A	201	TRP	3.7
1	A	543	GLY	3.4
1	A	197	GLY	3.4
1	A	378	ASP	3.2
1	A	11	ILE	2.9
1	A	200	SER	2.8
1	A	325	LYS	2.8
1	A	196	TYR	2.6
1	A	198	ASN	2.4
1	A	309	MET	2.4
1	A	33	GLN	2.4
1	A	73	LEU	2.4
1	A	350	GLY	2.4
1	A	162	ILE	2.4
1	A	352	SER	2.4
1	A	28	ILE	2.2
1	A	224	ALA	2.2
1	A	54	ASN	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PEG	A	603	7/7	0.89	0.18	2.13	39,55,63,67	0
2	CL	A	601	1/1	0.97	0.15	-0.20	48,48,48,48	0
3	PGE	A	602	10/10	0.93	0.12	-0.82	44,48,53,54	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.