



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:21 PM GMT

PDB ID : 4LCW  
Title : The structure of human MAIT TCR in complex with MR1-K43A-RL-6-Me-7OH  
Authors : Patel, O.; Rossjohn, J.  
Deposited on : 2013-06-24  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	
1	C	271	
2	B	99	
2	F	99	
3	D	203	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	203	 80% 11% 7%
4	E	245	 94% 5%
4	H	245	 91% 6%

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 12797 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major histocompatibility complex class I-related gene protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			2150	1379	370	390	11			
1	C	256	Total	C	N	O	S	0	0	0
			2064	1321	359	373	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	EXPRESSION TAG	UNP Q95460
A	43	ALA	LYS	ENGINEERED MUTATION	UNP Q95460
A	261	SER	CYS	ENGINEERED MUTATION	UNP Q95460
C	0	MET	-	EXPRESSION TAG	UNP Q95460
C	43	ALA	LYS	ENGINEERED MUTATION	UNP Q95460
C	261	SER	CYS	ENGINEERED MUTATION	UNP Q95460

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	96	Total	C	N	O	S	0	0	0
			769	494	131	142	2			
2	F	96	Total	C	N	O	S	0	0	0
			755	486	126	141	2			

- Molecule 3 is a protein called MAIT T cell receptor alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	200	Total	C	N	O	S	0	0	0
			1532	968	248	307	9			
3	G	188	Total	C	N	O	S	0	2	0
			1426	917	225	275	9			

- Molecule 4 is a protein called MAIT T cell receptor beta chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	243	Total	C	N	O	S	0	1	0
			1893	1193	329	362	9			
4	H	240	Total	C	N	O	S	0	0	0
			1822	1153	310	350	9			

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is 1-DEOXY-1-(7-HYDROXY-6-METHYL-2,4-DIOXO-3,4-DIHYDROPTERIDIN-8(2H)-YL)-D-RIBITOL (three-letter code: 1VY) (formula:  $C_{12}H_{16}N_4O_7$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			23	12	4	7		
6	C	1	Total	C	N	O	0	0
			23	12	4	7		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	65	Total	O	0	0
			65	65		
7	B	32	Total	O	0	0
			32	32		
7	C	51	Total	O	0	0
			51	51		
7	D	60	Total	O	0	0
			60	60		
7	E	59	Total	O	0	0
			59	59		
7	F	16	Total	O	0	0
			16	16		
7	G	32	Total	O	0	0
			32	32		
7	H	19	Total	O	0	0
			19	19		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major histocompatibility complex class I-related gene protein




- Molecule 1: Major histocompatibility complex class I-related gene protein



- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin





- Molecule 3: MAIT T cell receptor alpha chain







- Molecule 3: MAIT T cell receptor alpha chain



- Molecule 4: MAIT T cell receptor beta chain



- Molecule 4: MAIT T cell receptor beta chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	215.91Å 69.36Å 142.83Å 90.00° 104.30° 90.00°	Depositor
Resolution (Å)	35.14 – 2.40 75.02 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (35.14-2.40) 99.7 (75.02-2.40)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.05 (at 2.40Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, $R_{free}$	0.179 , 0.227 0.191 , 0.240	Depositor DCC
$R_{free}$ test set	4021 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.3	Xtriage
Anisotropy	0.548	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 47.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 80362 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12797	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 1VY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/2215	0.67	0/3014
1	C	0.52	0/2124	0.66	0/2888
2	B	0.49	0/792	0.67	0/1080
2	F	0.48	0/778	0.66	0/1064
3	D	0.53	0/1567	0.72	0/2130
3	G	0.51	0/1464	0.69	0/1989
4	E	0.52	0/1948	0.71	0/2656
4	H	0.48	0/1873	0.70	0/2562
All	All	0.51	0/12761	0.69	0/17383

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2150	0	2030	17	0
1	C	2064	0	1932	10	0
2	B	769	0	715	3	0
2	F	755	0	687	2	0
3	D	1532	0	1426	5	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	1426	0	1336	11	0
4	E	1893	0	1766	4	0
4	H	1822	0	1664	10	0
5	A	6	0	8	0	0
6	A	23	0	16	1	0
6	C	23	0	16	1	0
7	A	65	0	0	0	0
7	B	32	0	0	0	0
7	C	51	0	0	0	0
7	D	60	0	0	0	0
7	E	59	0	0	0	0
7	F	16	0	0	0	0
7	G	32	0	0	0	0
7	H	19	0	0	0	0
All	All	12797	0	11596	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:94:ARG:HH22	6:C:301:1VY:H14	1.53	0.74
1:A:94:ARG:HH22	6:A:302:1VY:H14	1.57	0.70
3:D:1:GLY:HA2	3:D:27:SER:H	1.60	0.66
1:A:77:LEU:HD13	1:A:92:TYR:HB2	1.80	0.63
3:D:150:ASP:HB2	3:D:177:LYS:HD2	1.82	0.62
4:H:155:HIS:HB3	4:H:216:TYR:HB2	1.83	0.59
1:C:214:TRP:HB2	1:C:225:ILE:HG12	1.84	0.58
1:A:259:GLU:HG2	1:A:264:HIS:CD2	2.39	0.57
1:A:259:GLU:HG2	1:A:264:HIS:HD2	1.69	0.57
1:A:226:ASP:HB2	1:C:82:ARG:HH12	1.70	0.56
1:C:77:LEU:HD13	1:C:92:TYR:HB2	1.87	0.55
3:D:54:LEU:HD13	3:D:63:PHE:HB2	1.89	0.55
1:C:63:THR:O	1:C:67:ARG:HG3	2.07	0.55
4:H:131:PRO:HG2	4:H:142:ALA:HB1	1.88	0.54
3:G:1:GLY:HA2	3:G:27:SER:H	1.71	0.54
4:E:86:THR:HG23	4:E:113:THR:HA	1.90	0.53
1:A:221:ILE:O	1:A:221:ILE:HG23	2.08	0.53
3:G:159:LEU:HB3	4:H:172:CYS:HB2	1.91	0.53
3:G:78:LEU:HG	3:G:107:ILE:HD12	1.92	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:MET:HG2	1:A:258:VAL:HG22	1.91	0.51
1:A:80:LEU:HD21	1:A:119:PHE:CZ	2.46	0.50
3:G:121:LEU:HB3	4:H:129:PHE:HB3	1.93	0.50
4:H:128:VAL:HG23	4:H:238:ALA:HB3	1.94	0.49
1:A:214:TRP:O	1:A:221:ILE:HG22	2.12	0.49
1:C:113:ALA:HB2	2:F:60:TRP:CE2	2.47	0.49
4:E:128:VAL:HG23	4:E:238:ALA:HB3	1.94	0.49
4:E:224:TRP:CZ2	4:E:226:GLN:HB2	2.48	0.48
4:E:119:LYS:O	4:E:228:ARG:NH2	2.45	0.48
1:A:156:TRP:HA	1:A:160:GLU:HB2	1.95	0.47
1:C:31:HIS:CD2	1:C:176:LEU:HD23	2.50	0.47
1:A:113:ALA:HB2	2:B:60:TRP:CE2	2.50	0.46
2:B:30:PHE:HZ	2:B:64:LEU:HD12	1.81	0.46
3:G:121:LEU:HB2	3:G:131:VAL:HG12	1.97	0.45
3:G:11:THR:HA	3:G:106:ILE:O	2.16	0.45
1:C:208:PRO:O	1:C:230:ILE:HD13	2.16	0.45
4:H:130:GLU:HG2	4:H:130:GLU:H	1.53	0.45
2:F:27:VAL:O	2:F:63:TYR:HA	2.17	0.45
4:H:131:PRO:CG	4:H:142:ALA:HB1	2.47	0.45
3:G:28:GLY:HA3	3:G:93[B]:SER:HB3	1.99	0.45
3:G:134:PHE:HB2	3:G:186:PHE:CE1	2.53	0.44
1:A:211:TYR:HB2	1:A:259:GLU:HB2	1.99	0.44
3:G:31:GLY:HA3	3:G:48:TYR:CE1	2.53	0.44
4:H:204:ASN:HB3	4:H:207:ASN:HD22	1.82	0.43
1:A:198:LEU:HD12	1:A:244:ILE:HD11	2.00	0.43
3:G:4:ILE:HD12	3:G:98:ILE:O	2.18	0.43
1:A:116:GLY:O	2:B:3:ARG:NH2	2.52	0.43
3:D:89:ALA:HB1	3:D:97:LEU:HD22	2.00	0.43
3:D:6:GLN:HE21	3:D:100:GLY:HA3	1.84	0.42
3:G:38:HIS:CD2	3:G:84:ALA:HB2	2.55	0.42
1:C:208:PRO:O	1:C:230:ILE:HG21	2.19	0.42
1:C:137:HIS:O	1:C:141:GLN:HG2	2.19	0.41
1:A:149:GLU:HG2	4:H:99:GLU:OE2	2.21	0.41
4:H:212:GLN:HG3	4:H:235:ILE:HG23	2.02	0.41
1:A:9:ARG:O	1:A:93:GLN:HA	2.21	0.40
1:A:80:LEU:HA	1:A:80:LEU:HD23	1.83	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/271 (96%)	251 (96%)	7 (3%)	2 (1%)	24	35
1	C	248/271 (92%)	243 (98%)	5 (2%)	0	100	100
2	B	94/99 (95%)	94 (100%)	0	0	100	100
2	F	94/99 (95%)	92 (98%)	1 (1%)	1 (1%)	17	25
3	D	198/203 (98%)	194 (98%)	4 (2%)	0	100	100
3	G	184/203 (91%)	178 (97%)	6 (3%)	0	100	100
4	E	242/245 (99%)	237 (98%)	5 (2%)	0	100	100
4	H	238/245 (97%)	232 (98%)	6 (2%)	0	100	100
All	All	1558/1636 (95%)	1521 (98%)	34 (2%)	3 (0%)	52	69

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	74	GLU
1	A	193	PRO
1	A	222	VAL

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/240 (94%)	218 (97%)	7 (3%)	47	69
1	C	213/240 (89%)	207 (97%)	6 (3%)	51	72

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	83/94 (88%)	79 (95%)	4 (5%)	31	49
2	F	80/94 (85%)	76 (95%)	4 (5%)	30	48
3	D	167/180 (93%)	160 (96%)	7 (4%)	36	56
3	G	150/180 (83%)	140 (93%)	10 (7%)	20	31
4	E	201/211 (95%)	196 (98%)	5 (2%)	55	76
4	H	189/211 (90%)	182 (96%)	7 (4%)	41	62
All	All	1308/1450 (90%)	1258 (96%)	50 (4%)	40	60

All (50) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	ARG
1	A	54	LEU
1	A	57	ASP
1	A	73	PHE
1	A	100	LEU
1	A	149	GLU
1	A	226	ASP
2	B	4	THR
2	B	64	LEU
2	B	70	PHE
2	B	89	GLN
1	C	57	ASP
1	C	73	PHE
1	C	82	ARG
1	C	100	LEU
1	C	149	GLU
1	C	259	GLU
3	D	25	GLN
3	D	27	SER
3	D	54	LEU
3	D	62	SER
3	D	81	LYS
3	D	93	SER
3	D	145	GLN
4	E	97	THR
4	E	99	GLU
4	E	103	GLU
4	E	194	ARG
4	E	237	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	F	35	ILE
2	F	36	GLU
2	F	70	PHE
2	F	96	ASP
3	G	52	ASP
3	G	66	ARG
3	G	70	TYR
3	G	78	LEU
3	G	93[A]	SER
3	G	93[B]	SER
3	G	120	GLN
3	G	131	VAL
3	G	182	CYS
3	G	189	SER
4	H	67	SER
4	H	97	THR
4	H	99	GLU
4	H	112	LEU
4	H	130	GLU
4	H	166	GLU
4	H	172	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	123	ASN
1	A	148	HIS
1	A	203	HIS
1	A	264	HIS
1	C	148	HIS
1	C	264	HIS
3	D	3	ASN
3	D	6	GLN
4	E	208	HIS
2	F	17	ASN
2	F	31	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GOL	A	301	-	5,5,5	0.14	0	5,5,5	0.56	0
6	1VY	A	302	-	20,24,24	1.61	4 (20%)	17,35,35	2.26	6 (35%)
6	1VY	C	301	-	20,24,24	1.83	3 (15%)	17,35,35	2.46	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	301	-	-	0/4/4/4	0/0/0/0
6	1VY	A	302	-	-	0/14/14/14	0/2/2/2
6	1VY	C	301	-	-	0/14/14/14	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	301	1VY	C9-C6	-4.58	1.41	1.50
6	A	302	1VY	C9-C6	-4.20	1.42	1.50
6	A	302	1VY	C8A-N8	-2.10	1.36	1.39
6	A	302	1VY	C4A-N5	2.10	1.36	1.33

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	301	1VY	C1'-N8	2.52	1.51	1.48
6	A	302	1VY	C6-N5	4.11	1.36	1.32
6	C	301	1VY	C6-N5	4.83	1.36	1.32

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	301	1VY	C4-C4A-C8A	-3.59	117.64	119.94
6	C	301	1VY	C4A-C4-N3	-2.71	119.89	123.59
6	A	302	1VY	C4A-C4-N3	-2.68	119.93	123.59
6	A	302	1VY	C4-C4A-C8A	-2.64	118.25	119.94
6	C	301	1VY	O2'-C2'-C3'	2.19	114.53	109.02
6	C	301	1VY	C4-C4A-N5	2.28	121.48	118.72
6	A	302	1VY	C9-C6-N5	2.34	120.18	116.94
6	A	302	1VY	C4-C4A-N5	2.39	121.62	118.72
6	C	301	1VY	C9-C6-N5	2.46	120.35	116.94
6	A	302	1VY	O2'-C2'-C3'	2.74	115.89	109.02
6	A	302	1VY	C4-N3-C2	6.73	121.07	115.25
6	C	301	1VY	C4-N3-C2	7.67	121.88	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	302	1VY	1	0
6	C	301	1VY	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	264/271 (97%)	-0.17	1 (0%) 93 93	17, 30, 53, 70	1 (0%)
1	C	256/271 (94%)	-0.10	1 (0%) 93 93	16, 32, 58, 75	0
2	B	96/99 (96%)	-0.19	0 100 100	16, 33, 55, 70	0
2	F	96/99 (96%)	-0.07	0 100 100	23, 47, 64, 85	0
3	D	200/203 (98%)	-0.19	1 (0%) 91 91	12, 28, 54, 64	0
3	G	188/203 (92%)	0.15	3 (1%) 74 74	17, 39, 69, 84	0
4	E	243/245 (99%)	-0.18	0 100 100	18, 29, 48, 69	1 (0%)
4	H	240/245 (97%)	-0.01	2 (0%) 87 87	19, 41, 63, 79	0
All	All	1583/1636 (96%)	-0.09	8 (0%) 91 91	12, 33, 61, 85	2 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	H	96	TRP	4.2
3	G	149	SER	3.3
3	G	132	CYS	3.3
3	D	1	GLY	2.9
1	A	221	ILE	2.7
1	C	246	LEU	2.5
3	G	143	VAL	2.5
4	H	221	ASN	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
6	1VY	C	301	23/23	0.97	0.16	0.70	19,26,37,43	0
5	GOL	A	301	6/6	0.88	0.16	0.21	37,45,49,50	0
6	1VY	A	302	23/23	0.96	0.12	-1.05	25,33,38,40	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.