



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:21 PM GMT

PDB ID : 4LD0
Title : T. thermophilus RuvC in complex with Holliday junction substrate
Authors : Gorecka, K.M.; Komorowska, W.; Nowotny, M.
Deposited on : 2013-06-24
Resolution : 3.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1268 (4.02-3.50)
Clashscore	102246	1407 (4.02-3.50)
Ramachandran outliers	100387	1346 (4.02-3.50)
Sidechain outliers	100360	1342 (4.02-3.50)
RSRZ outliers	91569	1276 (4.02-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	169	<div> <div>4%</div> <div>69%</div> <div>23%</div> <div>7%</div> </div>
1	B	169	<div> <div>71%</div> <div>20%</div> <div>7%</div> </div>
2	C	31	<div> <div>3%</div> <div>26%</div> <div>65%</div> <div>10%</div> </div>
3	D	13	<div> <div>54%</div> <div>38%</div> <div>8%</div> </div>
4	F	11	<div> <div>36%</div> <div>55%</div> <div>9%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Crossover junction endodeoxyribonuclease RuvC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	158	Total	C	N	O	S	0	0	0
			1153	742	206	201	4			
1	B	157	Total	C	N	O	S	0	0	0
			1090	695	200	191	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP Q5SJC4
A	-1	PRO	-	EXPRESSION TAG	UNP Q5SJC4
A	0	HIS	-	EXPRESSION TAG	UNP Q5SJC4
A	70	GLN	GLU	ENGINEERED MUTATION	UNP Q5SJC4
B	-2	GLY	-	EXPRESSION TAG	UNP Q5SJC4
B	-1	PRO	-	EXPRESSION TAG	UNP Q5SJC4
B	0	HIS	-	EXPRESSION TAG	UNP Q5SJC4
B	70	GLN	GLU	ENGINEERED MUTATION	UNP Q5SJC4

- Molecule 2 is a DNA chain called DNA 31-MER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	31	Total	C	N	O	P	0	0	0
			631	302	112	187	30			

- Molecule 3 is a DNA chain called DNA 13-MER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	13	Total	C	N	O	P	0	0	0
			259	126	42	79	12			


- Molecule 4 is a DNA chain called DNA 11-MER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	11	Total	C	N	O	P	0	0	0
			228	108	45	64	11			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Crossover junction endodeoxyribonuclease RuvC



- Molecule 1: Crossover junction endodeoxyribonuclease RuvC



- Molecule 2: DNA 31-MER




- Molecule 3: DNA 13-MER



- Molecule 4: DNA 11-MER





4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	106.36Å 106.36Å 132.65Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.05 – 3.75 44.22 – 3.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.05-3.75) 100.0 (44.22-3.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 3.66Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.267 , 0.329 0.261 , 0.329	Depositor DCC
R_{free} test set	463 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	142.7	Xtriage
Anisotropy	0.317	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 123.8	EDS
Estimated twinning fraction	0.147 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	1 of 9636 reflections (0.010%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3361	wwPDB-VP
Average B, all atoms (Å ²)	181.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	1/1175 (0.1%)	1.13	6/1596 (0.4%)
1	B	0.49	0/1109	1.16	10/1507 (0.7%)
2	C	0.68	0/706	1.44	7/1088 (0.6%)
3	D	0.62	0/288	1.37	2/442 (0.5%)
4	F	0.70	0/256	1.41	2/393 (0.5%)
All	All	0.56	1/3534 (0.0%)	1.26	27/5026 (0.5%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	86	TRP	CB-CG	-5.59	1.40	1.50

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	128	MET	CG-SD-CE	22.30	135.88	100.20
1	B	128	MET	CG-SD-CE	21.97	135.35	100.20
1	B	45	LYS	CD-CE-NZ	15.53	147.42	111.70
1	A	45	LYS	CD-CE-NZ	14.69	145.49	111.70
1	A	37	LYS	CD-CE-NZ	13.52	142.79	111.70
1	B	37	LYS	CD-CE-NZ	12.47	140.38	111.70
2	C	15	DC	O4'-C4'-C3'	-10.92	99.45	106.00
2	C	12	DG	O4'-C1'-N9	7.05	112.94	108.00
1	B	48	VAL	CG1-CB-CG2	7.01	122.12	110.90
2	C	14	DC	O4'-C1'-N1	6.92	112.84	108.00
1	A	54	ARG	CG-CD-NE	6.85	126.19	111.80
2	C	15	DC	O4'-C1'-N1	6.65	112.66	108.00
1	B	54	ARG	CD-NE-CZ	5.99	131.98	123.60
1	B	54	ARG	CA-CB-CG	-5.93	100.34	113.40
2	C	13	DA	O4'-C1'-N9	5.66	111.96	108.00
2	C	18	DT	N3-C4-O4	5.58	123.25	119.90
3	D	11	DT	N3-C4-O4	5.58	123.25	119.90
4	F	22	DG	C4'-C3'-C2'	-5.53	98.12	103.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	39	SER	N-CA-CB	-5.32	102.52	110.50
1	B	1	MET	CG-SD-CE	5.32	108.70	100.20
4	F	22	DG	O4'-C4'-C3'	-5.31	102.38	104.50
1	B	54	ARG	NE-CZ-NH1	-5.30	117.65	120.30
3	D	6	DC	O4'-C4'-C3'	-5.25	102.40	104.50
2	C	16	DT	O4'-C4'-C3'	-5.12	102.45	104.50
1	A	47	ARG	CB-CG-CD	5.11	124.88	111.60
1	A	54	ARG	CB-CG-CD	5.10	124.87	111.60
1	B	47	ARG	CG-CD-NE	-5.03	101.23	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1153	0	1158	26	1
1	B	1090	0	1050	21	0
2	C	631	0	352	25	1
3	D	259	0	150	7	0
4	F	228	0	124	13	0
All	All	3361	0	2834	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:14:DC:H42	2:C:20:DG:H1	1.12	0.96
2:C:9:DT:H3	4:F:25:DA:H61	1.28	0.79
3:D:12:DC:N3	4:F:22:DG:O6	2.18	0.77
2:C:14:DC:N4	2:C:20:DG:H1	1.85	0.73
3:D:12:DC:O2	4:F:22:DG:N1	2.21	0.66
2:C:12:DG:O6	2:C:22:DC:N3	2.31	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:9:DT:H3	4:F:25:DA:N6	1.96	0.62
1:A:40:PRO:HG3	2:C:28:DG:P	2.41	0.61
2:C:5:DC:H2''	2:C:6:DG:C8	2.37	0.60
1:A:92:LEU:HD13	1:B:48:VAL:HG11	1.82	0.60
2:C:5:DC:N3	4:F:29:DG:O6	2.36	0.58
1:A:80:LEU:HD11	2:C:26:DC:H4'	1.86	0.58
1:B:80:LEU:HD11	4:F:26:DG:H4'	1.84	0.58
2:C:24:DA:H2''	2:C:25:DT:H5'	1.87	0.57
1:A:108:MET:HE1	2:C:13:DA:OP1	2.05	0.56
3:D:11:DT:H2''	3:D:12:DC:H5'	1.87	0.56
3:D:1:DA:H2''	3:D:2:DT:H5'	1.88	0.56
2:C:12:DG:N1	2:C:22:DC:O2	2.39	0.55
1:B:5:GLY:HA3	1:B:151:ALA:HB2	1.89	0.55
1:A:5:GLY:HA3	1:A:151:ALA:HB2	1.88	0.55
1:A:72:GLN:NE2	2:C:25:DT:O3'	2.40	0.55
2:C:31:DG:H1	3:D:3:DC:H42	1.54	0.55
1:A:82:TYR:HE1	1:B:84:VAL:HG12	1.72	0.55
1:A:48:VAL:HG12	1:A:90:ALA:HB2	1.90	0.54
4:F:27:DC:H2''	4:F:28:DC:C6	2.43	0.53
1:B:80:LEU:CD1	4:F:27:DC:H5'	2.38	0.53
2:C:20:DG:H2'	2:C:21:DT:C6	2.45	0.52
1:A:24:GLY:HA2	1:A:25:ALA:C	2.30	0.51
1:A:2:VAL:O	1:A:65:GLU:HG2	2.12	0.49
2:C:26:DC:H2''	2:C:27:DG:C8	2.48	0.48
4:F:24:DA:H2''	4:F:25:DA:H5'	1.95	0.48
1:A:85:GLY:HA3	1:B:85:GLY:HA3	1.96	0.48
2:C:1:DC:H2'	2:C:2:DA:C8	2.48	0.48
3:D:12:DC:N3	4:F:22:DG:C6	2.80	0.48
1:B:42:GLU:HG2	1:B:46:GLU:HB3	1.96	0.48
1:A:80:LEU:CD1	2:C:27:DG:H5'	2.44	0.47
1:A:8:PRO:HA	1:A:13:LEU:HD12	1.97	0.47
1:B:80:LEU:HD12	4:F:27:DC:H5'	1.96	0.47
2:C:2:DA:H2'	2:C:3:DA:C8	2.50	0.47
1:B:140:ARG:HE	1:B:141:PRO:HA	1.79	0.46
1:B:8:PRO:HA	1:B:13:LEU:HD12	1.98	0.46
1:A:82:TYR:CE1	1:B:84:VAL:HG12	2.50	0.46
1:B:19:ALA:HB2	1:B:31:LEU:HD11	1.98	0.45
1:A:55:VAL:O	1:A:59:LEU:HG	2.16	0.45
1:A:5:GLY:CA	1:A:151:ALA:HB2	2.46	0.45
2:C:31:DG:H1	3:D:3:DC:N4	2.15	0.45
1:B:5:GLY:CA	1:B:151:ALA:HB2	2.47	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:10:ILE:HG22	4:F:27:DC:O5'	2.17	0.45
1:A:80:LEU:HD12	2:C:27:DG:H5'	1.99	0.44
1:A:28:ALA:O	1:A:133:LEU:HD23	2.18	0.43
1:B:28:ALA:O	1:B:133:LEU:HD23	2.18	0.43
1:A:19:ALA:HB2	1:A:31:LEU:HD11	2.00	0.43
1:A:27:LYS:HA	1:A:132:ILE:HG23	1.99	0.43
1:A:37:LYS:O	1:A:50:ARG:NH2	2.40	0.43
1:B:51:ILE:O	1:B:55:VAL:HG23	2.18	0.43
1:B:70:GLN:OE1	1:B:150:ILE:HD11	2.19	0.42
1:B:27:LYS:HA	1:B:132:ILE:HG23	2.01	0.42
1:A:1:MET:HB3	1:A:20:VAL:HB	2.00	0.42
1:B:114:LEU:O	1:B:128:MET:HG3	2.20	0.41
1:A:72:GLN:HB2	1:A:75:TYR:OH	2.20	0.41
2:C:5:DC:O2	4:F:29:DG:N1	2.46	0.41
2:C:4:DT:H1'	2:C:5:DC:H5'	2.03	0.41
1:B:44:ALA:O	1:B:48:VAL:HG12	2.22	0.40
1:A:70:GLN:OE1	1:A:150:ILE:HD11	2.21	0.40
1:A:40:PRO:HG3	2:C:28:DG:OP2	2.21	0.40
2:C:28:DG:C2	2:C:29:DC:C2	3.09	0.40
1:A:92:LEU:CD1	1:B:48:VAL:HG11	2.50	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:ARG:NH1	2:C:20:DG:OP1[4_655]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	152/169 (90%)	148 (97%)	3 (2%)	1 (1%)	26 72

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	151/169 (89%)	147 (97%)	2 (1%)	2 (1%)	15	61
All	All	303/338 (90%)	295 (97%)	5 (2%)	3 (1%)	19	66

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	78	ASN
1	B	78	ASN
1	B	165	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/123 (86%)	103 (97%)	3 (3%)	51	82
1	B	92/123 (75%)	87 (95%)	5 (5%)	27	68
All	All	198/246 (80%)	190 (96%)	8 (4%)	38	75

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	HIS
1	A	78	ASN
1	A	162	THR
1	B	48	VAL
1	B	60	HIS
1	B	73	PHE
1	B	140	ARG
1	B	162	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	158/169 (93%)	0.06	7 (4%) 38 25	89, 160, 266, 304	0
1	B	157/169 (92%)	-0.04	0 100 100	101, 164, 241, 306	0
2	C	31/31 (100%)	-0.68	1 (3%) 51 35	141, 183, 236, 243	0
3	D	13/13 (100%)	-0.35	0 100 100	153, 211, 254, 260	0
4	F	11/11 (100%)	-0.84	0 100 100	164, 207, 243, 252	0
All	All	370/393 (94%)	-0.09	8 (2%) 65 49	89, 168, 252, 306	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	115	ALA	3.0
1	A	123	GLU	2.6
2	C	25	DT	2.2
1	A	111	LYS	2.2
1	A	70	GLN	2.1
1	A	114	LEU	2.1
1	A	29	ARG	2.0
1	A	108	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.