



Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 01:05 AM BST

PDB ID : 2LFC
Title : Solution NMR Structure of Fumarate reductase flavoprotein subunit from Lactobacillus plantarum, Northeast Structural Genomics Consortium Target LpR145J
Authors : Liu, G.; Tang, Y.; Xiao, R.; Janjua, H.; Ciccocanti, C.; Wang, H.; Acton, T.; Everett, J.K.; Montelione, G.T.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2011-06-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 89%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	160	<div> <div style="width: 81%; background-color: green;"></div> <div style="width: 17%; background-color: cyan;"></div> <div style="width: 2%; background-color: yellow;"></div> <div style="width: 2%; background-color: orange;"></div> <div style="width: 2%; background-color: red;"></div> </div> <div>81% • 17%</div>

2 Ensemble composition and analysis

This entry contains 20 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:10-A:48, A:52-A:85, A:90-A:129, A:134-A:153 (133)	0.47	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 7, 8, 9, 10, 14, 15, 16
2	5, 6, 12, 13
3	11, 18, 20
4	4, 17, 19

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2584 atoms, of which 1275 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Fumarate reductase, flavoprotein subunit.

Mol	Chain	Residues	Atoms						Trace
1	A	160	Total	C	H	N	O	S	0
			2584	833	1275	226	247	3	

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INITIATING METHIONINE	UNP Q88SE0
A	153	LEU	-	EXPRESSION TAG	UNP Q88SE0
A	154	GLU	-	EXPRESSION TAG	UNP Q88SE0
A	155	HIS	-	EXPRESSION TAG	UNP Q88SE0
A	156	HIS	-	EXPRESSION TAG	UNP Q88SE0
A	157	HIS	-	EXPRESSION TAG	UNP Q88SE0
A	158	HIS	-	EXPRESSION TAG	UNP Q88SE0
A	159	HIS	-	EXPRESSION TAG	UNP Q88SE0
A	160	HIS	-	EXPRESSION TAG	UNP Q88SE0

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.2 Score per residue for model 2

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.3 Score per residue for model 3

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.4 Score per residue for model 4

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.5 Score per residue for model 5 (medoid)

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.6 Score per residue for model 6

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.7 Score per residue for model 7

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.8 Score per residue for model 8

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.9 Score per residue for model 9

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.10 Score per residue for model 10

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.11 Score per residue for model 11


- Molecule 1: Fumarate reductase, flavoprotein subunit



4.2.12 Score per residue for model 12


- Molecule 1: Fumarate reductase, flavoprotein subunit






4.2.13 Score per residue for model 13

- Molecule 1: Fumarate reductase, flavoprotein subunit





4.2.14 Score per residue for model 14

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.15 Score per residue for model 15

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.16 Score per residue for model 16

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.17 Score per residue for model 17

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.18 Score per residue for model 18

- Molecule 1: Fumarate reductase, flavoprotein subunit




4.2.19 Score per residue for model 19

- Molecule 1: Fumarate reductase, flavoprotein subunit



4.2.20 Score per residue for model 20

- Molecule 1: Fumarate reductase, flavoprotein subunit



5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry, molecular dynamics, simulated annealing, torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
CNS	structure solution	
CNS	geometry optimization	
CYANA	refinement	3.0
CYANA	geometry optimization	3.0
CYANA	structure solution	3.0
AutoStructure	refinement	2.1
TALOS+	geometry optimization	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2lfc_cs.str
Number of chemical shift lists	1
Total number of shifts	1899
Number of shifts mapped to atoms	1899
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	89%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1089	1066	1062	2±2
All	All	21780	21320	21240	46

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:31:ARG:HD2	1:A:115:THR:HG23	0.51	1.82	4	1
1:A:58:MET:SD	1:A:147:TYR:HB2	0.48	2.49	6	1
1:A:122:TYR:HB2	1:A:127:HIS:O	0.47	2.10	4	8
1:A:32:ILE:HB	1:A:45:ALA:HB1	0.47	1.87	2	1
1:A:31:ARG:NE	1:A:31:ARG:H	0.45	2.08	11	1
1:A:20:MET:SD	1:A:65:LYS:HG3	0.45	2.51	2	2
1:A:76:THR:HG22	1:A:79:GLU:HG2	0.45	1.88	14	2
1:A:112:LEU:HA	1:A:115:THR:HG22	0.45	1.88	11	1
1:A:26:ASN:HD21	1:A:30:ASP:HB3	0.45	1.72	6	1
1:A:123:VAL:HG22	1:A:138:LEU:HB3	0.44	1.90	12	4
1:A:90:ARG:NH1	1:A:106:GLY:HA3	0.44	2.28	3	1
1:A:58:MET:SD	1:A:149:ILE:HG12	0.44	2.53	11	1
1:A:119:TYR:O	1:A:123:VAL:HG23	0.43	2.14	12	9
1:A:11:THR:O	1:A:15:LYS:HG2	0.43	2.13	7	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:23:ILE:HD12	1:A:31:ARG:NH2	0.43	2.28	19	1
1:A:134:ASP:HB3	1:A:137:TYR:HD1	0.42	1.73	7	1
1:A:99:GLU:O	1:A:103:GLU:HG3	0.42	2.14	11	3
1:A:28:LYS:O	1:A:108:VAL:HB	0.41	2.14	10	1
1:A:81:LYS:O	1:A:85:GLU:HG2	0.41	2.15	4	1
1:A:124:GLN:HB3	1:A:140:GLN:NE2	0.41	2.31	18	1
1:A:26:ASN:HD21	1:A:30:ASP:HB2	0.41	1.76	19	1
1:A:119:TYR:HA	1:A:122:TYR:CE2	0.41	2.51	18	1
1:A:31:ARG:HD3	1:A:31:ARG:H	0.40	1.74	4	1
1:A:11:THR:HB	1:A:15:LYS:HE2	0.40	1.92	11	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/160 (83%)	121±2 (91±1%)	10±2 (8±1%)	2±1 (1±1%)	20	66
All	All	2660/3200 (83%)	2423 (91%)	202 (8%)	35 (1%)	20	66

All 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	74	ASP	16
1	A	21	GLY	7
1	A	75	PHE	3
1	A	143	GLY	2
1	A	31	ARG	2
1	A	37	ASN	2
1	A	29	GLY	1
1	A	34	ASN	1
1	A	35	GLU	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	118/140 (84%)	115±2 (98±1%)	3±2 (2±1%)	61 93
All	All	2360/2800 (84%)	2305 (98%)	55 (2%)	61 93

All 18 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	31	ARG	13
1	A	122	TYR	7
1	A	81	LYS	5
1	A	75	PHE	5
1	A	128	ASP	3
1	A	34	ASN	3
1	A	117	LYS	3
1	A	35	GLU	2
1	A	152	ARG	2
1	A	44	ASN	2
1	A	115	THR	2
1	A	61	ARG	2
1	A	24	TYR	1
1	A	144	GLU	1
1	A	125	ASP	1
1	A	142	GLU	1
1	A	118	ASN	1
1	A	150	GLU	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 89% for the well-defined parts and 83% for the entire structure.

7.1 Chemical shift list 1

File name: 2lfc_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1899
Number of shifts mapped to atoms	1899
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	150	0.12 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	141	0.48 ± 0.10	None needed (< 0.5 ppm)
$^{13}\text{C}'$	128	0.08 ± 0.08	None needed (< 0.5 ppm)
^{15}N	130	0.64 ± 0.40	None needed (imprecise)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 89%, i.e. 1500 atoms were assigned a chemical shift out of a possible 1683. 22 out of 22 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	614/659 (93%)	249/263 (95%)	249/266 (94%)	116/130 (89%)
Sidechain	731/845 (87%)	451/491 (92%)	270/317 (85%)	10/37 (27%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	155/179 (87%)	82/93 (88%)	72/77 (94%)	1/9 (11%)
Overall	1500/1683 (89%)	782/847 (92%)	591/660 (90%)	127/176 (72%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 83%, i.e. 1691 atoms were assigned a chemical shift out of a possible 2032. 24 out of 25 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	690/792 (87%)	282/316 (89%)	278/320 (87%)	130/156 (83%)
Sidechain	835/1004 (83%)	518/586 (88%)	305/373 (82%)	12/45 (27%)
Aromatic	166/236 (70%)	88/122 (72%)	77/93 (83%)	1/21 (5%)
Overall	1691/2032 (83%)	888/1024 (87%)	660/786 (84%)	143/222 (64%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

