



# Full wwPDB X-ray Structure Validation Report ⓘ

May 30, 2016 – 12:32 PM EDT

PDB ID : 4LHK  
Title : Structure of the N-terminal domain of the Lg-Flo1 adhesin (N-Lg-Flo1p) from the yeast *Saccharomyces pastorianus*, in complex with calcium and alpha-1,2-mannobiose  
Authors : Ielasi, F.S.; Willaert, R.G.  
Deposited on : 2013-07-01  
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027674  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027674


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2417 (1.76-1.72)
Clashscore	102246	2570 (1.76-1.72)
Ramachandran outliers	100387	2544 (1.76-1.72)
Sidechain outliers	100360	2544 (1.76-1.72)
RSRZ outliers	91569	2420 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	239	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>9%</div> <div>9%</div> </div> </div>
1	B	239	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>8%</div> <div>9%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MAN	A	302	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Flocculin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	0	0
			1686	1081	256	341	8			
1	B	217	Total	C	N	O	S	0	0	0
			1686	1081	256	341	8			

There are 98 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	58	GLN	PRO	SEE REMARK 999	UNP B3IUB3
A	214	THR	-	SEE REMARK 999	UNP B3IUB3
A	215	GLU	-	SEE REMARK 999	UNP B3IUB3
A	216	VAL	-	SEE REMARK 999	UNP B3IUB3
A	217	ASN	-	SEE REMARK 999	UNP B3IUB3
A	218	ASP	-	SEE REMARK 999	UNP B3IUB3
A	219	ASP	-	SEE REMARK 999	UNP B3IUB3
A	220	PHE	-	SEE REMARK 999	UNP B3IUB3
A	221	GLU	-	SEE REMARK 999	UNP B3IUB3
A	222	GLY	-	SEE REMARK 999	UNP B3IUB3
A	223	TYR	-	SEE REMARK 999	UNP B3IUB3
A	224	VAL	-	SEE REMARK 999	UNP B3IUB3
A	225	TYR	-	SEE REMARK 999	UNP B3IUB3
A	226	SER	-	SEE REMARK 999	UNP B3IUB3
A	227	PHE	-	SEE REMARK 999	UNP B3IUB3
A	228	ASP	-	SEE REMARK 999	UNP B3IUB3
A	229	ASP	-	SEE REMARK 999	UNP B3IUB3
A	230	ASP	-	SEE REMARK 999	UNP B3IUB3
A	231	LEU	-	SEE REMARK 999	UNP B3IUB3
A	232	SER	-	SEE REMARK 999	UNP B3IUB3
A	233	GLN	-	SEE REMARK 999	UNP B3IUB3
A	234	SER	-	SEE REMARK 999	UNP B3IUB3
A	235	ASN	-	SEE REMARK 999	UNP B3IUB3
A	236	CYS	-	SEE REMARK 999	UNP B3IUB3
A	237	THR	-	SEE REMARK 999	UNP B3IUB3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	238	ILE	-	SEE REMARK 999	UNP B3IUB3
A	239	PRO	-	SEE REMARK 999	UNP B3IUB3
A	240	ASP	-	SEE REMARK 999	UNP B3IUB3
A	241	PRO	-	SEE REMARK 999	UNP B3IUB3
A	242	SER	-	SEE REMARK 999	UNP B3IUB3
A	243	LYS	-	SEE REMARK 999	UNP B3IUB3
A	244	HIS	-	SEE REMARK 999	UNP B3IUB3
A	245	THR	-	SEE REMARK 999	UNP B3IUB3
A	246	THR	-	SEE REMARK 999	UNP B3IUB3
A	247	SER	-	SEE REMARK 999	UNP B3IUB3
A	248	LEU	-	EXPRESSION TAG	UNP B3IUB3
A	249	GLU	-	EXPRESSION TAG	UNP B3IUB3
A	250	VAL	-	EXPRESSION TAG	UNP B3IUB3
A	251	LEU	-	EXPRESSION TAG	UNP B3IUB3
A	252	PHE	-	EXPRESSION TAG	UNP B3IUB3
A	253	GLN	-	EXPRESSION TAG	UNP B3IUB3
A	254	GLY	-	EXPRESSION TAG	UNP B3IUB3
A	255	PRO	-	EXPRESSION TAG	UNP B3IUB3
A	256	HIS	-	EXPRESSION TAG	UNP B3IUB3
A	257	HIS	-	EXPRESSION TAG	UNP B3IUB3
A	258	HIS	-	EXPRESSION TAG	UNP B3IUB3
A	259	HIS	-	EXPRESSION TAG	UNP B3IUB3
A	260	HIS	-	EXPRESSION TAG	UNP B3IUB3
A	261	HIS	-	EXPRESSION TAG	UNP B3IUB3
B	58	GLN	PRO	SEE REMARK 999	UNP B3IUB3
B	214	THR	-	SEE REMARK 999	UNP B3IUB3
B	215	GLU	-	SEE REMARK 999	UNP B3IUB3
B	216	VAL	-	SEE REMARK 999	UNP B3IUB3
B	217	ASN	-	SEE REMARK 999	UNP B3IUB3
B	218	ASP	-	SEE REMARK 999	UNP B3IUB3
B	219	ASP	-	SEE REMARK 999	UNP B3IUB3
B	220	PHE	-	SEE REMARK 999	UNP B3IUB3
B	221	GLU	-	SEE REMARK 999	UNP B3IUB3
B	222	GLY	-	SEE REMARK 999	UNP B3IUB3
B	223	TYR	-	SEE REMARK 999	UNP B3IUB3
B	224	VAL	-	SEE REMARK 999	UNP B3IUB3
B	225	TYR	-	SEE REMARK 999	UNP B3IUB3
B	226	SER	-	SEE REMARK 999	UNP B3IUB3
B	227	PHE	-	SEE REMARK 999	UNP B3IUB3
B	228	ASP	-	SEE REMARK 999	UNP B3IUB3
B	229	ASP	-	SEE REMARK 999	UNP B3IUB3
B	230	ASP	-	SEE REMARK 999	UNP B3IUB3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	231	LEU	-	SEE REMARK 999	UNP B3IUB3
B	232	SER	-	SEE REMARK 999	UNP B3IUB3
B	233	GLN	-	SEE REMARK 999	UNP B3IUB3
B	234	SER	-	SEE REMARK 999	UNP B3IUB3
B	235	ASN	-	SEE REMARK 999	UNP B3IUB3
B	236	CYS	-	SEE REMARK 999	UNP B3IUB3
B	237	THR	-	SEE REMARK 999	UNP B3IUB3
B	238	ILE	-	SEE REMARK 999	UNP B3IUB3
B	239	PRO	-	SEE REMARK 999	UNP B3IUB3
B	240	ASP	-	SEE REMARK 999	UNP B3IUB3
B	241	PRO	-	SEE REMARK 999	UNP B3IUB3
B	242	SER	-	SEE REMARK 999	UNP B3IUB3
B	243	LYS	-	SEE REMARK 999	UNP B3IUB3
B	244	HIS	-	SEE REMARK 999	UNP B3IUB3
B	245	THR	-	SEE REMARK 999	UNP B3IUB3
B	246	THR	-	SEE REMARK 999	UNP B3IUB3
B	247	SER	-	SEE REMARK 999	UNP B3IUB3
B	248	LEU	-	EXPRESSION TAG	UNP B3IUB3
B	249	GLU	-	EXPRESSION TAG	UNP B3IUB3
B	250	VAL	-	EXPRESSION TAG	UNP B3IUB3
B	251	LEU	-	EXPRESSION TAG	UNP B3IUB3
B	252	PHE	-	EXPRESSION TAG	UNP B3IUB3
B	253	GLN	-	EXPRESSION TAG	UNP B3IUB3
B	254	GLY	-	EXPRESSION TAG	UNP B3IUB3
B	255	PRO	-	EXPRESSION TAG	UNP B3IUB3
B	256	HIS	-	EXPRESSION TAG	UNP B3IUB3
B	257	HIS	-	EXPRESSION TAG	UNP B3IUB3
B	258	HIS	-	EXPRESSION TAG	UNP B3IUB3
B	259	HIS	-	EXPRESSION TAG	UNP B3IUB3
B	260	HIS	-	EXPRESSION TAG	UNP B3IUB3
B	261	HIS	-	EXPRESSION TAG	UNP B3IUB3

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	2	Total	C	O	0	0
			23	12	11		
2	B	2	Total	C	O	0	0
			23	12	11		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Ca 1	0	0
3	A	1	Total 1	Ca 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	96	Total 96	O 96	0	0
4	B	156	Total 156	O 156	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.62Å 85.48Å 68.24Å 90.00° 93.72° 90.00°	Depositor
Resolution (Å)	35.13 – 1.73 35.13 – 1.73	Depositor EDS
% Data completeness (in resolution range)	97.1 (35.13-1.73) 97.1 (35.13-1.73)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.13 (at 1.73Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.137 , 0.193 0.150 , 0.200	Depositor DCC
$R_{free}$ test set	2275 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.1	Xtriage
Anisotropy	0.424	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 41.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3672	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.93	0/1736	0.89	1/2375 (0.0%)
1	B	0.94	2/1736 (0.1%)	0.91	0/2375
All	All	0.94	2/3472 (0.1%)	0.90	1/4750 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	59	TYR	CE1-CZ	-5.32	1.31	1.38
1	B	74	SER	CB-OG	-5.11	1.35	1.42

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	129	PHE	CB-CG-CD2	-5.62	116.86	120.80

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1686	0	1578	12	0
1	B	1686	0	1578	15	0
2	A	23	0	19	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	23	0	20	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	96	0	0	2	0
4	B	156	0	0	5	0
All	All	3672	0	3195	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:181:THR:HG22	4:B:492:HOH:O	1.63	0.96
1:A:67:THR:HG22	4:A:493:HOH:O	1.93	0.67
1:A:128:LYS:HD2	1:A:176:ASP:OD1	1.93	0.66
1:B:36:LYS:HE3	1:B:228:ASP:OD1	2.00	0.62
1:B:181:THR:CG2	4:B:492:HOH:O	2.32	0.60
1:B:59:TYR:HA	1:B:63:LYS:HB2	1.84	0.59
1:B:81:ILE:HD13	1:B:195:TYR:CD2	2.42	0.55
1:A:59:TYR:HA	1:A:63:LYS:HB2	1.92	0.50
2:A:301:MAN:H1	2:A:302:MAN:C5	2.43	0.49
1:A:68:LYS:O	4:A:492:HOH:O	2.20	0.47
1:B:119:PRO:HD2	1:B:187:TYR:O	2.15	0.47
1:B:37:ASN:O	1:B:75:GLY:HA2	2.15	0.47
1:A:136:ALA:HA	1:A:194:VAL:O	2.15	0.46
1:A:167:LYS:HE3	2:A:302:MAN:H3	1.97	0.46
1:A:119:PRO:HD2	1:A:187:TYR:O	2.16	0.46
2:A:301:MAN:H1	2:A:302:MAN:O5	2.15	0.46
1:B:36:LYS:HE3	4:B:515:HOH:O	2.18	0.43
1:B:42:ASN:OD1	1:B:69:LYS:HD3	2.19	0.43
1:B:133:ASP:HA	1:B:134:ASP:HA	1.86	0.42
1:B:36:LYS:HD2	4:B:515:HOH:O	2.18	0.42
1:B:67:THR:OG1	1:B:68:LYS:N	2.52	0.42
1:A:133:ASP:HA	1:A:134:ASP:HA	1.90	0.41
1:A:84:ASP:OD1	1:A:201:LEU:CD2	2.69	0.41
1:A:59:TYR:CE1	1:A:64:TYR:HA	2.56	0.41
1:B:42:ASN:CG	1:B:69:LYS:HD3	2.41	0.41
2:B:302:MAN:H2	4:B:540:HOH:O	2.22	0.41
1:B:59:TYR:CE1	1:B:64:TYR:HA	2.56	0.41
1:B:83:TYR:HB3	1:B:202:ALA:HB3	2.04	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:PRO:HG3	1:A:183:MET:HB3	2.04	0.40
1:A:39:MET:HB3	1:A:114:THR:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/239 (90%)	208 (97%)	7 (3%)	0	100	100
1	B	215/239 (90%)	205 (95%)	10 (5%)	0	100	100
All	All	430/478 (90%)	413 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/207 (90%)	186 (100%)	1 (0%)	92	87
1	B	187/207 (90%)	187 (100%)	0	100	100
All	All	374/414 (90%)	373 (100%)	1 (0%)	94	91

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	234	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	A	58	GLN
1	A	76	GLN
1	A	121	GLN
1	B	37	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MAN	A	301	3,2	12,12,12	1.11	2 (16%)	17,17,17	1.13	1 (5%)
2	MAN	A	302	2	11,11,12	0.90	0	15,15,17	1.52	4 (26%)
2	MAN	B	301	3,2	12,12,12	0.42	0	17,17,17	0.50	0
2	MAN	B	302	2	11,11,12	0.25	0	15,15,17	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAN	A	301	3,2	-	0/2/22/22	0/1/1/1
2	MAN	A	302	2	-	0/2/19/22	0/1/1/1
2	MAN	B	301	3,2	-	0/2/22/22	0/1/1/1
2	MAN	B	302	2	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	MAN	O2-C2	-2.35	1.37	1.43
2	A	301	MAN	O4-C4	-2.07	1.38	1.43

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	302	MAN	O2-C2-C1	-3.01	103.21	109.23
2	A	302	MAN	O5-C1-C2	-2.47	106.94	110.89
2	A	302	MAN	C1-C2-C3	-2.24	106.84	109.55
2	A	302	MAN	O3-C3-C2	-2.08	106.20	110.01
2	A	301	MAN	O1-C1-O5	2.10	116.18	110.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	MAN	2	0
2	A	302	MAN	3	0
2	B	302	MAN	1	0

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	217/239 (90%)	0.09	6 (2%) 56 62	13, 21, 36, 44	0
1	B	217/239 (90%)	0.13	7 (3%) 51 57	13, 20, 34, 55	0
All	All	434/478 (90%)	0.11	13 (2%) 54 60	13, 21, 35, 55	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	240	ASP	4.2
1	B	239	PRO	3.5
1	B	241	PRO	2.9
1	B	242	SER	2.7
1	A	206	VAL	2.6
1	A	241	PRO	2.6
1	A	240	ASP	2.5
1	B	193	ILE	2.5
1	A	242	SER	2.5
1	A	204	LEU	2.2
1	B	206	VAL	2.1
1	A	182	TYR	2.1
1	B	175	THR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.



The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	MAN	A	302	11/12	0.89	0.12	3.31	25,29,38,41	0
2	MAN	B	301	12/12	0.93	0.11	1.03	13,18,22,23	0
2	MAN	A	301	12/12	0.97	0.09	0.80	13,15,22,23	0
2	MAN	B	302	11/12	0.94	0.17	-	27,33,41,50	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	CA	A	303	1/1	1.00	0.09	0.13	15,15,15,15	0
3	CA	B	303	1/1	1.00	0.07	-0.29	16,16,16,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.