



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:26 AM GMT

PDB ID : 3LSI
Title : Pyranose 2-oxidase T169A, tetragonal
Authors : Tan, T.C.; Spadiut, O.; Divne, C.
Deposited on : 2010-02-12
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	623	
1	B	623	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pyranose 2-oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	575	Total	C	N	O	S	0	0	0
			4532	2862	776	870	24			
1	B	576	Total	C	N	O	S	0	0	0
			4540	2867	777	871	25			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	169	ALA	THR	ENGINEERED	UNP Q7ZA32
B	169	ALA	THR	ENGINEERED	UNP Q7ZA32

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	507	Total	O	0	0
			507	507		
3	B	497	Total	O	0	0
			497	497		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyranose 2-oxidase



• Molecule 1: Pyranose 2-oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	101.30 Å 101.30 Å 249.77 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.10 – 1.90 29.11 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.10-1.90) 99.6 (29.11-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 1.91 Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, R_{free}	0.164 , 0.213 0.178 , 0.219	Depositor DCC
R_{free} test set	3090 reflections (3.11%)	DCC
Wilson B-factor (Å ²)	16.8	Xtriage
Anisotropy	0.215	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 102585 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10182	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.81 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.3323e-03.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.03	5/4648 (0.1%)	0.95	13/6320 (0.2%)
1	B	1.02	5/4656 (0.1%)	0.94	9/6330 (0.1%)
All	All	1.02	10/9304 (0.1%)	0.94	22/12650 (0.2%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	310	GLU	CB-CG	7.82	1.67	1.52
1	B	310	GLU	CG-CD	7.50	1.63	1.51
1	A	310	GLU	CG-CD	7.47	1.63	1.51
1	B	310	GLU	CB-CG	7.20	1.65	1.52
1	A	482	GLU	CG-CD	6.74	1.62	1.51
1	B	371	GLU	CB-CG	-5.95	1.40	1.52
1	A	478	GLU	CD-OE2	5.92	1.32	1.25
1	B	61	TYR	CD1-CE1	5.50	1.47	1.39
1	A	408	TRP	CE3-CZ3	5.17	1.47	1.38
1	B	576	LYS	CE-NZ	5.09	1.61	1.49

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	470	ASP	CB-CG-OD1	8.09	125.58	118.30
1	B	383	ARG	NE-CZ-NH1	-7.46	116.57	120.30
1	B	383	ARG	NE-CZ-NH2	7.42	124.01	120.30
1	A	112	MET	CG-SD-CE	-7.41	88.34	100.20
1	B	472	ARG	NE-CZ-NH2	-6.98	116.81	120.30
1	B	470	ASP	CB-CG-OD2	-6.95	112.05	118.30
1	A	139	ARG	NE-CZ-NH2	-6.82	116.89	120.30
1	A	521	ASP	CB-CG-OD2	6.31	123.98	118.30
1	A	82	SER	CB-CA-C	-6.30	98.12	110.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	81	ASP	CB-CG-OD1	-6.13	112.78	118.30
1	A	569	ASP	CB-CG-OD1	-5.65	113.22	118.30
1	A	48	ASP	CB-CG-OD1	5.61	123.35	118.30
1	B	261	ASP	CB-CG-OD1	5.53	123.28	118.30
1	B	406	ASP	CB-CG-OD1	5.48	123.23	118.30
1	A	234	ARG	NE-CZ-NH1	-5.44	117.58	120.30
1	B	192	ASP	CB-CG-OD1	5.41	123.17	118.30
1	A	245	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	B	310	GLU	OE1-CD-OE2	-5.29	116.96	123.30
1	A	174	ARG	NE-CZ-NH2	-5.20	117.70	120.30
1	A	139	ARG	CG-CD-NE	-5.14	101.00	111.80
1	A	201	LYS	CD-CE-NZ	-5.11	99.96	111.70
1	A	147	ASP	CB-CG-OD1	5.05	122.85	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4532	0	4377	52	0
1	B	4540	0	4386	50	0
2	A	53	0	30	1	0
2	B	53	0	30	3	0
3	A	507	0	0	13	0
3	B	497	0	0	14	0
All	All	10182	0	8823	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:460:GLN:HB3	3:A:1821:HOH:O	1.57	1.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:459:VAL:HA	3:A:1442:HOH:O	1.61	1.00
1:B:101:ASP:O	1:B:104:VAL:HG12	1.65	0.96
1:A:459:VAL:HG12	1:A:460:GLN:O	1.66	0.94
1:A:132:GLN:HG2	3:A:1850:HOH:O	1.65	0.94
1:B:365:GLN:HE22	1:B:459:VAL:HG22	1.38	0.85
1:B:132:GLN:HG2	3:B:1822:HOH:O	1.80	0.82
1:A:478:GLU:HB3	3:A:1931:HOH:O	1.78	0.82
1:B:132:GLN:CG	3:B:1822:HOH:O	2.29	0.80
1:B:457:GLY:HA3	3:B:1373:HOH:O	1.82	0.80
1:A:45:ILE:CG2	1:A:45:ILE:O	2.34	0.75
1:B:382:ILE:HD13	3:B:1450:HOH:O	1.86	0.74
1:A:132:GLN:CG	3:A:1850:HOH:O	2.29	0.70
1:A:45:ILE:O	1:A:45:ILE:HG22	1.91	0.69
1:A:167:HIS:CD2	2:A:801:FAD:HM82	2.27	0.69
1:B:225:LYS:HE2	3:B:1650:HOH:O	1.94	0.68
1:A:285:ARG:NH1	1:A:299:HIS:HD2	1.92	0.68
1:B:576:LYS:HE2	3:B:1942:HOH:O	1.93	0.68
1:A:459:VAL:HG12	1:A:460:GLN:N	2.10	0.66
1:B:167:HIS:CD2	2:B:801:FAD:HM82	2.31	0.66
1:A:167:HIS:ND1	3:A:1300:HOH:O	2.28	0.66
1:A:44:ASP:HB2	1:A:71:LYS:HZ3	1.64	0.63
1:A:404:HIS:HE1	3:A:1385:HOH:O	1.81	0.63
1:B:218:ARG:HD2	3:B:1026:HOH:O	1.99	0.61
1:B:404:HIS:HE1	3:B:1539:HOH:O	1.83	0.61
1:B:246:ARG:HG3	1:B:246:ARG:O	2.01	0.60
1:B:299:HIS:HB2	1:B:310:GLU:OE2	2.02	0.60
1:A:63:ARG:HD3	1:A:259:VAL:O	2.02	0.59
1:A:285:ARG:NH1	1:A:299:HIS:CD2	2.70	0.58
1:A:230:TYR:O	1:A:233:GLN:HG3	2.04	0.57
1:A:403:LYS:HE2	3:A:1973:HOH:O	2.04	0.56
1:B:341:ASN:HD22	1:B:342:PRO:HD2	1.72	0.55
1:A:81:ASP:O	1:B:81:ASP:O	2.25	0.54
1:A:450:HIS:O	1:A:470:ASP:HB2	2.07	0.54
1:B:201:LYS:HE2	1:B:205:TYR:OH	2.07	0.53
1:A:285:ARG:HH12	1:A:299:HIS:CD2	2.26	0.53
1:A:89:HIS:CE1	1:A:91:LYS:HB2	2.44	0.53
1:A:44:ASP:HB2	1:A:71:LYS:NZ	2.23	0.52
1:B:285:ARG:NH2	3:B:1974:HOH:O	2.31	0.52
1:B:157:VAL:HG21	1:B:324:HIS:HE1	1.75	0.52
1:A:459:VAL:CG1	1:A:460:GLN:N	2.72	0.52
1:B:132:GLN:NE2	3:B:1822:HOH:O	2.38	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:558:ASP:OD2	1:A:561:GLU:HG3	2.10	0.51
1:A:158:THR:HG22	1:A:160:VAL:HG22	1.93	0.51
1:B:471:TRP:CH2	1:B:526:SER:HA	2.45	0.51
1:B:81:ASP:OD1	1:B:81:ASP:N	2.42	0.50
1:B:576:LYS:CE	3:B:1942:HOH:O	2.56	0.50
1:A:47:TYR:O	1:A:313:ALA:HA	2.11	0.49
1:B:47:TYR:O	1:B:313:ALA:HA	2.12	0.49
1:B:167:HIS:HD2	2:B:801:FAD:HM82	1.77	0.49
1:B:285:ARG:NE	3:B:1974:HOH:O	2.25	0.48
1:B:576:LYS:HD2	1:B:576:LYS:N	2.28	0.48
1:A:218:ARG:HD2	3:A:1032:HOH:O	2.14	0.47
1:B:432:GLU:HB2	1:B:433:PRO:HD2	1.97	0.47
1:A:89:HIS:ND1	1:A:91:LYS:HB2	2.30	0.47
1:A:461:GLN:HG3	3:B:1832:HOH:O	2.14	0.46
1:B:89:HIS:CE1	1:B:91:LYS:HB2	2.50	0.46
1:B:328:LEU:HD23	1:B:328:LEU:O	2.15	0.46
1:B:444:PRO:HD2	1:B:445:TRP:CZ3	2.51	0.46
1:A:223:LEU:HD13	1:A:236:PHE:HB3	1.97	0.46
1:A:56:PRO:HD3	1:A:165:SER:HB3	1.97	0.46
1:A:223:LEU:CD1	1:A:236:PHE:HB3	2.46	0.46
1:B:50:VAL:HG13	1:B:313:ALA:HB2	1.97	0.45
1:B:137:PHE:CE2	1:B:139:ARG:HG3	2.51	0.45
1:A:403:LYS:CE	3:A:1973:HOH:O	2.61	0.45
1:B:299:HIS:CB	1:B:310:GLU:OE2	2.64	0.45
1:B:218:ARG:HG3	1:B:430:ASP:OD2	2.17	0.45
1:A:336:GLN:NE2	1:A:344:ASN:O	2.50	0.45
1:A:288:ARG:HD2	1:A:292:ASN:OD1	2.17	0.45
1:B:489:ASP:O	1:B:490:LYS:HD3	2.17	0.45
1:A:91:LYS:NZ	3:A:1361:HOH:O	2.43	0.44
1:B:288:ARG:HD2	1:B:292:ASN:OD1	2.16	0.44
1:B:46:LYS:HD3	1:B:47:TYR:N	2.33	0.44
1:A:346:PRO:HG2	1:A:350:PRO:HA	1.99	0.44
1:A:459:VAL:O	3:A:1649:HOH:O	2.21	0.44
1:A:218:ARG:HG3	1:A:430:ASP:OD2	2.18	0.44
1:A:461:GLN:CB	3:B:1832:HOH:O	2.66	0.44
1:B:570:SER:HB3	1:B:580:LEU:O	2.18	0.44
1:A:324:HIS:HD2	1:A:327:GLN:OE1	2.02	0.43
1:A:293:SER:HA	1:A:574:GLY:O	2.18	0.43
1:B:489:ASP:OD1	1:B:490:LYS:HG2	2.18	0.43
1:A:230:TYR:O	1:A:231:LYS:C	2.57	0.43
1:B:349:LEU:N	1:B:350:PRO:CD	2.81	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:ILE:HG23	1:A:45:ILE:O	2.16	0.43
1:B:101:ASP:O	1:B:104:VAL:CG1	2.51	0.43
1:A:178:GLU:O	1:A:178:GLU:HG3	2.18	0.43
1:A:44:ASP:CB	1:A:71:LYS:NZ	2.82	0.43
1:A:219:HIS:HB2	1:A:433:PRO:HA	2.01	0.43
1:B:408:TRP:C	1:B:408:TRP:CD1	2.93	0.42
1:B:341:ASN:HD22	1:B:342:PRO:CD	2.31	0.42
1:B:45:ILE:N	1:B:45:ILE:HD13	2.35	0.42
1:B:433:PRO:O	1:B:450:HIS:HA	2.19	0.42
1:A:404:HIS:CE1	3:A:1385:HOH:O	2.62	0.42
1:B:159:ARG:HA	2:B:801:FAD:O2B	2.19	0.42
1:A:47:TYR:CD2	1:A:73:ALA:HB2	2.55	0.42
1:A:81:ASP:C	1:A:81:ASP:OD1	2.59	0.41
1:B:478:GLU:HG3	1:B:511:THR:OG1	2.21	0.41
1:B:81:ASP:O	1:B:83:GLY:N	2.45	0.40
1:B:110:GLN:HG3	1:B:158:THR:HG23	2.03	0.40
1:A:471:TRP:CH2	1:A:526:SER:HA	2.56	0.40
1:B:340:PRO:HD3	1:B:486:TRP:CG	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	573/623 (92%)	552 (96%)	20 (4%)	1 (0%)	52	42
1	B	574/623 (92%)	557 (97%)	15 (3%)	2 (0%)	46	35
All	All	1147/1246 (92%)	1109 (97%)	35 (3%)	3 (0%)	46	35

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	82	SER
1	B	458	ALA
1	A	45	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	502/541 (93%)	486 (97%)	16 (3%)	46	35
1	B	503/541 (93%)	489 (97%)	14 (3%)	51	41
All	All	1005/1082 (93%)	975 (97%)	30 (3%)	48	38

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	ASP
1	A	91	LYS
1	A	168	TRP
1	A	178	GLU
1	A	206	PHE
1	A	223	LEU
1	A	235	ASP
1	A	286	VAL
1	A	341	ASN
1	A	385	THR
1	A	408	TRP
1	A	450	HIS
1	A	460	GLN
1	A	516	GLU
1	A	593	ASN
1	A	618	PHE
1	B	45	ILE
1	B	168	TRP
1	B	185	LYS
1	B	206	PHE
1	B	211	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	246	ARG
1	B	336	GLN
1	B	341	ASN
1	B	408	TRP
1	B	450	HIS
1	B	460	GLN
1	B	560	LYS
1	B	576	LYS
1	B	593	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	299	HIS
1	A	324	HIS
1	A	341	ASN
1	A	404	HIS
1	B	324	HIS
1	B	341	ASN
1	B	344	ASN
1	B	404	HIS
1	B	460	GLN
1	B	461	GLN
1	B	611	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FAD	A	801	-	48,58,58	1.26	8 (16%)	54,89,89	3.36	18 (33%)
2	FAD	B	801	-	48,58,58	1.12	2 (4%)	54,89,89	3.15	15 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	801	-	-	0/30/50/50	0/6/6/6
2	FAD	B	801	-	-	0/30/50/50	0/6/6/6

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	FAD	O2B-C2B	-2.16	1.37	1.43
2	A	801	FAD	C4-N3	2.19	1.37	1.33
2	A	801	FAD	C1'-N10	2.20	1.50	1.48
2	A	801	FAD	C2A-N3A	2.24	1.36	1.32
2	A	801	FAD	C2A-N1A	2.29	1.38	1.33
2	A	801	FAD	C5'-C4'	2.62	1.55	1.51
2	A	801	FAD	C8M-C8	2.75	1.56	1.51
2	B	801	FAD	C2A-N3A	3.27	1.38	1.32
2	A	801	FAD	C4-C4X	3.80	1.48	1.41
2	B	801	FAD	C4-C4X	3.91	1.49	1.41

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	FAD	N3A-C2A-N1A	-12.22	119.54	128.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	FAD	N3A-C2A-N1A	-10.13	121.14	128.89
2	B	801	FAD	C4-C4X-C10	-8.29	114.64	119.94
2	A	801	FAD	C4-C4X-C10	-6.55	115.75	119.94
2	A	801	FAD	C4X-C4-N3	-6.13	115.21	123.59
2	A	801	FAD	C4X-C10-N10	-6.08	116.94	120.52
2	B	801	FAD	C4X-C4-N3	-3.75	118.46	123.59
2	B	801	FAD	C4B-O4B-C1B	-3.45	105.92	109.72
2	B	801	FAD	C4X-C10-N10	-3.36	118.54	120.52
2	A	801	FAD	C4B-O4B-C1B	-3.35	106.03	109.72
2	A	801	FAD	O2'-C2'-C1'	-3.25	101.96	109.94
2	A	801	FAD	C7M-C7-C6	-2.10	114.58	120.28
2	A	801	FAD	O4'-C4'-C5'	2.00	114.55	110.19
2	B	801	FAD	O3B-C3B-C4B	2.10	117.34	111.05
2	A	801	FAD	O3'-C3'-C2'	2.75	115.68	108.75
2	A	801	FAD	O4B-C4B-C3B	2.76	110.71	105.15
2	B	801	FAD	O4B-C4B-C5B	2.85	119.53	109.32
2	A	801	FAD	O2A-PA-O3P	3.04	118.86	105.09
2	B	801	FAD	O2B-C2B-C3B	3.06	121.79	111.83
2	A	801	FAD	O4B-C1B-N9A	3.13	114.66	108.10
2	A	801	FAD	C5B-C4B-C3B	3.23	128.04	115.21
2	A	801	FAD	O3B-C3B-C4B	3.27	120.86	111.05
2	B	801	FAD	O4B-C4B-C3B	3.28	111.76	105.15
2	A	801	FAD	C1'-N10-C9A	3.60	122.90	118.86
2	A	801	FAD	C4-C4X-N5	3.61	123.10	118.72
2	B	801	FAD	O4B-C1B-N9A	3.61	115.66	108.10
2	B	801	FAD	C4X-N5-C5X	3.75	121.07	116.76
2	B	801	FAD	C1'-N10-C9A	4.06	123.42	118.86
2	B	801	FAD	C4-C4X-N5	4.74	124.47	118.72
2	B	801	FAD	C2B-C1B-N9A	6.65	124.44	114.29
2	A	801	FAD	C2B-C1B-N9A	8.46	127.21	114.29
2	B	801	FAD	C4-N3-C2	9.73	123.66	115.25
2	A	801	FAD	C4-N3-C2	13.10	126.57	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	FAD	1	0
2	B	801	FAD	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	575/623 (92%)	-0.15	13 (2%) 64 67	12, 17, 33, 55	0
1	B	576/623 (92%)	-0.11	14 (2%) 62 66	13, 19, 35, 58	0
All	All	1151/1246 (92%)	-0.13	27 (2%) 64 67	12, 18, 34, 58	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	459	VAL	7.7
1	A	459	VAL	7.5
1	A	45	ILE	7.1
1	A	343	ALA	6.0
1	B	343	ALA	6.0
1	B	344	ASN	5.9
1	A	458	ALA	5.0
1	A	44	ASP	5.0
1	B	458	ALA	4.5
1	A	460	GLN	3.9
1	B	45	ILE	3.9
1	A	618	PHE	3.9
1	B	457	GLY	3.7
1	B	232	GLY	3.7
1	B	618	PHE	3.6
1	B	460	GLN	3.6
1	A	457	GLY	3.5
1	A	344	ASN	3.4
1	B	617	PRO	3.4
1	B	341	ASN	3.1
1	A	341	ASN	2.5
1	B	342	PRO	2.5
1	A	342	PRO	2.5
1	B	44	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	617	PRO	2.2
1	B	345	PRO	2.2
1	A	233	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FAD	B	801	53/53	0.98	0.08	-0.82	10,14,18,19	0
2	FAD	A	801	53/53	0.98	0.08	-0.83	9,13,17,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.