



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:30 PM GMT

PDB ID : 4LST
Title : Crystal structure of broadly and potently neutralizing antibody VRC01 in complex with HIV-1 clade C strain ZM176.66 gp120
Authors : Zhou, T.; Moquin, S.; Kwong, P.D.
Deposited on : 2013-07-23
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4549 (2.58-2.50)
Clashscore	102246	5292 (2.58-2.50)
Ramachandran outliers	100387	5194 (2.58-2.50)
Sidechain outliers	100360	5196 (2.58-2.50)
RSRZ outliers	91569	4561 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	355	<div> <div>14%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>• 5%</div> </div> </div>
2	H	224	<div> <div>5%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div>••</div> </div> </div>
3	L	210	<div> <div>4%</div> <div> <div></div> <div>90%</div> <div>10%</div> </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6171 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENVELOPE GLYCOPROTEIN GP120 of HIV-1 clade C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	339	Total	C	N	O	S	0	0	0
			2657	1666	461	509	21			

- Molecule 2 is a protein called HEAVY CHAIN OF ANTIBODY VRC01.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	222	Total	C	N	O	S	0	0	0
			1699	1072	295	321	11			

- Molecule 3 is a protein called LIGHT CHAIN OF ANTIBODY VRC01.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	210	Total	C	N	O	S	0	0	0
			1632	1022	279	326	5			

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	G	25	Total O 25 25	0	0
5	H	27	Total O 27 27	0	0
5	L	19	Total O 19 19	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ENVELOPE GLYCOPROTEIN GP120 of HIV-1 clade C



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.11Å 77.98Å 200.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.68 – 2.55 40.68 – 2.55	Depositor EDS
% Data completeness (in resolution range)	91.9 (40.68-2.55) 91.9 (40.68-2.55)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.69 (at 2.54Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: DEV_998)	Depositor
R, R_{free}	0.190 , 0.239 0.189 , 0.233	Depositor DCC
R_{free} test set	1653 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	74.0	Xtriage
Anisotropy	0.527	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 79.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 32745 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6171	wwPDB-VP
Average B, all atoms (Å ²)	104.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	G	0.27	0/2714	0.46	0/3682
2	H	0.27	0/1743	0.50	0/2369
3	L	0.28	0/1669	0.47	0/2265
All	All	0.27	0/6126	0.47	0/8316

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2657	0	2585	26	0
2	H	1699	0	1671	12	0
3	L	1632	0	1574	11	0
4	G	112	0	104	3	0
5	G	25	0	0	0	0
5	H	27	0	0	1	0
5	L	19	0	0	0	0
All	All	6171	0	5934	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:61:ARG:NH1	5:H:306:HOH:O	2.22	0.73
3:L:47:VAL:HG12	3:L:48:ILE:HG12	1.75	0.68
3:L:110:ARG:NH1	3:L:111:THR:O	2.27	0.67
1:G:335:GLU:HG2	1:G:414:ILE:HG13	1.77	0.67
1:G:340:LYS:HG2	1:G:344:LYS:HD2	1.79	0.64
2:H:18:MET:HG2	2:H:82(C):LEU:HD11	1.80	0.64
1:G:46:LYS:HD2	1:G:490:GLU:O	1.97	0.64
2:H:119:PRO:HB3	2:H:145:TYR:HB3	1.80	0.62
2:H:51:LEU:HD21	2:H:71:ARG:HB3	1.82	0.61
1:G:231:GLU:HB2	1:G:268:GLU:HG3	1.83	0.60
1:G:295:VAL:HG22	1:G:446:LYS:HG2	1.88	0.55
1:G:294:ILE:HD12	1:G:333:ILE:HD11	1.90	0.53
2:H:148:GLU:HB2	2:H:149:PRO:HA	1.90	0.53
1:G:230:ASN:HB3	1:G:233:PHE:HB2	1.91	0.51
2:H:152:VAL:HG22	2:H:198:VAL:HG22	1.93	0.50
1:G:69:TRP:HA	1:G:72:HIS:CE1	2.47	0.49
1:G:117:LYS:HB3	1:G:203:GLN:HE21	1.78	0.48
3:L:60:ASP:N	3:L:60:ASP:OD1	2.47	0.48
1:G:339:ASN:O	1:G:343:GLN:HG3	2.14	0.47
1:G:233:PHE:HD1	4:G:503:NAG:H82	1.78	0.47
2:H:100:TYR:CE2	2:H:100(B):TRP:HB2	2.50	0.47
1:G:390:LEU:HD21	1:G:416:LEU:HD11	1.96	0.47
1:G:298:ARG:NH2	1:G:439:ILE:O	2.43	0.46
3:L:151:LYS:NZ	3:L:154:ASN:HA	2.30	0.46
1:G:59:LYS:HB2	1:G:62:GLU:HB2	1.98	0.46
2:H:34:LEU:HD23	2:H:78:ALA:HB2	1.99	0.45
1:G:378:CYS:HB3	1:G:383:PHE:CE1	2.52	0.45
3:L:107:ASP:OD2	3:L:142:TYR:OH	2.29	0.44
1:G:233:PHE:CD1	4:G:503:NAG:H82	2.52	0.44
2:H:38:ARG:HD3	2:H:90:TYR:CE2	2.53	0.44
3:L:35:TRP:CZ3	3:L:88:CYS:HB3	2.52	0.43
1:G:207:LYS:NZ	1:G:381:GLU:OE2	2.42	0.43
2:H:139:GLY:HA2	2:H:154:TRP:CH2	2.53	0.43
1:G:65:VAL:HB	1:G:115:SER:HB2	2.00	0.43
3:L:58:ILE:HA	3:L:59:PRO:HD3	1.82	0.43
1:G:45:TRP:O	1:G:46:LYS:HG3	2.18	0.42
3:L:4:LEU:HD12	3:L:25:THR:HB	2.01	0.42
2:H:87:THR:HG23	2:H:110:ILE:HA	2.01	0.42
3:L:151:LYS:HG2	3:L:156:LEU:HD23	2.00	0.42
1:G:52:LEU:HD11	1:G:100:MET:HG2	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:337:LYS:O	1:G:341:THR:HG23	2.20	0.42
1:G:121:LYS:HE3	1:G:426:MET:HE1	2.01	0.41
1:G:118:PRO:HD2	1:G:203:GLN:HE22	1.86	0.41
1:G:121:LYS:HB3	1:G:201:VAL:HG23	2.03	0.41
3:L:117:VAL:O	3:L:209:LYS:HE2	2.20	0.41
1:G:338:TRP:CZ2	1:G:390:LEU:HB3	2.56	0.41
2:H:29:PHE:CD2	2:H:76:ASP:HA	2.56	0.40
3:L:165:VAL:HG22	3:L:177:LEU:HD12	2.04	0.40
1:G:229:ASN:HB3	4:G:504:NAG:HN2	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	333/355 (94%)	319 (96%)	13 (4%)	1 (0%)	46	66
2	H	218/224 (97%)	208 (95%)	10 (5%)	0	100	100
3	L	208/210 (99%)	203 (98%)	4 (2%)	1 (0%)	34	54
All	All	759/789 (96%)	730 (96%)	27 (4%)	2 (0%)	46	66

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	412	THR
3	L	191	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	300/309 (97%)	296 (99%)	4 (1%)	76	91
2	H	191/193 (99%)	188 (98%)	3 (2%)	70	89
3	L	182/182 (100%)	182 (100%)	0	100	100
All	All	673/684 (98%)	666 (99%)	7 (1%)	82	94

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	45	TRP
1	G	116	LEU
1	G	230	ASN
1	G	492	LYS
2	H	34	LEU
2	H	57	VAL
2	H	71	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	72	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAG	G	501	1	14,14,15	0.44	0	15,19,21	1.37	2 (13%)
4	NAG	G	502	1	14,14,15	0.56	0	15,19,21	0.72	0
4	NAG	G	503	1	14,14,15	1.06	1 (7%)	15,19,21	1.48	2 (13%)
4	NAG	G	504	1	14,14,15	0.60	0	15,19,21	1.13	1 (6%)
4	NAG	G	505	1	14,14,15	0.44	0	15,19,21	0.80	0
4	NAG	G	506	1	14,14,15	0.49	0	15,19,21	0.82	1 (6%)
4	NAG	G	507	1	14,14,15	0.49	0	15,19,21	0.89	1 (6%)
4	NAG	G	508	1	14,14,15	0.54	0	15,19,21	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	501	1	-	0/6/23/26	0/1/1/1
4	NAG	G	502	1	-	0/6/23/26	0/1/1/1
4	NAG	G	503	1	-	0/6/23/26	0/1/1/1
4	NAG	G	504	1	-	0/6/23/26	0/1/1/1
4	NAG	G	505	1	-	0/6/23/26	0/1/1/1
4	NAG	G	506	1	-	0/6/23/26	0/1/1/1
4	NAG	G	507	1	-	0/6/23/26	0/1/1/1
4	NAG	G	508	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	503	NAG	O7-C7	-3.49	1.15	1.23

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	G	501	NAG	C2-N2-C7	-3.91	118.02	123.04
4	G	503	NAG	C4-C3-C2	-3.17	106.30	111.23
4	G	503	NAG	C3-C2-N2	-2.33	104.98	110.56
4	G	507	NAG	C1-O5-C5	2.33	115.20	112.25
4	G	506	NAG	C1-O5-C5	2.48	115.40	112.25
4	G	501	NAG	C1-O5-C5	2.60	115.55	112.25
4	G	504	NAG	C1-O5-C5	3.40	116.56	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	503	NAG	2	0
4	G	504	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	339/355 (95%)	0.92	49 (14%) 3 3	70, 99, 153, 185	0
2	H	222/224 (99%)	0.26	12 (5%) 29 34	67, 98, 148, 208	0
3	L	210/210 (100%)	0.14	8 (3%) 44 50	73, 102, 154, 199	0
All	All	771/789 (97%)	0.52	69 (8%) 12 13	67, 99, 154, 208	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	472	GLY	7.3
1	G	45	TRP	6.9
1	G	124	GLY	6.3
2	H	216	CYS	6.1
2	H	175	LEU	6.0
2	H	211	VAL	5.7
1	G	410	SER	5.6
3	L	216	CYS	5.4
1	G	492	LYS	5.1
1	G	473	GLY	5.0
3	L	1	GLU	4.7
1	G	471	GLY	4.4
1	G	440	LYS	4.3
3	L	2	ILE	3.6
1	G	491	ILE	3.6
1	G	283	THR	3.5
1	G	300	GLY	3.4
1	G	372	THR	3.3
1	G	284	ILE	3.3
2	H	136	ALA	3.2
1	G	481	SER	3.2
1	G	82	GLN	3.2
1	G	257	THR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	44	VAL	3.1
1	G	299	PRO	3.0
3	L	83	PHE	3.0
3	L	204	ARG	3.0
1	G	412	THR	3.0
2	H	54	GLY	3.0
1	G	46	LYS	3.0
1	G	411	ASN	2.9
1	G	455	THR	2.9
1	G	474	ASP	2.7
2	H	56	ALA	2.7
1	G	371	ILE	2.7
2	H	134	GLY	2.7
1	G	84	MET	2.7
2	H	174	GLY	2.7
1	G	285	ILE	2.7
1	G	256	SER	2.6
1	G	454	LEU	2.6
1	G	198	GLY	2.5
1	G	272	ILE	2.5
1	G	274	SER	2.5
2	H	133	GLY	2.5
1	G	470	PRO	2.5
1	G	86	LEU	2.5
3	L	182	THR	2.4
1	G	479	TRP	2.4
1	G	477	ASP	2.4
2	H	129	LYS	2.3
1	G	281	ALA	2.3
1	G	442	ASN	2.3
1	G	367	GLY	2.3
1	G	260	LEU	2.3
1	G	478	ASN	2.3
1	G	427	TRP	2.3
1	G	255	VAL	2.2
1	G	225	ILE	2.2
3	L	214	GLY	2.2
1	G	452	LEU	2.2
1	G	201	VAL	2.2
2	H	173	SER	2.2
1	G	271	VAL	2.1
1	G	286	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	365	SER	2.1
3	L	127	LEU	2.1
1	G	451	GLY	2.1
2	H	57	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NAG	G	508	14/15	0.92	0.30	1.48	144,160,167,168	0
4	NAG	G	502	14/15	0.94	0.18	0.64	107,139,155,170	0
4	NAG	G	505	14/15	0.93	0.19	0.40	92,115,138,140	0
4	NAG	G	503	14/15	0.79	0.17	-0.47	106,128,132,136	0
4	NAG	G	501	14/15	0.95	0.13	-0.79	66,94,107,107	0
4	NAG	G	507	14/15	0.92	0.23	-	139,160,163,164	0
4	NAG	G	504	14/15	0.84	0.31	-	130,148,161,161	0
4	NAG	G	506	14/15	0.82	0.42	-	154,173,184,191	0

6.5 Other polymers [i](#)

There are no such residues in this entry.