



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:27 AM GMT

PDB ID : 3LXZ
Title : Structure of probable Glutathione S-transferase(PP0183) from Pseudomonas putida
Authors : Ramagopal, U.A.; Toro, R.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2010-02-25
Resolution : 1.76 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1609 (1.76-1.76)
Clashscore	102246	1730 (1.76-1.76)
Ramachandran outliers	100387	1711 (1.76-1.76)
Sidechain outliers	100360	1711 (1.76-1.76)
RSRZ outliers	91569	1610 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	<div> <div>3%</div> <div> <div></div> <div>90%</div> <div>6%</div> <div>5%</div> </div> </div>
1	B	229	<div> <div>3%</div> <div> <div></div> <div>91%</div> <div>•</div> <div>5%</div> </div> </div>
1	C	229	<div> <div>4%</div> <div> <div></div> <div>86%</div> <div>9%</div> <div>5%</div> </div> </div>
1	D	229	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>9%</div> <div>5%</div> </div> </div>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 7538 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutathione S-transferase family protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	218	Total	C	N	O	S	Se	0	1	0
			1699	1101	272	316	2	8			
1	B	217	Total	C	N	O	S	Se	0	5	0
			1729	1120	279	320	2	8			
1	C	218	Total	C	N	O	S	Se	0	6	0
			1747	1127	286	324	2	8			
1	D	217	Total	C	N	O	S	Se	0	3	0
			1710	1105	279	316	2	8			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MSE	-	EXPRESSION TAG	UNP Q88RE7
A	1	SER	-	EXPRESSION TAG	UNP Q88RE7
A	221	GLU	-	EXPRESSION TAG	UNP Q88RE7
A	222	GLY	-	EXPRESSION TAG	UNP Q88RE7
A	223	HIS	-	EXPRESSION TAG	UNP Q88RE7
A	224	HIS	-	EXPRESSION TAG	UNP Q88RE7
A	225	HIS	-	EXPRESSION TAG	UNP Q88RE7
A	226	HIS	-	EXPRESSION TAG	UNP Q88RE7
A	227	HIS	-	EXPRESSION TAG	UNP Q88RE7
A	228	HIS	-	EXPRESSION TAG	UNP Q88RE7
B	0	MSE	-	EXPRESSION TAG	UNP Q88RE7
B	1	SER	-	EXPRESSION TAG	UNP Q88RE7
B	221	GLU	-	EXPRESSION TAG	UNP Q88RE7
B	222	GLY	-	EXPRESSION TAG	UNP Q88RE7
B	223	HIS	-	EXPRESSION TAG	UNP Q88RE7
B	224	HIS	-	EXPRESSION TAG	UNP Q88RE7
B	225	HIS	-	EXPRESSION TAG	UNP Q88RE7
B	226	HIS	-	EXPRESSION TAG	UNP Q88RE7
B	227	HIS	-	EXPRESSION TAG	UNP Q88RE7
B	228	HIS	-	EXPRESSION TAG	UNP Q88RE7
C	0	MSE	-	EXPRESSION TAG	UNP Q88RE7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	SER	-	EXPRESSION TAG	UNP Q88RE7
C	221	GLU	-	EXPRESSION TAG	UNP Q88RE7
C	222	GLY	-	EXPRESSION TAG	UNP Q88RE7
C	223	HIS	-	EXPRESSION TAG	UNP Q88RE7
C	224	HIS	-	EXPRESSION TAG	UNP Q88RE7
C	225	HIS	-	EXPRESSION TAG	UNP Q88RE7
C	226	HIS	-	EXPRESSION TAG	UNP Q88RE7
C	227	HIS	-	EXPRESSION TAG	UNP Q88RE7
C	228	HIS	-	EXPRESSION TAG	UNP Q88RE7
D	0	MSE	-	EXPRESSION TAG	UNP Q88RE7
D	1	SER	-	EXPRESSION TAG	UNP Q88RE7
D	221	GLU	-	EXPRESSION TAG	UNP Q88RE7
D	222	GLY	-	EXPRESSION TAG	UNP Q88RE7
D	223	HIS	-	EXPRESSION TAG	UNP Q88RE7
D	224	HIS	-	EXPRESSION TAG	UNP Q88RE7
D	225	HIS	-	EXPRESSION TAG	UNP Q88RE7
D	226	HIS	-	EXPRESSION TAG	UNP Q88RE7
D	227	HIS	-	EXPRESSION TAG	UNP Q88RE7
D	228	HIS	-	EXPRESSION TAG	UNP Q88RE7

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	148	Total O 148 148	0	0
2	B	181	Total O 181 181	0	0
2	C	170	Total O 170 170	0	0
2	D	154	Total O 154 154	0	0

- Molecule 1: Glutathione S-transferase family protein



- [illegible]

- [illegible]

-

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	48.05Å 66.36Å 88.66Å 89.96° 85.27° 87.04°	Depositor
Resolution (Å)	50.00 – 1.76 30.58 – 1.76	Depositor EDS
% Data completeness (in resolution range)	90.6 (50.00-1.76) 90.5 (30.58-1.76)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.98 (at 1.76Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.196 , 0.229 0.206 , 0.237	Depositor DCC
R_{free} test set	4901 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.6	EDS
Estimated twinning fraction	0.064 for -h,k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 97746 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7538	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	1/1726 (0.1%)	0.73	2/2318 (0.1%)
1	B	0.74	1/1755 (0.1%)	0.72	0/2355
1	C	0.75	0/1773	0.75	0/2377
1	D	0.75	1/1735 (0.1%)	0.72	0/2328
All	All	0.74	3/6989 (0.0%)	0.73	2/9378 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	205	GLU	CG-CD	5.22	1.59	1.51
1	A	159	CYS	CB-SG	-5.11	1.73	1.81
1	B	13	TYR	CD1-CE1	5.08	1.47	1.39

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	141	ARG	NE-CZ-NH2	-5.60	117.50	120.30
1	A	138	ARG	NE-CZ-NH1	5.26	122.93	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1699	0	1717	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1729	0	1755	6	0
1	C	1747	0	1767	22	0
1	D	1710	0	1737	12	0
2	A	148	0	0	1	0
2	B	181	0	0	3	0
2	C	170	0	0	2	0
2	D	154	0	0	6	0
All	All	7538	0	6976	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 45 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:208:MSE:SE	1:C:212:MSE:HE3	2.21	0.90
1:C:211:PHE:CD2	1:C:212:MSE:HE2	2.08	0.87
1:C:211:PHE:CD2	1:C:212:MSE:CE	2.61	0.83
1:C:211:PHE:HD2	1:C:212:MSE:CE	2.02	0.71
1:C:47:GLY:HA2	2:C:245:HOH:O	1.93	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/229 (95%)	213 (98%)	4 (2%)	0	100	100
1	B	220/229 (96%)	215 (98%)	5 (2%)	0	100	100
1	C	222/229 (97%)	219 (99%)	3 (1%)	0	100	100
1	D	218/229 (95%)	214 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	877/916 (96%)	861 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/180 (99%)	175 (98%)	4 (2%)	60	35
1	B	184/180 (102%)	179 (97%)	5 (3%)	52	27
1	C	185/180 (103%)	182 (98%)	3 (2%)	70	52
1	D	181/180 (101%)	177 (98%)	4 (2%)	60	35
All	All	729/720 (101%)	713 (98%)	16 (2%)	63	35

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	171	LYS
1	B	181	PHE
1	D	120	LEU
1	B	117[B]	VAL
1	D	171	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	39	GLN
1	C	166	ASN
1	C	189	GLN
1	D	189	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/229 (91%)	0.01	6 (2%) 55 61	14, 21, 33, 47	0
1	B	209/229 (91%)	-0.07	6 (2%) 55 61	13, 18, 32, 47	0
1	C	210/229 (91%)	-0.04	10 (4%) 34 39	11, 18, 30, 49	0
1	D	209/229 (91%)	-0.01	6 (2%) 55 61	13, 19, 33, 55	0
All	All	838/916 (91%)	-0.03	28 (3%) 50 56	11, 19, 33, 55	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	217	SER	4.6
1	A	1	SER	3.9
1	B	1	SER	3.9
1	B	217	SER	3.6
1	A	215	ILE	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.