



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:43 PM GMT

PDB ID : 4MJR
Title : E. coli sliding clamp in complex with (S)-Carprofen
Authors : Yin, Z.; Oakley, A.J.
Deposited on : 2013-09-04
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3202 (1.64-1.60)
Clashscore	102246	3500 (1.64-1.60)
Ramachandran outliers	100387	3411 (1.64-1.60)
Sidechain outliers	100360	3410 (1.64-1.60)
RSRZ outliers	91569	3207 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	<div> <div>2%</div> <div>87%</div> <div>11%</div> </div>
1	B	366	<div> <div>5%</div> <div>91%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PGE	A	401	-	-	-	X
3	PEG	A	402	-	-	-	X
6	CA	B	405	-	-	-	X

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 6404 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase III subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	12	0
			2839	1793	488	536	22			
1	B	365	Total	C	N	O	S	0	14	0
			2884	1824	497	543	20			

- Molecule 2 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	6	4		
2	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Cl	0	0
			2	2		
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is (2S)-2-(6-CHLORO-9H-CARBAZOL-2-YL)PROPANOIC ACID (three-letter code: 0LA) (formula: C₁₅H₁₂ClNO₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Cl	N	O	0	0
			19	15	1	1	2		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Ca	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	310	Total	O	0	0
			310	310		
7	B	300	Total	O	0	0
			300	300		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase III subunit beta



- Molecule 1: DNA polymerase III subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	40.89Å 64.19Å 71.88Å 74.01° 82.99° 84.03°	Depositor
Resolution (Å)	21.27 – 1.62 21.27 – 1.62	Depositor EDS
% Data completeness (in resolution range)	95.1 (21.27-1.62) 86.4 (21.27-1.62)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.88 (at 1.62Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.189 , 0.225 0.197 , 0.231	Depositor DCC
R_{free} test set	4235 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	16.2	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 45.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 84106 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6404	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CA, PGE, 0LA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/2922	0.67	0/3954
1	B	0.44	0/2977	0.67	0/4033
All	All	0.43	0/5899	0.67	0/7987

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2839	0	2869	28	0
1	B	2884	0	2920	38	1
2	A	10	0	14	0	0
2	B	10	0	14	0	0
3	A	7	0	10	0	0
3	B	21	0	30	3	0
4	A	1	0	0	0	0
4	B	2	0	0	0	0
5	A	19	0	11	6	0
6	B	1	0	0	0	0
7	A	310	0	0	7	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	300	0	0	3	0
All	All	6404	0	5868	65	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:346[B]:SER:CB	1:B:362[B]:MET:CE	1.86	1.52
1:B:346[B]:SER:HB3	1:B:362[B]:MET:CE	0.90	1.36
1:B:346[B]:SER:HB3	1:B:362[B]:MET:HE3	1.26	1.14
1:B:346[B]:SER:CB	1:B:362[B]:MET:HE1	1.63	1.10
1:B:346[B]:SER:HB3	1:B:362[B]:MET:HE2	1.10	1.05
1:B:362[B]:MET:HA	1:B:362[B]:MET:HE2	1.49	0.93
1:B:346[B]:SER:HB3	1:B:362[B]:MET:HE1	0.88	0.88
1:A:152:ARG:HH11	5:A:404:OLA:H10	1.42	0.85
1:B:346[B]:SER:CB	1:B:362[B]:MET:HE3	1.92	0.81
1:B:346[B]:SER:CB	1:B:362[B]:MET:HE2	1.81	0.80
1:A:184:ILE:HD11	1:A:188:LEU:HD11	1.62	0.80
1:B:362[B]:MET:CE	1:B:362[B]:MET:HA	2.19	0.73
1:A:205:ARG:HG3	7:A:725:HOH:O	1.93	0.68
1:B:214:LEU:HD11	1:B:225:ALA:HB1	1.79	0.65
1:A:136:LYS:NZ	7:A:760:HOH:O	2.27	0.65
1:B:150:ASP:OD1	1:B:152:ARG:HD3	1.97	0.65
1:B:346[B]:SER:CA	1:B:362[B]:MET:HE2	2.30	0.60
1:A:239:GLY:N	7:A:768:HOH:O	2.34	0.59
1:B:26[B]:THR:HG21	7:B:575:HOH:O	2.03	0.58
1:B:366:LEU:HD12	1:B:366:LEU:O	2.04	0.57
1:B:120:ASP:OD2	3:B:402:PEG:O2	2.23	0.56
1:A:237:VAL:HG12	7:A:768:HOH:O	2.05	0.55
1:B:136:LYS:HG3	1:B:204:MET:HE1	1.91	0.53
1:A:93:GLU:HG3	1:A:98:LEU:HD22	1.92	0.52
1:B:346[B]:SER:OG	1:B:362[B]:MET:HE1	2.06	0.51
1:A:362:MET:HG2	5:A:404:OLA:H1	1.92	0.51
1:A:152:ARG:NH1	5:A:404:OLA:H10	2.19	0.51
1:B:73:ARG:HG3	7:B:665:HOH:O	2.09	0.51
1:B:120:ASP:CG	3:B:402:PEG:O2	2.48	0.51
1:B:249:PRO:HD2	1:B:348:GLN:HE21	1.76	0.51
1:A:6:GLU:HG3	7:A:776:HOH:O	2.11	0.50
1:B:258:ALA:HB2	1:B:308:VAL:HG12	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:ASP:C	7:A:768:HOH:O	2.51	0.49
1:A:152:ARG:HD2	5:A:404:OLA:C14	2.43	0.49
1:A:307:ASP:OD1	1:B:103:ARG:NH2	2.42	0.48
1:A:184:ILE:CD1	1:A:188:LEU:HD11	2.39	0.48
1:B:258:ALA:HB3	1:B:263:LEU:HD11	1.94	0.48
1:A:348:GLN:NE2	1:A:350:GLU:OE2	2.43	0.48
1:B:263:LEU:HD12	1:B:336:VAL:CG1	2.42	0.48
1:B:214:LEU:HD13	1:B:227:VAL:HG22	1.95	0.47
1:A:19:GLY:HA3	7:A:778:HOH:O	2.14	0.47
1:B:214:LEU:CD1	1:B:227:VAL:HG22	2.44	0.47
1:B:33:LEU:HG	1:B:72:ALA:HB2	1.97	0.46
1:B:197[B]:ARG:HB2	1:B:197[B]:ARG:HH11	1.81	0.45
1:A:144:PHE:CD2	1:A:326:ASP:HB3	2.51	0.45
1:A:254:LYS:HD3	1:A:315[B]:MET:HE2	1.99	0.45
1:A:346:SER:HB3	1:A:362:MET:SD	2.57	0.45
1:B:346[B]:SER:CA	1:B:362[B]:MET:CE	2.82	0.43
1:B:177:LEU:HD13	1:B:247:VAL:HG21	2.00	0.43
1:B:146:MET:HE1	1:B:197[B]:ARG:HA	2.01	0.42
1:B:197[B]:ARG:NH1	1:B:197[B]:ARG:HB2	2.34	0.42
1:A:296:ASN:HB2	1:A:297:PRO:CD	2.49	0.42
1:B:120:ASP:OD2	3:B:402:PEG:O1	2.31	0.42
1:A:12:LYS:HB3	1:A:13:PRO:HD3	2.01	0.42
1:B:146:MET:HE1	1:B:197[A]:ARG:HA	2.02	0.41
1:A:51:MET:HE1	1:A:198:LYS:HG2	2.02	0.41
1:B:251:ASN:ND2	7:B:700:HOH:O	2.53	0.41
1:A:271:ALA:HB2	1:A:321:VAL:HG21	2.01	0.41
1:A:152:ARG:HH11	5:A:404:OLA:C14	2.23	0.41
1:A:261:ASP:O	1:A:265:GLN:HG2	2.21	0.41
1:A:140:GLU:CG	1:A:204[A]:MET:HE2	2.51	0.41
1:A:152:ARG:HD2	5:A:404:OLA:H11	2.03	0.41
1:B:147:ALA:O	1:B:197[B]:ARG:NH1	2.52	0.41
1:B:214:LEU:HD13	1:B:227:VAL:CG2	2.51	0.40
1:A:136:LYS:HE3	1:A:204[A]:MET:HE1	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:154:TYR:OH	7:A:726:HOH:O[1_556]	2.12	0.08

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	368/366 (100%)	362 (98%)	6 (2%)	0	100	100
1	B	375/366 (102%)	366 (98%)	9 (2%)	0	100	100
All	All	743/732 (102%)	728 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/313 (100%)	307 (98%)	7 (2%)	60	30
1	B	321/313 (103%)	314 (98%)	7 (2%)	60	30
All	All	635/626 (101%)	621 (98%)	14 (2%)	58	30

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	52	GLU
1	A	71	PRO
1	A	95	GLU
1	A	120	ASP
1	A	143	GLN
1	A	176	ARG
1	B	176	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	184	ILE
1	B	192	SER
1	B	247	VAL
1	B	251	ASN
1	B	298	GLU
1	B	366	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	HIS
1	A	36	GLN
1	A	91	GLN
1	A	143	GLN
1	A	191	HIS
1	B	191	HIS
1	B	212	ASN
1	B	251	ASN
1	B	295	ASN
1	B	348	GLN
1	B	355	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PGE	A	401	-	9,9,9	0.61	0	8,8,8	0.53	0
3	PEG	A	402	-	6,6,6	0.46	0	5,5,5	0.18	0
5	OLA	A	404	-	18,21,21	1.53	3 (16%)	25,31,31	1.70	9 (36%)
3	PEG	B	401	-	6,6,6	0.45	0	5,5,5	0.21	0
3	PEG	B	402	-	6,6,6	0.49	0	5,5,5	0.61	0
2	PGE	B	403	-	9,9,9	0.48	0	8,8,8	0.60	0
3	PEG	B	404	-	6,6,6	0.52	0	5,5,5	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PGE	A	401	-	-	0/7/7/7	0/0/0/0
3	PEG	A	402	-	-	0/4/4/4	0/0/0/0
5	OLA	A	404	-	-	0/4/8/8	0/3/3/3
3	PEG	B	401	-	-	0/4/4/4	0/0/0/0
3	PEG	B	402	-	-	0/4/4/4	0/0/0/0
2	PGE	B	403	-	-	0/7/7/7	0/0/0/0
3	PEG	B	404	-	-	0/4/4/4	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	404	OLA	C4B-C8A	2.07	1.48	1.42
5	A	404	OLA	C4A-C9A	2.08	1.48	1.42
5	A	404	OLA	C5-C6	2.30	1.41	1.36

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	404	OLA	C2-C3-CL	-3.17	114.15	119.35
5	A	404	OLA	C6-C5-C4B	-2.76	117.31	121.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	404	0LA	C8-C7-C10	-2.13	115.83	121.80
5	A	404	0LA	C11-C10-C7	-2.11	108.18	112.32
5	A	404	0LA	C2-C1-C9A	-2.07	118.63	120.88
5	A	404	0LA	C8-C8A-C4B	-2.05	118.52	121.14
5	A	404	0LA	C1-C2-C3	2.19	122.00	119.28
5	A	404	0LA	C4-C3-CL	2.55	123.36	119.74
5	A	404	0LA	C4-C4A-C9A	3.38	121.68	120.34

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	404	0LA	6	0
3	B	402	PEG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	360/366 (98%)	0.03	8 (2%) 65 63	9, 17, 30, 41	3 (0%)
1	B	365/366 (99%)	0.14	18 (4%) 33 29	9, 19, 35, 51	2 (0%)
All	All	725/732 (99%)	0.09	26 (3%) 46 42	9, 18, 33, 51	5 (0%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	366	LEU	6.7
1	A	19	GLY	5.7
1	A	20	PRO	4.9
1	A	27	LEU	4.0
1	B	211	ASP	3.8
1	B	23	GLY	3.7
1	B	252	PRO	3.3
1	B	25	PRO	3.3
1	B	209	GLY	3.2
1	B	24	ARG	3.2
1	B	210	GLY	3.0
1	B	251	ASN	2.9
1	A	28	PRO	2.7
1	A	153	TYR	2.5
1	B	153	TYR	2.5
1	B	208	ASP	2.5
1	B	284[A]	TYR	2.4
1	B	253	ASP	2.4
1	B	148	HIS	2.4
1	A	120	ASP	2.3
1	B	212	ASN	2.3
1	A	238	ASP	2.2
1	B	332	LYS	2.1
1	B	297	PRO	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	187	SER	2.0
1	B	278	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	PEG	A	402	7/7	0.84	0.29	4.11	31,33,34,35	0
6	CA	B	405	1/1	0.85	0.21	4.02	56,56,56,56	0
2	PGE	A	401	10/10	0.67	0.24	2.87	26,30,33,33	0
5	OLA	A	404	19/19	0.91	0.15	1.81	17,24,34,35	19
3	PEG	B	404	7/7	0.68	0.19	1.71	42,43,44,44	0
3	PEG	B	402	7/7	0.74	0.21	1.65	26,30,33,33	0
2	PGE	B	403	10/10	0.81	0.15	1.02	37,40,42,44	0
4	CL	B	406	1/1	0.99	0.12	-0.01	23,23,23,23	0
3	PEG	B	401	7/7	0.90	0.09	-0.22	29,30,34,35	0
4	CL	A	403	1/1	1.00	0.02	-2.05	22,22,22,22	0
4	CL	B	407	1/1	0.98	0.03	-2.09	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.