



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 08:55 PM GMT

PDB ID : 1MQB
Title : Crystal Structure of Ephrin A2 (ephA2) Receptor Protein Kinase
Authors : Nowakowski, J.; Cronin, C.N.; McRee, D.E.; Knuth, M.W.; Nelson, C.; Pavletich, N.; Rogers, J.; Sang, B.C.; Scheibe, D.N.; Swanson, R.V.; Thompson, D.A.
Deposited on : 2002-09-16
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	333	
1	B	333	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4349 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ephrin type-A receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	0	0
			2101	1349	360	374	18			
1	B	265	Total	C	N	O	S	0	0	0
			2120	1362	365	375	18			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	568	MET	-	CLONING ARTIFACT	UNP P29317
A	569	SER	-	CLONING ARTIFACT	UNP P29317
A	570	TYR	-	CLONING ARTIFACT	UNP P29317
A	571	TYR	-	CLONING ARTIFACT	UNP P29317
A	572	HIS	-	CLONING ARTIFACT	UNP P29317
A	573	HIS	-	CLONING ARTIFACT	UNP P29317
A	574	HIS	-	CLONING ARTIFACT	UNP P29317
A	575	HIS	-	CLONING ARTIFACT	UNP P29317
A	576	HIS	-	CLONING ARTIFACT	UNP P29317
A	577	HIS	-	CLONING ARTIFACT	UNP P29317
A	578	ASP	-	CLONING ARTIFACT	UNP P29317
A	579	TYR	-	CLONING ARTIFACT	UNP P29317
A	580	ASP	-	CLONING ARTIFACT	UNP P29317
A	581	ILE	-	CLONING ARTIFACT	UNP P29317
A	582	PRO	-	CLONING ARTIFACT	UNP P29317
A	583	THR	-	CLONING ARTIFACT	UNP P29317
A	584	THR	-	CLONING ARTIFACT	UNP P29317
A	585	GLU	-	CLONING ARTIFACT	UNP P29317
A	586	ASN	-	CLONING ARTIFACT	UNP P29317
A	587	LEU	-	CLONING ARTIFACT	UNP P29317
A	588	TYR	-	CLONING ARTIFACT	UNP P29317
A	589	PHE	-	CLONING ARTIFACT	UNP P29317
A	590	GLN	-	CLONING ARTIFACT	UNP P29317
A	591	GLY	-	CLONING ARTIFACT	UNP P29317
A	592	ALA	-	CLONING ARTIFACT	UNP P29317

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	593	MET	-	CLONING ARTIFACT	UNP P29317
A	594	GLY	-	CLONING ARTIFACT	UNP P29317
A	595	SER	-	CLONING ARTIFACT	UNP P29317
B	568	MET	-	CLONING ARTIFACT	UNP P29317
B	569	SER	-	CLONING ARTIFACT	UNP P29317
B	570	TYR	-	CLONING ARTIFACT	UNP P29317
B	571	TYR	-	CLONING ARTIFACT	UNP P29317
B	572	HIS	-	CLONING ARTIFACT	UNP P29317
B	573	HIS	-	CLONING ARTIFACT	UNP P29317
B	574	HIS	-	CLONING ARTIFACT	UNP P29317
B	575	HIS	-	CLONING ARTIFACT	UNP P29317
B	576	HIS	-	CLONING ARTIFACT	UNP P29317
B	577	HIS	-	CLONING ARTIFACT	UNP P29317
B	578	ASP	-	CLONING ARTIFACT	UNP P29317
B	579	TYR	-	CLONING ARTIFACT	UNP P29317
B	580	ASP	-	CLONING ARTIFACT	UNP P29317
B	581	ILE	-	CLONING ARTIFACT	UNP P29317
B	582	PRO	-	CLONING ARTIFACT	UNP P29317
B	583	THR	-	CLONING ARTIFACT	UNP P29317
B	584	THR	-	CLONING ARTIFACT	UNP P29317
B	585	GLU	-	CLONING ARTIFACT	UNP P29317
B	586	ASN	-	CLONING ARTIFACT	UNP P29317
B	587	LEU	-	CLONING ARTIFACT	UNP P29317
B	588	TYR	-	CLONING ARTIFACT	UNP P29317
B	589	PHE	-	CLONING ARTIFACT	UNP P29317
B	590	GLN	-	CLONING ARTIFACT	UNP P29317
B	591	GLY	-	CLONING ARTIFACT	UNP P29317
B	592	ALA	-	CLONING ARTIFACT	UNP P29317
B	593	MET	-	CLONING ARTIFACT	UNP P29317
B	594	GLY	-	CLONING ARTIFACT	UNP P29317
B	595	SER	-	CLONING ARTIFACT	UNP P29317

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
2	B	1	Total 31	C 10	N 6	O 12	P 3	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	39	Total O 39 39	0	0
3	B	27	Total O 27 27	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Ephrin type-A receptor 2



- Molecule 1: Ephrin type-A receptor 2



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	72.13 Å 72.13 Å 241.62 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 2.30	Depositor
% Data completeness (in resolution range)	93.7 (40.00-2.30)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.236 , 0.288	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4349	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.78	0/2145	0.89	5/2888 (0.2%)
1	B	0.78	0/2164	0.94	12/2913 (0.4%)
All	All	0.78	0/4309	0.92	17/5801 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	708	ASP	CB-CG-OD2	8.65	126.09	118.30
1	B	757	ASP	CB-CG-OD2	7.21	124.78	118.30
1	B	687	PRO	N-CD-CG	-6.79	93.02	103.20
1	A	657	ARG	NE-CZ-NH2	-6.15	117.23	120.30
1	B	687	PRO	N-CA-CB	-6.04	95.96	102.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	686	LYS	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	654	GLU	Peptide
1	B	686	LYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2101	0	2133	26	0
1	B	2120	0	2162	30	0
2	A	31	0	13	1	0
2	B	31	0	13	2	0
3	A	39	0	0	0	0
3	B	27	0	0	0	0
All	All	4349	0	4321	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 57 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:807:MET:HE2	1:A:850:MET:HB3	1.40	0.98
1:A:807:MET:HE2	1:A:850:MET:CB	2.06	0.84
1:A:748:ASN:HD22	1:A:748:ASN:C	1.86	0.76
1:B:795:THR:HG22	1:B:798:SER:H	1.51	0.75
1:B:808:TRP:O	1:B:812:THR:HB	1.93	0.68

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/333 (77%)	248 (96%)	8 (3%)	1 (0%)	39	48
1	B	259/333 (78%)	244 (94%)	13 (5%)	2 (1%)	24	27
All	All	516/666 (78%)	492 (95%)	21 (4%)	3 (1%)	30	36

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	687	PRO
1	B	687	PRO
1	B	653	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/287 (79%)	199 (88%)	27 (12%)	6	7
1	B	229/287 (80%)	190 (83%)	39 (17%)	2	2
All	All	455/574 (79%)	389 (86%)	66 (14%)	4	3

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	614	THR
1	B	648	LEU
1	B	863	LYS
1	B	617	LYS
1	B	633	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	732	ASN
1	B	852	GLN
1	A	748	ASN
1	A	674	ASN
1	B	674	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ANP	A	1000	-	27,33,33	1.46	4 (14%)	30,52,52	2.87	6 (20%)
2	ANP	B	1001	-	27,33,33	1.50	5 (18%)	30,52,52	2.62	7 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	1000	-	-	1/12/38/38	0/3/3/3
2	ANP	B	1001	-	-	0/12/38/38	0/3/3/3

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	ANP	C2-N1	2.13	1.38	1.33
2	B	1001	ANP	C2-N1	2.34	1.38	1.33
2	B	1001	ANP	PB-O1B	2.65	1.49	1.46
2	A	1000	ANP	C2-N3	2.77	1.37	1.32
2	A	1000	ANP	PB-O1B	2.98	1.49	1.46

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	ANP	N3-C2-N1	-13.31	118.70	128.89
2	B	1001	ANP	N3-C2-N1	-11.83	119.84	128.89
2	A	1000	ANP	O1B-PB-N3B	-4.09	105.62	111.90
2	B	1001	ANP	C2'-C1'-N9	-3.83	108.44	114.29
2	A	1000	ANP	C4-C5-N7	-3.02	106.70	109.48

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1000	ANP	O1G-PG-N3B-PB

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1000	ANP	1	0
2	B	1001	ANP	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.